UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under All Rights Reserved license:

Loisel, J., Yu, Z., Beilman, D. W., Camill, P., Alm, J., Amesbury, M. J., Anderson, D., Andersson, S., Bochicchio, C., Barber, K., Belyea, L. R., Bunbury, J., Chambers, Frank M ORCID: 0000-0002-0998-2093, Charman, D. J., De Vleeschouwer, F., Fia kiewicz-Kozie, B., Finkelstein, S. A., Ga ka, M., Garneau, M., Hammarlund, D., Hinchcliffe, W., Holmquist, J., Hughes, P., Jones, M. C., Klein, E. S., Kokfelt, U., Korhola, A., Kuhry, P., Lamarre, A., Lamentowicz, M., Large, D., Lavoie, M., MacDonald, G., Magnan, G., Makila, M., Mallon, G., Mathijssen, P., Mauquoy, D., McCarroll, Julia, Moore, T. R., Nichols, J., O'Reilly, B., Oksanen, P., Packalen, M., Peteet, D., Richard, P. J., Robinson, S., Ronkainen, T., Rundgren, M., Sannel, A. B. K., Tarnocai, C., Thom, T., Tuittila, E.-S., Turetsky, M., Valiranta, M., van der Linden, M., van Geel, B., van Bellen, S., Vitt, D., Zhao, Y. and Zhou, W. (2014) A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation. *Holocene*, 24 (9). pp. 1028-1042. doi:10.1177/0959683614538073

Official URL: <http://dx.doi.org/10.1177/0959683614538073>

DOI: <http://dx.doi.org/10.1177/0959683614538073>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/2504>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

This is a for peer-review, pre-print version of the following published document:

Loisel, J. and Yu, Z. and Beilman, D. W. and Camill, P. and Alm, J. and Amesbury, M. J. and Anderson, D. and Andersson, S. and Bochicchio, C. and Barber, K. and Belyea, L. R. and Bunbury, J. and Chambers, Frank M and Charman, D. J. and De Vleeschouwer, F. and Fia kiewicz-Kozie, B. and Finkelstein, S. A. and Ga ka, M. and Garneau, M. and Hammarlund, D. and Hinchcliffe, W. and Holmquist, J. and Hughes, P. and Jones, M. C. and Klein, E. S. and Kokfelt, U. and Korhola, A. and Kuhry, P. and Lamarre, A. and Lamentowicz, M. and Large, D. and Lavoie, M. and MacDonald, G. and Magnan, G. and Makila, M. and Mallon, G. and Mathijssen, P. and Mauquoy, D. and McCarroll, J. and Moore, T. R. and Nichols, J. and O'Reilly, B. and Oksanen, P. and Packalen, M. and Peteet, D. and Richard, P. J. and Robinson, S. and Ronkainen, T. and Rundgren, M. and Sannel, A. B. K. and Tarnocai, C. and Thom, T. and Tuittila, E.-S. and Turetsky, M. and Valiranta, M. and van der Linden, M. and van Geel, B. and van Bellen, S. and Vitt, D. and Zhao, Y. and Zhou, W. (2014). *A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation*. *The Holocene*, 24 (9), 1028-1042. ISSN 0959-6836

Published in *The Holocene*, and available online at:

<http://hol.sagepub.com/content/24/9/1028>

We recommend you cite the published (post-print) version.

The URL for the published version is <http://dx.doi.org/10.1177/0959683614538073>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation

Journal:	<i>The Holocene</i>
Manuscript ID:	HOL-13-0128.R1
Manuscript Type:	Paper
Date Submitted by the Author:	20-Dec-2013
Complete List of Authors:	Loisel, Julie; Lehigh University, Earth and Environmental Sciences Yu, Zicheng; Lehigh University, Department of Earth and Environmental Sciences; Beilman, David; University of Hawaii at Manoa, Camill, Philip; Bowdoin, Environmental Studies Program and Department of Biology Carbon Network, Holocene; Lehigh University, Earth and Environmental Sciences
Keywords:	Northern peatlands, Data synthesis, Climate change, Soil carbon and nitrogen, Biogeochemical cycles, Long-term ecosystem dynamics
Abstract:	<p>Here we present results from the most comprehensive compilation of Holocene peat soil properties with associated carbon and nitrogen accumulation rates for northern peatlands. Our database consists of 268 peat cores from 215 sites located north of 45°N. It encompasses regions within which peat carbon data have only recently become available, such as the West Siberia Lowlands, the Hudson Bay Lowlands, Kamchatka in Far East Russia, and the Tibetan Plateau. For all northern peatlands, carbon content in organic matter was estimated at $42 \pm 3\%$ (S.D.) for Sphagnum peat, $51 \pm 2\%$ for non-Sphagnum peat, and at $49 \pm 2\%$ overall. Dry bulk density averaged 0.12 ± 0.07 g cm⁻³, organic matter bulk density averaged 0.11 ± 0.05 g cm⁻³, and total carbon content in peat averaged $47 \pm 6\%$. In general, large differences were found between Sphagnum and non-Sphagnum peat types in terms of peat properties. Time-weighted peat carbon accumulation rates averaged 23 ± 2 (S.E.M.) g C m⁻² yr⁻¹ during the Holocene on the basis of 151 peat cores from 127 sites, with the highest rates of carbon accumulation ($25\text{--}28$ g C m⁻² yr⁻¹) recorded during the early Holocene when the climate was warmer than the present. Furthermore, we estimate the northern peatland carbon and nitrogen pools at 436 and 10 gigatons, respectively. The database is publicly available at https://peatlands.lehigh.edu.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For Peer Review

A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation

Running title

Northern peatland database and synthesis

Authors

Julie Loisel^{1*}, Zicheng Yu^{1*}, David W. Beilman², Philip Camill³, Jukka Alm⁴, Matthew J. Amesbury⁵, David Anderson⁶, Sofia Andersson⁷, Christopher Bochicchio¹, Keith Barber⁸, Lisa R. Belyea⁹, Joan Bunbury¹⁰, Frank M. Chambers¹¹, Daniel J. Charman⁵, François De Vleeschouwer¹², Barbara Fiałkiewicz-Kozieł¹³, Sarah A. Finkelstein¹⁴, Mariusz Gałka¹³, Michelle Garneau¹⁵, Dan Hammarlund¹⁶, William Hinchcliffe⁵, James Holmquist¹⁷, Paul Hughes⁸, Miriam C. Jones¹⁸, Eric S. Klein¹, Ulla Kokfelt¹⁹, Atte Korhola²⁰, Peter Kuhry⁷, Alexandre Lamarre¹⁵, Mariusz Lamentowicz¹³, David Large²¹, Martin Lavoie²², Glen MacDonald¹⁷, Gabriel Magnan¹⁵, Markku Mäkilä²³, Gunnar Mallon⁸, Paul Mathijssen²⁰, Dmitri Mauquoy²⁴, Julia McCarroll¹¹, Tim R. Moore²⁵, Jonathan Nichols²⁶, Benjamin O'Reilly¹⁴, Pirita Oksanen²⁷, Maara Packalen²⁸, Dorothy Peteet²⁶, Pierre J.H. Richard²⁹, Stephen Robinson³⁰, Tiina Ronkainen²⁰, Mats Rundgren¹⁶, A. Britta K. Sannel⁷, Charles Tarnocai³¹, Tim Thom³², Eeva-Stiina Tuittila⁴, Merritt Turetsky³³, Minna Väliranta²⁰, Marjolein van der Linden³⁴, Bas van Geel³⁵, Simon van Bellen²³, Dale Vitt³⁶, Yan Zhao³⁷, Weijian Zhou³⁸

Revised manuscript submitted on 19 December 2013 as a Research Paper for the special issue *Holocene Peatland Carbon Dynamics in the Circum-Arctic Region*.

Affiliations

¹Department of Earth and Environmental Sciences, Lehigh University, Bethlehem, PA 18015, USA

²Department of Geography, University of Hawaii – Manoa, Honolulu, HI 96822, USA

³Department of Earth and Oceanographic Sciences, Bowdoin College, Brunswick, ME 04011, USA

⁴School of Forest Sciences, University of Eastern Finland, Joensuu, FI 80101, Finland

⁵Department of Geography, University of Exeter, Exeter, EX4 4RJ, UK

⁶Department of Geography, Eton College, Windsor, Berkshire SL4 6DW, UK

⁷Department of Physical Geography and Quaternary Geology, Stockholm University, Stockholm, 106 91, Sweden

⁸Geography and Environment, University of Southampton, Southampton, SO17 1BJ, UK

⁹School of Geography, Queen Mary University of London, London, E1 4NS, UK

¹⁰Department of Geography and Earth Science, University of Wisconsin – La Crosse, La Crosse, WI 54601, USA

¹¹Centre for Environmental Change and Quaternary Research, University of Gloucestershire, Cheltenham, GL50 4AZ, UK

¹²CNRS and Université de Toulouse, Castanet Tolosan, 31326, France

¹³Department of Biogeography and Paleoecology, Adam Mickiewicz University, Poznan, 61-680, Poland

¹⁴Department of Earth Sciences, University of Toronto, Toronto, ON M5S 3B1, Canada

¹⁵Departement de Géographie and GEOTOP, Université du Québec – Montréal, Montréal, QC H3C 3P8, Canada

¹⁶Department of Geology, Lund University, Lund, SE-223 62, Sweden

- 65
66 ¹⁷Department of Geography, University of California – Los Angeles, Los Angeles, CA
67 90095, USA
68
69 ¹⁸U.S. Geological Survey, Reston, VA 20192, USA
70
71 ¹⁹Department of Geosciences and Natural Resource Management, University of
72 Copenhagen, Copenhagen, DK-1350, Denmark
73
74 ²⁰Department of Environmental Sciences, University of Helsinki, Helsinki, FIN-00014,
75 Finland
76
77 ²¹Department of Chemical and Environmental Engineering, University of Nottingham,
78 Nottingham, NG7 2RD, UK
79
80 ²²Département de Géographie and Centre d'études nordiques, Université Laval, Québec,
81 QC G1V 0A6, Canada
82
83 ²³Geological Survey of Finland, P.O. Box 96, Espoo, 02151, Finland
84
85 ²⁴School of Geosciences, University of Aberdeen, Aberdeen, AB24 3UF, UK
86
87 ²⁵Department of Geography and Global Environmental and Climate Change Centre,
88 McGill University, Montreal, QC H3A 0B9, Canada
89
90 ²⁶Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964, USA
91
92 ²⁷Centre for Economic Development, Transport and the Environment, Vaasa, 65101,
93 Finland
94
95 ²⁸Department of Geography, University of Toronto, Toronto, ON M5S 3G3, Canada
96
97 ²⁹Département de Géographie, Université de Montréal, Montréal, QC H2V 2B8, Canada
98
99 ³⁰Champlain College – Dublin Campus, Dublin, Ireland
100
101 ³¹Agriculture and Agri-Food Canada, Ottawa, ON K1A 0C6, Canada
102
103 ³²Yorkshire Peat Partnership, Yorkshire Wildlife Trust, York, YO24 1GN, UK
104
105 ³³Department of Integrative Biology, University of Guelph, Guelph, ON N1G 2W1,
106 Canada
107
108 ³⁴BIAX Consult, Zaandam, 1506 AL, The Netherlands
109

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

110 ³⁵Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam,
111 Amsterdam, 1098 XH, The Netherlands
112
113 ³⁶Department of Plant Biology, Southern Illinois University, Carbondale, IL 62901, USA
114
115 ³⁷Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of
116 Sciences, Beijing, 100101, China
117
118 ³⁸Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, 710075 Shaanxi,
119 China
120

121 Corresponding authors

122 Julie Loisel email: jul208@lehigh.edu phone: 610-758-3660
123 Zicheng Yu email: ziy2@lehigh.edu phone: 610-758-6751

125 Abstract

126 Here we present results from the most comprehensive compilation of Holocene peat soil
127 properties with associated carbon and nitrogen accumulation rates for northern peatlands.
128 Our database consists of 268 peat cores from 215 sites located north of 45°N. It
129 encompasses regions within which peat carbon data have only recently become available,
130 such as the West Siberia Lowlands, the Hudson Bay Lowlands, Kamchatka in Far East
131 Russia, and the Tibetan Plateau. For all northern peatlands, carbon content in organic
132 matter was estimated at $42 \pm 3\%$ (S.D.) for *Sphagnum* peat, $51 \pm 2\%$ for non-*Sphagnum*
133 peat, and at $49 \pm 2\%$ overall. Dry bulk density averaged $0.12 \pm 0.07 \text{ g cm}^{-3}$, organic
134 matter bulk density averaged $0.11 \pm 0.05 \text{ g cm}^{-3}$, and total carbon content in peat
135 averaged $47 \pm 6\%$. In general, large differences were found between *Sphagnum* and non-*Sphagnum*
136 *Sphagnum* peat types in terms of peat properties. Time-weighted peat carbon
137 accumulation rates averaged $23 \pm 2 \text{ (S.E.M.) g C m}^{-2} \text{ yr}^{-1}$ during the Holocene on the

basis of 151 peat cores from 127 sites, with the highest rates of carbon accumulation (25-28 g C m⁻² yr⁻¹) recorded during the early Holocene when the climate was warmer than the present. Furthermore, we estimate the northern peatland carbon and nitrogen pools at 436 and 10 gigatons, respectively. The database is publicly available at <https://peatlands.lehigh.edu>.

Keywords

Northern peatlands, Data synthesis, Climate change, Soil carbon and nitrogen, Biogeochemical cycles, Long-term ecosystem dynamics

Introduction

Of all terrestrial ecosystems, peatlands are arguably the most efficient at sequestering carbon (C) over long time scales. Northern peatlands cover approximately 4,000,000 km² or 3% of the global land area (Maltby and Immirzi, 1993) and have accumulated about 500 gigatons of C (GtC) mostly during the Holocene, equivalent to ~ 30% of the present-day global soil organic carbon (SOC) pool (Gorham, 1991; Bridgham et al., 2006; Yu et al., 2010). These ecosystems have also played a dynamic role in the Holocene global C cycle as important sinks of carbon dioxide (CO₂) and major sources of methane (CH₄) to the atmosphere (Frolking and Roulet, 2007; Korhola et al., 2010; Yu, 2011). As climate warming positively affects both plant growth and organic matter decomposition, recent and projected climate change could shift the balance between peat production and organic matter decomposition, potentially affecting the peatland C-sink capacity and modifying peat C fluxes to the atmosphere (Frolking et al., 2011; Yu, 2012). This prediction

1
2
3 161 particularly holds true for the northern high-latitude regions, where the intensity of
4
5 162 climate change is expected to be greatest (McGuire et al., 2009). The peatland C cycle –
6
7
8 163 climate feedback remains difficult to assess, however, because of (1) limited
9
10 164 understanding of peatland responses to climate change (Frolking et al., 2011), (2) data
11
12 165 gaps and large uncertainties in regional peatland C stocks (Yu, 2012), and (3) non-linear
13
14 166 peatland responses to external forcing (Belyea, 2009).
15
16
17 167
18
19
20 168 Very little is known about the nitrogen (N) budget that accompanies C accumulation in
21
22 169 northern peatlands (but see Limpens et al., 2006 for a review). Assuming a net C
23
24 170 sequestration of ~270 GtC (Yu, 2012) and a C/N ratio of 20-30 for fen peat (Bergner et
25
26 171 al., 1990; Rydin and Jeglum, 2013; results therein) during the early Holocene (11-7 ka),
27
28 172 about 10-13 GtN would have been required to build such peat deposits. It is therefore
29
30 173 possible that northern peatlands have been playing an undocumented, dynamic role in the
31
32 174 Holocene global N cycle as important sinks of N, potentially limiting the amount of N
33
34 175 available for other ecosystems at the global scale (McLauchlan et al., 2013).
35
36
37 176 Alternatively, if the main N input to peatlands was through N₂ fixation by cyanobacteria,
38
39 177 these microorganisms might have been more important in driving the C cycle in
40
41 178 peatlands than previously thought. Overall, studying the coupling between N and C
42
43 179 cycling in northern peatlands is essential for a better understanding of how key
44
45 180 biogeochemical processes interact in these systems and for predicting the future of peat C
46
47 181 stocks.
48
49
50 182
51
52
53
54
55
56
57
58
59
60

Here we present the most comprehensive compilation of Holocene C and N data for northern peatlands. This synthesis encompasses regions within which peat C and N data have only recently become available, such as the West Siberian Lowlands in Asian Russia, the Hudson Bay Lowlands in Canada, Kamchatka in the Russian Far East, and the Tibetan Plateau. In addition, we present the most comprehensive synthesis of peat soil properties (such as bulk density, organic matter content, C and N content) from the northern hemisphere. Also, this new database and synthesis work represent a major expansion from Yu et al.'s (2009) synthesis on Holocene peat C dynamics, which was based on 33 sites (vs. 127 sites as reported in this paper). Finally, it constitutes a natural continuation of Charman et al.'s (2013) recent study on peat C accumulation in northern peatlands during the last millennium.

In addition to filling regional data and knowledge gaps, the main objectives of this paper are to (1) describe a database of peat soil properties and synthesize this information for different peat types, time intervals, and geographic regions, and (2) produce time series of Holocene peat C and N accumulation rates in 500-yr bins for comparison with climate history. Key differences in Holocene peat properties between different regions and peatland types are also discussed in light of their implications for long-term peat C stocks. Of particular importance and relevance are the differences between *Sphagnum* and non-*Sphagnum* peat types. Finally, we present new estimates for northern peat C and N stocks for northern peatlands on the basis of the expanded database.

Database and analysis

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

206 Database

207 We compiled a dataset of 268 published and unpublished Holocene peat records from
208 215 sites located in North America and Eurasia (Figure 1, Supplementary Table 1). The
209 difference in the number of peat cores and peatland sites is due to the fact that, in a few
210 instances, multiple cores were collected from a single peatland. As these multiple cores
211 were not designed as true replicates in the original publications, each of these cores was
212 considered as an independent record of peat properties in the present study. However,
213 only the oldest core for each site was used for estimating peat inception age. Finally,
214 when calculating peat C accumulation rates, multiple cores from a single site were each
215 attributed an equal fraction of the weight for that site. For example, for a site with three
216 cores, the peat C accumulation history of each core only accounted for 1/3 of the site's
217 record.

218
219 The latitude of most peatland sites ranges from 45 to 69 °N. The cutoff at 45 °N
220 represents the southern limit for defining what is considered to be the area contributing to
221 the C cycle of the Arctic region (McGuire et al., 2009). Four high-elevation sites found in
222 China (the Tibetan Plateau) and Japan were also included, as they developed under
223 similar 'northern' climatic conditions. The name and coordinates of these four sites are as
224 follows: Zoige (33.5 °N, 102.6 °E), Hongyuan (32.8 °N, 102.5 °E), Hani (42.2 °N, 126.5
225 °E), and Utsai (42.4 °N, 140.2 °E). A total of 155 cores originate from Eurasia,
226 including 112 cores from Russia. The remaining 113 cores come from North America.
227 Approximately 40% of all cores were collected from ombrotrophic bogs (n = 110 cores)
228 and 20% were extracted from minerotrophic fens (n = 50 cores). The remainder (40%)

229 was collected in peatlands currently affected by permafrost ($n = 108$ cores). Note that
230 peatland type was identified independently for each site by the original investigators.
231 From an ecosystem functioning perspective, distinguishing bogs from fens and
232 permafrost peatlands is important, as these peatland types are characterized by different
233 hydrological regimes, vegetation communities, and peat-growth trajectories, all of which
234 impact long-term rates of peat C sequestration (Rydin and Jeglum, 2013). Bogs are
235 mineral-poor, rain-fed peatland ecosystems with relatively low plant net primary
236 production (NPP) and slow peat decomposition rates. In contrast, fens are hydrologically
237 connected to surface or ground water, thereby receiving more mineral nutrients.
238 Generally speaking, fens have greater NPP but also faster peat decay rates than bogs
239 (Blodau, 2002). Finally, in the sub-Arctic and Arctic regions, peatland hydrology,
240 structure, and peat C balance are sensitive to the underlying permafrost aggradation and
241 degradation dynamics (Camill, 1999; Turetsky et al., 2007). For the analysis, peat
242 plateaus (87 out of 108 cores), permafrost bogs (18 out of 108 cores), and collapse scars
243 (3 out of 108 cores) were grouped under the peatland type ‘permafrost peatlands’.
244 Original peatland categories can be found in Supplementary Table 1.
245
246 The database was built to include as many peat records as possible. Therefore, we
247 included any peat core that was extracted north of 45 °N (or at high elevation) and for
248 which bulk density or organic matter bulk density data were available. Information
249 related to peat-core location, peatland type, peat properties, age, and data source can be
250 found in Supplementary Table 1. Additional information related to the type of coring
251 device used and the year of coring can be found in the original publications. Data used in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

252 this synthesis are readily accessible from the Holocene Peatland Carbon Network website
253 (<https://peatlands.lehigh.edu>). This database will be useful in future studies of ecosystem
254 – C cycle – climate interactions and for modeling long-term peatland dynamics.

255

256 In the following sub-sections, we present the criteria for site selection and the protocols
257 used to develop the database. In an effort to only analyze and synthesize *peat* samples,
258 inorganic-rich horizons often found at the base of the peat cores were removed from the
259 database. When available, stratigraphic information was used to distinguish peat vs. non-
260 peat material. For example, gyttja (organic-rich lake sediments) was excluded from the
261 dataset, as well as marshy, clayey, and silty sediments. When stratigraphic information
262 was not available, a bulk density value of 0.5 g cm^{-3} was used as a cut-off between peat
263 and non-peat material. This value was chosen on the basis of stratigraphic information
264 from peatland records where peaty sediments with bulk density values up to 0.5 g cm^{-3}
265 were identified. We acknowledge that this cut-off value is arbitrary, and that our dataset
266 likely contains some non-peat samples. Inorganic horizons (e.g., tephra layers) were also
267 excluded from the database.

268

269 Peat properties

270 A total of 232 peat cores (181 sites) were used for characterizing peat properties, though
271 not all cores have all types of peat properties available (Figure 2a). This dataset contains
272 139 cores from Eurasia (including 109 cores from Russia) and 93 cores from North
273 America (Figure 1). While approximately half of these cores were sampled and analyzed
274 at high resolution (1-5 cm increments), the remainder was sampled at lower resolution,

typically at 10 cm increments. Dry bulk density (BD; g cm⁻³), organic matter content (OM%; gravimetric %), as well as elemental C and N concentration values were compiled and synthesized. On the basis of these raw datasets, C/N mass ratio and organic matter bulk density (OMBD; g OM cm⁻³) were calculated. These peat geochemical values are examined in light of peat stratigraphy, peat ages, and geographic regions. The following paragraphs briefly describe the protocols used to obtain these values.

Peat stratigraphic information was obtained for 83 peat cores ('peat types' in Figure 2a) for which plant macrofossil analysis or detailed peat description had been performed following standard techniques (e.g., Troels-Smith, 1955; Mauquoy and van Geel, 2007). This peat stratigraphic information was condensed into the following five peat types: *Sphagnum*, herbaceous, woody, brown moss, and humified peat. In a few cases, the investigators only ascribed a general peat type to the samples (e.g., 'bog' vs. 'fen' peat, or '*Sphagnum*' vs. 'herbaceous' peat). In these cases, the uncertainty associated with classifying peat samples mostly relates to the uniformity of naming convention used among the investigators. For example, a peat sample that contains sizable fractions of brown moss and humified peat may be classified by an investigator as 'brown moss peat' and by another one as 'humified peat'. We recognize that, due to their nature, brown moss and humified peat types might be less uniform than *Sphagnum* or herbaceous peat types. We included as much stratigraphic information as possible in the database, though ambiguous or imprecise descriptions were left out to avoid further confusions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

297 Dry bulk density and organic matter content were determined following standard
298 procedures (Dean, 1974; Chambers et al., 2011). Peat samples of a known fresh volume
299 were either freeze-dried or oven-dried at ca. 100 °C until constant weight was reached
300 and weighed to determine bulk density, then burned at 500-600 °C for one to four hours
301 and weighed again to determine organic matter content. The accuracy of these
302 measurements mostly depends on sample handling (care must be taken to prevent peat
303 compaction in the field and in the laboratory) and the analytical error associated with
304 weighing. The product of bulk density (BD; g cm⁻³) and organic matter content (OM%)
305 of each peat sample was used to calculate organic matter bulk density (OMBD; g OM
306 cm⁻³), also referred to as ash-free bulk density (AFBD) or organic bulk density (OBD) in
307 the literature (Yu et al., 2003; Björck and Clemmensen, 2004). We compiled a total of
308 21,220 bulk density measurements, 18,973 organic matter content values, and computed
309 18,544 organic matter bulk density values (Figure 2a).
310
311 Total peat C and N content were directly measured by combustion and elemental analysis
312 of dry peat samples (Chambers et al., 2011). We compiled 3741 C and 3365 N
313 measurements (Figure 2a). We also computed a total of 3362 C/N mass ratio values.
314
315 Finally, the regression between peat C content (C%) and peat organic matter content
316 (OM%) for each peat type is presented as an estimate for C content in organic matter
317 (OC%). A total of 995 samples were used in this analysis. The slope of each one of these
318 regressions is interpreted as the ‘conversion factor’ from OM% to OC%, such that it

provides an indirect way for estimating the C% content of ash-free peat for investigators who do not perform elemental C measurements directly.

Peat-core chronology

Peat-core chronologies were almost exclusively based on radiocarbon (^{14}C) dates that were determined mostly by accelerator mass spectrometry (AMS) on terrestrial plant macrofossils or bulk peat (Piotrowska et al., 2011). A few older chronologies were based on conventional ^{14}C dating of bulk samples. Because no systematic offset has been observed in the ^{14}C age of bulk vs. non-woody plant macrofossils (G.M. MacDonald, pers. comm. 2013), the use of bulk dates is justifiable. For the purpose of this study, all ^{14}C dates were calibrated to calendar years before present (cal. BP) using the program CALIB 6.1.0 (Stuiver and Reimer, 1993) with the IntCal09 calibration dataset (Reimer et al., 2009). In this paper, ages are reported in thousands of calibrated years before present (ka).

Age-depth relationships were established for all continuous peat cores for which at least five age determinations were available ($n = 151$ cores). Except for a few palynostratigraphic and tephrochronologic markers, nearly all records have chronologies exclusively based on ^{14}C ages (Supplementary Table 1). Chronologies were obtained through linear interpolation of calibrated ages between dated horizons. Single-age estimates were taken from the mid-point of each calibrated 2σ probability distribution. This parsimonious approach captures general patterns of temporal changes in peat accumulation and allows for analysis of peat C accumulation trajectories (e.g., Telford et

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

342 al., 2004). Although more sophisticated approaches are possible (e.g., Charman et al.,
343 2013), we seek to make the fewest or simplest assumptions for this analysis because the
344 temporal resolution target for peat C accumulation rate calculations is relatively low (at
345 500 years). In the cases where the original investigator identified hiatuses (e.g., peat loss
346 caused by erosion or fire) or depositional anomalies (e.g., thick tephra layers that
347 interrupted peat accumulation) along their peat records, these gaps were taken into
348 consideration when building age-depth relationships (Glaser et al., 2012). Otherwise, peat
349 records were assumed to be continuous.

350

351 In an effort to assess the representativeness of our samples in terms of peatland inception
352 timing, peat basal ages were compiled and compared to results from large datasets
353 (MacDonald et al., 2006; Gorham et al., 2007; Korhola et al., 2010). Peat inception ages
354 from 199 sites (Supplementary Table 1) were summed and binned in 500-year intervals.

355

356 Long-term rates of carbon and nitrogen accumulation

357 A total of 151 peat cores from 127 sites were used for estimating rates of peat C
358 accumulation. This dataset contains 96 cores from 78 North American sites and 55 cores
359 from 49 Eurasian sites (Figure 1). Of the 33 sites presented in Yu et al.'s (2009) study, 25
360 were used in the present study (Supplementary Table 1). The remaining 8 sites did not
361 fulfill our dating quality criterion (presented below). The dating quality of each record
362 was determined by the quotient of the calibrated peat basal age and the number of age
363 determinations. For example, a 10,000-year-old peat core with a chronology constrained
364 by 10 ¹⁴C dates was attributed a dating quality of one date per 1000 years. About 58% of

our 151 cores were characterized by an acceptable dating quality of one to two ^{14}C dates per 1000 years (Figure 2b). These resolutions are well suited to capture millennial-scale variations in C accumulation. Several peat cores with more than two ^{14}C dates per 1000 years ($n = 35$ cores) were available from North America and Europe. The lower dating quality cores (<1 date per 1000 years,) were unevenly distributed and comprise 44% of the North American records and 40% of the Eurasian records.

The long-term rate of peat C accumulation was calculated for each core following one of the following five approaches (Supplementary Table 1): (1) whenever possible, peat core chronology was combined with bulk density and C% for each depth increment ($n = 47$ cores); (2) in the cases where direct C measurements were lacking, peat core chronology was combined with organic matter bulk density measurements and a mean organic C value of 49% in organic matter ($n = 57$ cores); (3) for cores that lacked organic matter bulk density and direct C%, peat core chronology was combined with bulk density measurements and a mean C content of 47% in total peat ($n = 3$ cores); (4) whenever neither bulk density nor C% was directly available from the cores, long-term rate of peat C accumulation was calculated for each core by combining time-dependent bulk densities (0.08 g cm^{-3} at 0-0.5 ka; 0.12 g cm^{-3} at 0.5-6 ka; 0.14 g cm^{-3} at 6-12 ka) with a mean C content of 47% in total peat for each dated interval ($n = 32$ cores), and (5) peat C accumulation rates for the remaining 12 cores were directly obtained from published figures and tables. For all the cores, time-weighted peat C accumulation rates were summed and binned in 500-year intervals. It is important to note that such reconstructions are ‘apparent rates’ that are different from true rates of C accumulation in peatlands,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

388 because decomposition processes have been affecting old peat layers for thousands of
389 years (Turunen et al., 2002). Finally, the long-term rate of peat N accumulation was
390 calculated by combining our binned peat-C accumulation rates with time-dependent C/N
391 values (65 at 0-6 ka, and 40 at 6-10 ka).

392

393 **Results**

394 *Peat Properties*

395 *Descriptive statistics of dataset*

396 The frequency distribution of each peat property is shown in Figure 3. Mean values and
397 standard deviations for all peat properties are presented by peat type in Table 1, and by
398 region in Table 2.

399

400 Bulk density values (n = 21,220) ranges from 0.003 to 0.498 g cm⁻³, with a mean value of
401 0.118 ± 0.069 g cm⁻³ (1 standard deviation (S.D.)). A one-way analysis of variance
402 (ANOVA) reveals an effect of peat type on bulk density ($F(10709) = 941, p < 0.0001$),
403 with all peat types significantly different from each other on the basis of post-hoc
404 Tukey's LSD tests ($p < 0.0001$). In increasing order, mean bulk density of the peat types
405 is *Sphagnum* < Woody < Herbaceous < Brown Moss < Humified (Table 1).

406

407 Organic matter content (n = 18,973) has a mean value of 90.7 ± 13% (1 S.D.) and a
408 median of 95.7%. The ANOVA reveals an effect of peat type on organic matter content
409 ($F(9512) = 349, p < 0.0001$), with all peat types significantly different from each other

(Tukey's LSD: $p < 0.0001$). In decreasing order, mean organic matter content of the peat types is *Sphagnum* > Woody > Herbaceous > Brown Moss > Humified (Table 1).

412

Organic matter bulk density ($n = 18,544$) ranges from 0.003 to 0.452 g OM cm⁻³, with a mean value of 0.105 ± 0.051 g OM cm⁻³ (1 S.D.). The ANOVA reveals an effect of peat type on organic matter density ($F(9081) = 942$, $p < 0.0001$), with all peat types significantly different from each other (Tukey's LSD: $p < 0.0001$). In increasing order, mean organic matter density of the peat types is *Sphagnum* < Herbaceous < Woody < Brown Moss < Humified (Table 1).

419

C content in total peat ($n = 3741$) ranges from 30 to 60%, with a mean value of $46.8 \pm 6.1\%$ (1 S.D.) and a median of 47.8%. While the lowest values (< 35%) are almost exclusively associated with samples from Alaska, western Canada, Fennoscandia, and eastern Russia, the highest values (> 55%) are characteristic of sites located in the western European Islands and Fennoscandia. The ANOVA reveals an effect of peat type on C% ($F(2494) = 161$, $p < 0.0001$), with *Sphagnum* samples significantly different from other peat types (Tukey's LSD: $p < 0.0001$). Herbaceous and woody peats are distinct from other types, but indistinguishable from one another (Tukey's LSD: $p = 0.238$).

Likewise, humified and brown moss peats are distinct from other types, but indistinguishable from each other (Tukey's LSD: $p = 0.448$). In increasing order, mean C% of the peat types is *Sphagnum* < Humified = Brown Moss < Herbaceous = Woody (Table 1). The frequency distribution of C% in peat is also characterized by a second, though minor, mode at 40% (Figure 3d). The latter is mostly associated with *Sphagnum*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

433 peat samples, as their average C% is significantly lower than those for other peat types
434 (Table 1). This difference is likely caused by the high content of complex and recalcitrant
435 compounds found in *Sphagnum* tissues such as lipids and waxes, which have lower C%
436 than more labile biopolymers such as cellulose (Cagnon et al., 2009).
437
438 On the basis of 995 samples for which both OM% and C% were quantified, we
439 developed conversion factors (slopes of linear regressions) for several peat types to
440 estimate OC% (Figure 4). There is a noticeable difference between the slope of
441 *Sphagnum* peat (0.423 ± 0.030 ; $n = 454$) and that of non-*Sphagnum* peat (0.514 ± 0.024 ;
442 $n = 308$). The C content in organic matter (OC%) for *Sphagnum* peat is smaller than
443 expected at $42.3 \pm 3.0\%$ (e.g., Bauer et al., 2006; Beilman et al., 2009; Table 3). As the
444 majority of our *Sphagnum* samples for this specific analysis are younger than 0.5 ka (304
445 out of 454 samples) and extracted from raised bogs, it is very likely that our estimated
446 OC% biases towards young and undecomposed *Sphagnum*. This ‘young *Sphagnum* peat
447 effect’ heavily influenced the slope of the overall relation between C% and OM% (0.467
448 ± 0.045) due to the overrepresentation of *Sphagnum* samples in the dataset (454 out of
449 995 samples). To minimize this bias when estimating mean OC% in peat, all 304 young
450 *Sphagnum* samples were removed from the dataset, yielding an overall conversion factor
451 of 0.492 ± 0.024 .
452
453 N content in peat ($n = 3365$) ranges from 0.04 to 3.39%, with a mean value of $1.2 \pm 0.7\%$
454 (1 S.D.). The frequency distribution is asymmetric and characterized by a mode at 0.65%
455 (Figure 3e). The latter is largely due to the overrepresentation of *Sphagnum* peat samples

456 (having low N content) in our database. The ANOVA reveals an effect of peat type on N
 457 content ($F(2504) = 666, p < 0.0001$) with *Sphagnum* and herbaceous peat types
 458 significantly different from each other and from all other types (Tukey's LSD: $p <$
 459 0.0001). Humified and brown moss peats are distinct from other types, but
 460 indistinguishable from each other (Tukey's LSD: $p = 0.113$). Likewise, woody and brown
 461 moss peats are indistinguishable from one another (Tukey's LSD: $p = 0.240$). In
 462 increasing order, mean N% of the peat types is *Sphagnum* < Woody = Brown Moss =
 463 Humified < Herbaceous (Table 1).
 464
 465 C/N mass ratio ($n = 3362$) ranges from 12 to 217, with a mean value of 55 ± 33 (1 S.D.).
 466 The frequency distribution is asymmetric and characterized by a mode at 25 (Figure 3f).
 467 While the distribution mode (25) is associated with non-*Sphagnum* peat types, the
 468 distribution mean (55) is skewed towards *Sphagnum* peat samples owing to their
 469 overrepresentation in our database (Figures 3f). The ANOVA reveals an effect of peat
 470 type on C/N ratio ($F(2501) = 174, p < 0.0001$) with *Sphagnum* peat significantly
 471 different from all other types (Tukey's LSD: $p < 0.0001$). Woody peat is distinguishable
 472 from all peat types except for brown moss peat (Tukey's LSD: $p = 0.665$). Herbaceous
 473 and humified peat types are indistinguishable from one another (Tukey's LSD: $p =$
 474 0.721). In decreasing order, the mean C/N ratio of peat types is *Sphagnum* > Woody =
 475 Brown Moss > Humified = Herbaceous (Table 1).
 476
 477 Temporal changes in peat properties

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

478 We find a decreasing trend in bulk density over the Holocene (Figure 5a), with the
479 densest peat characterizing the oldest samples (8-10 ka) and the least-dense peat
480 characterizing the youngest samples (0-2 ka). This trend is most likely attributable to the
481 progressive decomposition and subsequent compaction of peat over time, as well as to
482 higher ash content in early-stage peat (likely from fens). Conversely, a clear increasing
483 trend in organic matter content (OM%) is found over the Holocene (Figure 5b), with the
484 greatest OM% characterizing the youngest samples (0-2 ka) and the least OM%
485 characterizing the oldest samples (8-10 ka). This trend is most likely attributable to
486 higher inorganic material inputs during early-stage peatland development as well as to a
487 greater loss of OM in the deeper portions of peat profiles. Organic matter bulk density
488 (OMBD) remains relatively constant over the Holocene (Figure 5c) because of the
489 opposite trends exhibited by BD and OM% (Figures 5a, 5b). The only exceptions are the
490 low OMBD values characterizing the youngest samples (< 0.5 ka), probably due to the
491 large proportion of young, undecomposed *Sphagnum* peat samples.
492
493 C content in peat remains uniform over the Holocene (Figure 5d), except for slightly
494 lower C% during the late Holocene. We find a decreasing trend in N% over the
495 Holocene, such that young peat deposits are associated with low N% (Figure 5e). Peat
496 deposits older than 6 ka are mostly associated with low C/N ratios, whereas peat samples
497 younger than 6 ka are characterized by high C/N values (Figure 5f).
498
499 In general, *Sphagnum* peat is characterized by lower BD, OMBD, C%, N%, and C/N
500 ratio than samples composed of non-*Sphagnum* peat (Figure 6). Therefore, peatland

development could explain much of the aforementioned temporal trends (Figure 5), as early-stage rich fens are typically characterized by non-*Sphagnum* peat, whereas late-stage poor fens and bogs are *Sphagnum*-dominated (Figure 6h).

Spatial differences in peat properties

Significant differences in bulk density are found at the regional scale (Table 2). The densest peat is observed in Alaska (mean = $0.168 \pm 0.087 \text{ g cm}^{-3}$) and western Canada (mean = $0.166 \pm 0.076 \text{ g cm}^{-3}$), whereas the least dense peat is found from the western European Islands (mean = $0.055 \pm 0.027 \text{ g cm}^{-3}$). Organic matter bulk density values follow a similar pattern across these regions (Table 2). These differences are strongly correlated with peat types and sample ages, with the Alaskan and western Canadian samples largely constituted of herbaceous, humified, and brown moss peat types. OM% does not vary much between regions (> 90% in all regions), with the notable exception of Alaskan and eastern Russian/Asian peatlands that exhibit mean values of $76.6 \pm 18.8\%$ and $80.3 \pm 16.7\%$, respectively (Table 2). Aeolian dust and tephra ash inputs to some peatlands in Alaska, Kamchatka, and Japan might partly explain such low OM% values.

Peat inception ages and long-term rates of carbon and nitrogen accumulation

Calibrated ages (mid-point) for peat inception range from 0.6 to 15 ka and the frequency distribution is characterized by a mode at 11-9 ka (Figure 2c). The latter corresponds with peat inception peaks in the West Siberian Lowlands and in Alaska. In general, our samples are in agreement with much larger networks of peat basal ages (Smith et al.,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

523 2004; MacDonald et al., 2006; Gorham et al., 2007; Korhola et al., 2010; Ruppel et al.,
524 2013; Yu et al., 2013).

525

526 The time-weighted long-term rate of C accumulation averages $22.9 \pm 2.0 \text{ g C m}^{-2} \text{ yr}^{-1}$
527 (standard error of mean (S.E.M.); Figure 7b). Values exhibit an increasing trend that
528 initially peaks during the early Holocene between 10 and 7.5 ka at $27.0 \pm 2.6 \text{ g C m}^{-2} \text{ yr}^{-1}$.
529 This peak is largely caused by rapid peat accumulation in Alaska, the Western Siberia
530 Lowlands, and southeastern Canada. The remainder of the Holocene is characterized by a
531 decreasing trend in C accumulation rates from 24 to $18 \text{ g C m}^{-2} \text{ yr}^{-1}$ and a time-weighted
532 mean at $22.0 \pm 1.9 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Figure 7b). There is a notable minimum value between 3
533 and 1.5 ka at $18\text{-}19 \text{ g C m}^{-2} \text{ yr}^{-1}$. Lack of decomposition probably explains most of the
534 apparent increase in accumulation over the past millennium ($24\text{-}32 \text{ g C m}^{-2} \text{ yr}^{-1}$), as
535 young peat appears to be accumulating more quickly than old peat simply because the
536 former has undergone less decomposition than the latter (Clymo, 1984).

537

538 The time-weighted long-term rate of N accumulation averages $0.5 \pm 0.04 \text{ g N m}^{-2} \text{ yr}^{-1}$
539 (S.E.M.; Figure 7c). While the mid and late Holocene (6-0 ka) are characterized by the
540 lowest rates of N accumulation at $0.34 \text{ g N m}^{-2} \text{ yr}^{-1}$, the highest rates ($0.61 \text{ g N m}^{-2} \text{ yr}^{-1}$)
541 occur between 12 and 6 ka (Figure 7c). This trend mirrors that of C accumulation (Figure
542 7b), as C and N sequestration rates are both mainly influenced by peat density and its
543 accumulation rate. The low rates of N accumulation over the past 6 ka might also relate
544 to the increasing presence and persistence of *Sphagnum* (having high C/N ratio and low
545 N concentration) in northern peatlands (Figures 6 and 7).

546

547 **Discussion**

548 *Representativeness of the database for northern peatlands*

549 The present database contains the most comprehensive compilation of peat properties and
550 C accumulation records for northern peatlands. The previous large-scale synthesis (Yu et
551 al., 2009) only contained 33 sites and lacked records from the Hudson Bay lowlands and
552 the Russian Far East, and had limited sites from West Siberia and the western European
553 Islands. The present database fills gaps from these regions.

554

555 However, European Russia, East Siberia, and the Russian Far East clearly remain poorly
556 studied regions in terms of northern peat C stocks and accumulation histories (Figure 1).
557 A wetland map by Stolbovoi and McCallum (2002) suggests that shallow peaty deposits
558 (<50 cm) interspersed with few deeper peat bogs (>50 cm) dominate the Far East Russian
559 landscape. Most of these deeper peatlands are presumably found in Kamchatka and
560 Sakhalin (Stolbovoi, 2002). This broad portrait is, however, based on fewer than 30 soil
561 profiles from across East Siberia and Far East Russia (Stolbovoi et al., 2001), making it
562 difficult to evaluate the importance of this region in the northern peatland C cycle. In
563 general, peat C stocks in Eastern Russia may not be as massive as those from West
564 Siberia or European Russia (Stolbovoi and McCallum, 2002). Therefore, understanding
565 how these shallow peatlands in East Siberia and the Russian Far East have developed
566 during the Holocene would provide useful end-members of climate controls of peat C
567 accumulation, but these peatlands do not seem to represent a large missing C stock.

568

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

569 *Northern peatland soil properties: key findings and uncertainties*

570 Peat-carbon stocks

571 Several studies have quantified the soil C density and total C pool of peatlands using

572 different approaches (e.g., Armentano and Menges, 1986; Gorham, 1991; Yu et al.,

573 2010). These methods have led to total C pool estimates for northern peatlands that vary

574 by at least a factor of two, from 234 to 547 GtC (Lappalainen, 1996; Yu et al., 2010; see

575 Yu, 2012 for a review). Many of these studies have combined mean peat depth, modern

576 peatland area, and a single mean C density value (BD x C% or OMBD x OC%) in their

577 calculations (e.g., Gorham, 1991). Applying Gorham's (1991) mean peat depth and

578 peatland area estimates to the mean BD and C% results from our database yields a C pool

579 estimate of 436 GtC ($2.3 \text{ m} \times 3.42 \text{ Mkm}^2 \times 0.118 \text{ g cm}^{-3} \times 47\% \text{ C}$). However, it is well

580 documented that most peatlands undergo a shift from herbaceous to *Sphagnum* peat

581 during their developmental history (Hughes, 2000; Figure 6h) and that different BD, C%,

582 and rates of peat accumulation are associated with fen and bog peats (e.g., Vitt et al.,

583 2000; Figure 6). We also know that peat-C accumulation rates have varied

584 asynchronously between regions throughout the Holocene as a result of regional changes

585 in hydroclimatic conditions (e.g., Yu et al., 2009; Charman et al., 2013). Therefore, we

586 argue that reconstructing Holocene changes in peat C accumulation on the basis of

587 *measured* peat C density and reliable peat-core chronologies constitutes a step forward in

588 providing the best possible peat C stock estimates (see Yu et al., 2010 for an example). It

589 also allows for quantifying spatial and temporal differences in rates of peat C

590 accumulation, as well as the temporal trajectories of peat C fluxes to the atmosphere

(MacDonald et al., 2006; Yu et al., 2013). However, better maps of the present peatland area (and its change over time) are still needed to improve current peat C stock estimates.

Carbon content in organic matter

For each peat layer, peat C density can be estimated by the product of either (1) bulk density and C content in total peat (BD x C%), or (2) ash-free bulk density and C content in organic matter (OMBD x OC%). It could be argued that the first option is preferable when estimating peat C stocks, as it produces values that are directly comparable to routine soil C measurements from other terrestrial ecosystems. However, the present database clearly indicates that the majority of peatland scientists routinely analyze organic matter content (OM%; n = 18,973 samples) rather than C% (n = 3741 samples) along peat cores. To provide a way to estimate OC% from OM%, we developed the following conversion factors: $42.3 \pm 3.0\%$ for *Sphagnum* peat, $51.4 \pm 2.4\%$ for non-*Sphagnum* peat, and $49.2 \pm 2.4\%$ overall (Figure 4, Table 1).

While the overall peat and the non-*Sphagnum* peat conversion factors are in line with those from previous studies (e.g., Gorham, 1991; Vitt et al., 2000; Turunen et al., 2002; Beilman et al., 2009), the *Sphagnum* peat factor is lower than other estimates (e.g., Bauer et al., 2006; Beilman et al., 2009; Table 3). Indeed, our mean OC% *Sphagnum* value at 42.3% is close to that of surface *Sphagnum* tissues, suggesting that it constitutes a valid estimate for ash-free and poorly decomposed *Sphagnum* peat. As previously mentioned, this bias towards low OC% is due to a large number of *Sphagnum* samples younger than 0.5 ka (304 out of 454 *Sphagnum* samples).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

614

615 Although each one of the three conversion factor slopes was significant ($p < 0.0001$),

616 there is a noticeable scatter in the data (Figure 4) that cannot solely be explained by the ~

617 1% analytical error associated with the loss-on-ignition procedure (Heiri et al., 2001).

618 The progressive accumulation of recalcitrant C in old samples (lignin ~ 60% C vs.

619 cellulose ~ 42% C), assuming it occurs at a greater rate than the loss of OM in the deeper

620 portions of peat profiles, could explain why C% appears higher than our OC% conversion

621 factors (Cagnon et al., 2009). The presence of inorganic C, particularly for the humified

622 and brown moss peat types, could also explain these results.

623

624 Oligotrophication and the fen-to-bog transition in northern peatlands

625 *Sphagnum* and non-*Sphagnum* peat types were characterized by very different peat

626 properties, with *Sphagnum* peat having lower BD, OMBD, C%, N%, and C/N ratio than

627 non-*Sphagnum* peat (Figure 6). These differences become important when estimating

628 Holocene peat C fluxes, as the proportion of *Sphagnum*-dominated peat records increases

629 during the late Holocene due to the fen-to-bog transition (Figure 6h). For example, much

630 stronger CH₄ emissions are associated with fens than bogs (e.g., Pelletier et al., 2007). In

631 terms of C sequestration rates, the systematically higher organic C density of non-

632 *Sphagnum* peat suggests that higher accumulation rates are possible in fens than in bogs

633 (Figure 6g), assuming optimal hydroclimatic conditions leading to rapid peat burial. In

634 addition, as non-*Sphagnum* samples contain twice the N mass of *Sphagnum* peat (Figure

635 6e), early-stage fens have the ability to stock more N than late-stage bogs. Overall,

636 further studies on the timing of the fen-to-bog transition across the northern peatland

domain are needed to better our understanding of its impact on C sequestration and CH₄ emissions.

Holocene pattern of carbon accumulation in northern peatlands

The overall trajectory and shape of our Holocene peat C accumulation curve is similar to the synthesis from a much smaller dataset ($n = 33$; Yu et al., 2009). As such, an early Holocene peak during the Holocene Thermal Maximum (HTM) and an overall slowdown of C accumulation during the mid- and late-Holocene, particularly after 4 ka during the Neoglacial period and associated permafrost development, were found in both syntheses (Figure 7). However, the mean Holocene value of $22.9 \pm 2.0 \text{ g C m}^{-2} \text{ yr}^{-1}$ (1 S.E.) presented here is approximately 24% higher than the estimate in Yu et al.'s 2009 study ($18.6 \text{ g C m}^{-2} \text{ yr}^{-1}$). Our larger dataset likely better represents the northern peatland C accumulation rates. These results imply that current peat C stocks might be underestimated.

While the peak value at $27 \text{ g C m}^{-2} \text{ yr}^{-1}$ is about 23% higher than the time-weighted mean peat-C accumulation rate for the remainder of the Holocene at $22 \text{ g C m}^{-2} \text{ yr}^{-1}$, we only found a 2% difference in organic C density values between young ($0.053 \pm 0.02 \text{ g C cm}^{-3}$) and old ($0.057 \pm 0.03 \text{ g C cm}^{-3}$) peat samples. These results clearly show that the peak value during the early Holocene cannot be mainly attributed to presumably dense peat deposits that would be rich in recalcitrant C due to long-term decomposition and compaction. Instead, factors influencing the rate of peat burial such as peat type (*Sphagnum* vs. non-*Sphagnum* peat; Figure 6), growing season length, and other

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

660 environmental variables, must have been responsible for such high rates of C
661 sequestration during the early Holocene.
662
663 The Holocene Thermal Maximum (HTM) is a well-documented period of orbitally-
664 induced warm climate in the northern high-latitude region (Kaufman et al., 2004;
665 Renssen et al., 2012; Marcott et al., 2013) that reaches its maximum around 11 ka
666 (Berger and Loutre, 1991; Figure 7a). The peak in warm climatic conditions shows a
667 transgressional pattern across northern North America that moved eastward with the
668 waning Laurentide Ice Sheet during the early and mid Holocene (Kaufman et al., 2004).
669 This progressive increase in land availability coupled with warming summer conditions
670 have been proposed as the main controls on peatland inception and rapid C accumulation
671 across northern North America (Harden et al., 1992; Gorham et al., 2007, 2012; Yu et al.,
672 2009; Jones and Yu, 2010). In general, our results support the hypothesis that warm
673 summers could promote peat formation and C sequestration (Beilman et al., 2009; Yu et
674 al., 2009; Charman et al., 2013), as the highest rates of C accumulation broadly coincide
675 with the peak in summer insolation from 11 to 7 ka (Figure 7). We acknowledge that
676 sufficient water input was necessary to allow for peatland development. Furthermore, the
677 observed temporal asymmetry in peatland inception age and peaks in C accumulation
678 rates between Alaska, western Canada, and the Hudson Bay Lowlands follows the
679 transgressional pattern of the HTM. For example, peat inception and highest peat C
680 accumulation rates occur at 11-9 ka in Alaska, whereas they are delayed in western
681 Canada with peak values around 9-7 ka. These findings have important implications for
682 projecting the fate of peat-C stocks in a future warmer world.

683

684 The Neoglacial period is characterized by generally cooler and wetter conditions than the
685 HTM (Figure 7a; Marcott et al., 2013). Particularly low C accumulation rates coincide
686 with this time period across the northern peatland domain (Figure 7; Vitt et al., 2000;
687 Jones and Yu, 2010). Peat accumulation processes might even have stopped in some
688 regions (e.g., Peteet et al., 1998). The onset of permafrost aggradation in many peatlands
689 also occurred during the Neoglacial period (Zoltai, 1971, 1995; Vitt et al., 2000; Oksanen
690 et al., 2003; Sannel and Kuhry, 2008), reducing the peat C-sink capacity. In addition to
691 shorter and cooler growing seasons, lower C accumulation rates in permafrost sites likely
692 relates to a slower peat burial due to (1) more intense peat decomposition in the acrotelm
693 due to drier surface conditions, and (2) a slower rate of peat formation and associated C
694 inputs to soil because many peat plateaus are not *Sphagnum*-dominated. Overall, our
695 results support the notion that climatic changes such as the HTM and the Neoglacial
696 cooling impact C sequestration rates in peatlands.

697

698 ***Role of northern peatlands in the global nitrogen cycle***

699 As relatively few downcore peat N concentrations have been reported in the literature, it
700 was difficult to compare our mean value of 1.2% to previous estimates. Bragazza et al.
701 (2012) reported N content values of 0.7% for *Sphagnum fuscum* litter and 1.48% for
702 *Eriophorum vaginatum* (herbaceous) litter, in line with our results (Table 1). Similarly,
703 Turunen et al. (2004) documented peat N concentrations ranging from 0.35 to 2.25%
704 (mean value of 0.8%) for the uppermost sections of 23 *Sphagnum* bogs across

1
2
3 705 northeastern Canada. Overall, these values closely match our findings for *Sphagnum*
4
5 706 (0.7%) and herbaceous (1.7%) peat types (Table 1).
6
7
8 707
9
10 708 Using our peat C pool estimate of 436 Gt and assuming a mean C/N ratio of 45 yields a
11
12 709 peat N pool of 9.7 Gt, roughly equivalent to 10% of the global soil N pool at 95 Gt (Post
13
14 710 et al., 1985). This estimate is within the range proposed by Limpens et al. (2006) at 8-15
15
16 711 GtN. The Holocene time-weighted peat N accumulation rate of $0.5 \pm 0.04 \text{ g N m}^{-2} \text{ yr}^{-1}$
17
18 712 (S.E.; Figure 7) is also in line with a previous estimate of 0.19-0.48 $\text{g N m}^{-2} \text{ yr}^{-1}$ (Limpens
19
20 713 et al., 2006). While the mid and late Holocene (6-0 ka) are characterized by the lowest
21
22 714 rates of peat-N accumulation at $0.34 \text{ g N m}^{-2} \text{ yr}^{-1}$, the highest rates ($0.61 \text{ g N m}^{-2} \text{ yr}^{-1}$)
23
24 715 occur between 12 and 6 ka (Figure 7c). The low rates of N accumulation over the past 6
25
26 716 ka might also relate to the increasing presence and persistence of *Sphagnum* peat (having
27
28 717 high C/N ratio and low N concentration) across the northern peatlands (Figures 6 and 7).
29
30 718 Overall, given the bias toward *Sphagnum*-dominated sites in our database, N pools and N
31
32 719 accumulation rates are probably underestimated.
33
34 720
35
36 721 Rapid N sequestration in peatlands during the early Holocene might have contributed to
37
38 722 the global decline in reactive N availability for terrestrial ecosystems (McLauchlan et al.,
39
40 723 2013), pointing to a potentially important and undocumented role of northern peatlands in
41
42 724 the global N cycle. These results also raise the important question of N provenance: in the
43
44 725 absence of large rates of atmospheric N deposition during the early Holocene, the only
45
46 726 process that could account for such a large N pool in peatlands is N fixation, either
47
48
49
50
51 727 through symbiotic or asymbiotic processes (Limpens et al., 2006).
52
53
54
55
56
57
58
59
60

728

729 The fate of these large peat N stocks remains largely unknown under recent and projected
730 warming. Indeed, the importance of peatlands as sources of nitrous oxide (N₂O) is just
731 emerging (e.g., Repo et al., 2009; Marushchak et al., 2011; Palmer et al., 2012), and
732 studies have suggested that reduced surface moisture or increasing temperatures might
733 significantly promote the production, transformation, and transport of dissolved N, and
734 N₂O emissions to the atmosphere through denitrification (e.g., Kane et al., 2010). On the
735 contrary, some authors have speculated that the potential increase in peatland-N₂O
736 emissions from climate change may not be significant relative to the global N₂O budget
737 (e.g., Martikainen et al., 1993; Frolking et al., 2011). Overall, additional peat N cycling
738 studies are needed to address these remaining questions.

739

740 **Future directions**

741 Peat core analysis has been extensively used over the past 20 years for estimating rates of
742 peat C accumulation at local, regional, and continental scales (e.g., Mäkilä, 1997; Clymo
743 et al., 1998; Vitt et al., 2000; Turunen et al., 2002; Mäkilä and Saarnisto, 2008; Yu et al.,
744 2010; van Bellen et al., 2011; Gorham et al., 2012). The present study analyzed a new
745 database that comprises 268 peat records from 215 northern peatland sites. This
746 systematic analysis of peat properties and Holocene C accumulation rates is essential for
747 accurately addressing the following general research topics in the future: (1) describing
748 and quantifying spatial and temporal patterns of Holocene peatland C and N
749 accumulation; (2) assessing the sensitivity of C and N accumulation to climate change;
750 (3) estimating peatland soil organic carbon (SOC) and soil organic nitrogen (SON) pools

1
2
3 751 at regional and hemispheric scales, (4) furthering our understanding of peatland C cycle –
4
5 752 climate linkages, and (5) providing the scientific community with a large dataset for
6
7
8 753 developing and testing earth system and ecological models.
9

10 754

11
12
13 755 **Acknowledgements**

14
15 756 We acknowledge the peatland research community for sharing their datasets. The U.S.
16
17 757 NSF supported the synthesis work through grant ARC-1107981 to Lehigh University.
18
19 758 The collection and analysis of unpublished records used in this synthesis were supported
20
21 759 by the following funding agencies and research grants: Alaska (NSF ARC-1107981,
22
23 760 AGS-0628455, and EAR-0819717; USGS Climate Research and Development Program),
24
25 761 Canada (NSF ARC-1107981, EAR-0223271, EAR-0843685, and AGS-0628598; NSERC
26
27 762 CRDPJ-305605, CRDPJ-365867; Hydro-Québec), Fennoscandia and Western Siberia
28
29 763 (NSF OPP-9818496; Academy of Finland 201321 and 1133515; University of Helsinki),
30
31 764 Kamchatka (NSF ARC-1107981, ARC-1108116), and the United Kingdom (Yorkshire
32
33 765 Peat Partnership). Lehigh University’s Library and Technology Services staff is
34
35 766 acknowledged for its support in building the web interface for the peatland database.
36
37 767 Finally, comments from Paul Glaser and two other journal reviewers improved the
38
39 768 overall quality of the manuscript.
40
41
42
43
44
45

46 769

47
48 770 **References**

49
50 771 Armentano TV and Menges ES (1986) Patterns of change in the carbon balance of
51 772 organic soil-wetlands of the temperate zone. *Journal of Ecology* 74: 755-774.
52
53 773
54 774 Bauer IE, Bhatti JS, Cash KJ, Tarnocai, C and Robinson SD (2006) Developing statistical
55 775 models to estimate the carbon density of organic soils. *Canadian Journal of Soil Science*
56 776 86: 295-304.
57
58
59
60

- 777
778 Beilman DW, MacDonald GM, Smith LC and Reimer PJ (2009) Carbon accumulation in
779 peatlands of West Siberia over the last 2000 years. *Global Biogeochemical cycles* 23,
780 GB1012. doi: 10.1029/2007GB003112.
781
- 782 Belyea LR (2009) Non-linear dynamics of peatlands and potential feedbacks on the
783 climate system. In: Baird A, Belyea L, Comas X, Reeve A. and Slater L (Eds.), *Northern*
784 *peatlands and carbon cycling*, American Geophysical Union Monograph Series,
785 Washington D.C., USA, pp. 5-18.
786
- 787 Berger A and Loutre M-F (1991) Insolation values for the climate of the last 10 million
788 years. *Quaternary Science Reviews* 10: 297-317.
789
- 790 Bergner K, Albano Å and Bohlin E (1990) The content of peat: a compilation of
791 botanical, physical and chemical data of peat. Department of Agricultural Research,
792 Northern Sweden, Swedish University of Agricultural Sciences.
793
- 794 Björck S and Clemmensen LB (2004) Aeolian sediment in raised bog deposits, Halland,
795 SW Sweden: a new proxy record of Holocene winter storminess in southern Scandinavia?
796 *The Holocene* 14: 677-688.
797
- 798 Blodau C (2002) Carbon cycling in peatlands - A review of processes and controls,
799 *Environmental Reviews* 10: 111-134.
800
- 801 Bragazza L, Buttler A, Habermacher J, Brancaloni L, Gerdol R, Fritze H, Hanajik P,
802 Laiho R and Johnson D (2012) High nitrogen deposition alters the decomposition of bog
803 plant litter and reduces carbon accumulation. *Global Change Biology* 18: 1163-1172.
804
- 805 Bridgman SC, Megonigal JP, Keller JK, Bliss, NB and Trettin C (2006) The carbon
806 balance of North American wetlands. *Wetlands* 26: 889-916.
807
- 808 Buffam I, Carpenter SR, Yeck W, Hanson PC and Turner MG (2010) Filling holes in
809 regional carbon budgets: Predicting peat depth in a north temperate lake district. *Journal*
810 *of Geophysical Research – Biogeosciences* 115, G01005, doi:10.1029/ 2009JG001034.
811
- 812 Cagnon B, Py X, Guillot A, Stoeckli F and Chambat G (2009) Contributions of
813 hemicellulose, cellulose and lignin to the mass and the porous properties of chars and
814 steam activated carbons from various lignocellulosic precursors. *Bioresource Technology*
815 100: 292-298.
816
- 817 Camill P (1999) Peat accumulation and succession following permafrost thaw in the
818 boreal peatlands of Manitoba, Canada. *Ecoscience* 6: 592-602.
819
- 820 Chambers FM, Beilman DW and Yu Z (2011) Methods for determining peat
821 humification and for quantifying peat bulk density, organic matter and carbon content for
822 palaeostudies of climate and peatland carbon dynamics. *Mires and Peat* 7, article 7, 10 p.

823
824 Charman D, Beilman D, Blaauw M, Booth RK, Brewer S, Chambers F, Christen JA,
825 Gallego-Sala AV, Harrison SP, Hughes PDM, Jackson S, Korhola A, Mauquoy D,
826 Mitchell F, Prentice IC, van der Linden M, De Vleeschouwer F, Yu Z, Alm J, Bauer IE,
827 McCorish Y, Garneau M, Hohl V, Huang Y, Karofeld E, Le Roux G, Loisel J, Moschen
828 R, Nichols JE, Nieminen TM, MacDonald GM, Phadtare NR, Rausch N, Sillasoo Ü,
829 Swindles GT, Tuittila E-S, Ukonmaanaho L, Välranta M, van Bellen S, van Geel B, Vitt
830 D and Zhao Y (2013) Climate-related changes in peatland carbon accumulation during
831 the last millennium. *Biogeosciences* 10: 929-944. doi: 10.5194/bg-10-929-2013.
832
833 Clymo RS (1984) The limits to peat growth. *Philosophical Transactions of the Royal*
834 *Society of London, Series B, Biological Sciences* 303: 605-654.
835
836 Clymo RS, Turunen J and Tolonen K (1998) Carbon accumulation in peatlands. *Oikos*
837 81: 368-388.
838
839 Dean Jr WE (1974) Determination of carbonate and organic matter in calcareous
840 sediments and sedimentary rocks by loss on ignition: comparison with other methods.
841 *Journal of Sedimentary Petrology* 44: 242-248.
842
843 Frohking S and Roulet NT (2007) Holocene radiative forcing impact of northern peatland
844 carbon accumulation and methane emissions. *Global Change Biology* 13: 1-10.
845
846 Frohking S, Talbot J, Jones MC, Treat CC, Kauffman JB, Tuittila E-S and Roulet N
847 (2011) Peatlands in the Earth's 21st century coupled climate-carbon system.
848 *Environmental Reviews* 19: 371-396.
849
850 Glaser PH, Volin JC, Givnish TJ, Hansen BCS and Stricker CA (2012) Carbon and
851 sediment accumulation in the Everglades (USA) during the past 4000 years: rates,
852 drivers, and sources of error. *Journal of Geophysical Research-Biogeosciences* 117,
853 G03026. doi:10.1029/2011JG001821.
854
855 Gorham E (1990) Biotic impoverishment in northern peatlands. In: Woodwell GM (Ed.),
856 *The Earth in transition: Patterns and processes of biotic impoverishment*. Cambridge
857 University Press, New York, USA, pp. 65-98.
858
859 Gorham E (1991) Northern peatlands: role in the carbon cycle and probable responses to
860 climatic warming. *Ecological Applications* 1: 182-195.
861
862 Gorham E, Lehman C, Dyke A, Janssens J and Dyke L (2007) Temporal and spatial
863 aspects of peatland initiation following deglaciation in North America. *Quaternary*
864 *Science Reviews* 26: 300-311.
865
866 Gorham E, Lehman C, Dyke A, Clymo D and Janssens J. (2012) Long-term carbon
867 sequestration in North American peatlands. *Quaternary Science Reviews* 58: 77-82.
868

- 869 Harden JW, Sundquist ET, Stallard RF and Mark RK (1992) Dynamics of soil carbon
870 during deglaciation of the Laurentide ice sheet. *Science* 258: 1921-1924.
871
- 872 Heiri O, Lotter AF and Lemcke G (2001) Loss on ignition as a method for estimating
873 organic and carbonate content in sediments: reproducibility and comparability of results.
874 *Journal of Paleolimnology* 25: 101-110.
875
- 876 Hughes PDM (2000) A reappraisal of the mechanisms leading to ombrotrophy in British
877 raised mires. *Ecology Letters* 3: 7-9.
878
- 879 Ireland AW, Booth RK, Hotchkiss SC and Schmitz JE (2013) A comparative study of
880 within-basin and regional peatland development: implications for peatland carbon
881 dynamics. *Quaternary Science Reviews* 61: 85-95.
882
- 883 Jones MC and Yu Z (2010) Rapid deglacial and early Holocene expansion of peatlands in
884 Alaska. *Proceedings of the National Academy of Sciences* 107: 7347-7352.
885
- 886 Kane ES, Turetsky MR, Harden JW, McGuire AD and Waddington JM (2010) Seasonal
887 ice and hydrologic controls on dissolved organic carbon and nitrogen concentrations in a
888 boreal-rich fen. *Journal of Geophysical Research – Biogeosciences* 115, G04012. doi:
889 10.1029/2010JG001366.
890
- 891 Kaufman DS, Ager TA, Anderson NJ, Anderson PM, Andrews JT, Bartlein PJ, Brubaker
892 LB, Coats LL, Cwynar LC, Duvall ML, Dyke AS, Edwards ME, Eisner WR, Gajewski
893 K, Geirsdottir A, Hu FS, Jennings AE, Kaplan MR, Kerwin MW, Lozhkin AV,
894 MacDonald GM, Miller GH, Mock CJ, Oswald WW, Otto-Bliesner BL, Porinchu DF,
895 Ruhland KR, Smol JP, Steig EJ, Wolfey BB (2004) Holocene thermal maximum in the
896 western Arctic (0-180 °W). *Quaternary Science Reviews* 23: 529-560.
897
- 898 Korhola A, Ruppel M, Seppä H, Välranta M, Virtanen T and Weckström J (2010) The
899 importance of northern peatland expansion to the late-Holocene rise of atmospheric
900 methane. *Quaternary Science Reviews* 29: 611-617.
901
- 902 Lappalainen E (1996) General review on world peatlands and peat resources. In:
903 Lappalainen E (Ed.), *Global Peat Resources*. International Peat Society, Jyska, pp. 53-56.
904
- 905 Limpens J, Heijmans MPD and Berendse F (2006) The nitrogen cycle in boreal
906 peatlands. In: Wieder RK and Vitt DH (Eds.), *Boreal Peatland Ecosystems*. Ecological
907 Studies Vol. 188, Springer-Verlag, Berlin Heidelberg, Germany, pp. 195-230.
908
- 909 MacDonald GM, Beilman DW, Kremenetski KV, Sheng Y, Smith LC and Valichko AA
910 (2006) Rapid early development of circum-arctic peatlands and atmospheric CH₄ and
911 CO₂ variations. *Science* 314: 285-288.
912

- 913 Mäkilä M (1997) Holocene lateral expansion, peat growth and carbon accumulation on
 914 Haukkasuo, a raised bog in southeastern Finland. *Boreas* 26: 1-14. doi:10.1111/j.1502-
 915 3885.1997.tb00647.x.
 916
- 917 Mäkilä M and Saarnisto M (2008) Carbon accumulation in boreal peatlands during the
 918 Holocene – impacts of climate variations. In: Strack, M. (Ed.), *Peatlands and climate*
 919 *change*. International Peat Society, Jyväskylä, Finland, pp. 24-43.
 920
- 921 Maltby E and Immirzi P (1993) Carbon dynamics in peatlands and other wetland soils,
 922 regional and global perspectives. *Chemosphere* 27: 999-1023.
 923
- 924 Marcott SA, Shakun JD, Clark PU and Mix AC (2013) A reconstruction of regional and
 925 global temperature for the past 11,300 years. *Science* 339: 1198-1201.
 926
- 927 Martikainen PJ, Nykänen H, Crill P and Silvola J (1993) Effect of a lowered water-table
 928 on nitrous-oxide fluxes from northern peatlands. *Nature* 366: 51-53.
 929
- 930 Marushchak ME, Pitkämäki A, Koponen H, Biasi C, Seppälä M and Martikainen PJ
 931 (2011) Hot spots for nitrous oxide emissions found in different types of permafrost
 932 peatlands. *Global Change Biology* 17: 2601-2614.
 933
- 934 Mauquoy D and van Geel B (2007) Mire and Peat Macros. In: Elias S.A. (Ed.),
 935 *Encyclopedia of Quaternary Science*. vol. 3. Elsevier, Amsterdam, The Netherlands, pp.
 936 2315-2336.
 937
- 938 McGuire AD, Anderson LG, Christensen TR, Dallimore S, Guo L, Hayes DJ, Heimann
 939 M, Lorenson TD, MacDonald RW and Roulet N (2009) Sensitivity of the carbon cycle in
 940 the Arctic to climate change. *Ecological Monographs* 79: 523-555.
 941
- 942 McLaughlan KK, Williams JJ, Craine JM and Jeffers ES (2013) Changes in global
 943 nitrogen cycling during the Holocene epoch. *Nature* 495: 352-357.
 944
- 945 Oksanen PO, Kuhry P and Alekseeva RN (2003) Holocene development and permafrost
 946 history of the Usinsk Mire, northeast European Russia. *Géographie Physique et*
 947 *Quaternaire* 57: 169-187.
 948
- 949 Palmer K, Biasi C and Horn MA (2012) Contrasting denitrifier communities relate to
 950 contrasting N₂O emission patterns from acidic peat soils in arctic tundra. *The ISME*
 951 *Journal* 6: 1058-1077.
 952
- 953 Pelletier L, Moore TR, Roulet NT, Garneau M and Beaulieu-Audy V (2007) Methane
 954 fluxes from three peatlands in the La Grande Rivière watershed, James Bay lowland,
 955 Canada. *Journal of Geophysical Research – Biogeosciences* 112, G01018. doi:
 956 10.1029/2006JG000216.
 957
 958
 959
 960

- Peteet D, Andreev A, Bardeen W and Mistretta F (1998) Long-term Arctic peatland dynamics, vegetation and climate history of the Pur-Taz region, Western Siberia. *Boreas* 27: 115-126.
- Piotrowska N, Blaauw M, Mauquoy D and Chambers FM (2011) Constructing deposition chronologies for peat deposits using radiocarbon dating. *Mires and Peat* 7, article 10, 14 p.
- Post WM, Pastor J, Zinke PJ and Stangenberger AG (1985) Global patterns of soil nitrogen storage. *Nature* 317: 613-616.
- Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Bertrand C, Blackwell PG, Buck CE, Burr G, Cutler KB, Damon PE, Edwards RL, Fairbanks RG, Friedrich M, Guilderson TM, Hughen KA, Kromer B, McCormac FG, Manning S, Bronk Ramsey C, Reimer RW, Remmele S, Southon JR, Stuiver M, Talamo S, Taylor FW, van der Plicht J, Weyhenmeyer CE (2009) IntCal09 and Marine09 radiocarbon age calibration curves, 0–50,000 years cal BP. *Radiocarbon* 51: 1111-1150.
- Renssen H, Seppä H, Crosta X, Goose H and Roche DM (2012) Global characterization of the Holocene Thermal Maximum. *Quaternary Science Reviews* 48: 7-19.
- Repo ME, Susiluoto S, Lind SE, Jokinen S, Elsakov V, Biasi C, Virtanen T, Martikainen PJ (2009) Large N₂O emissions from cryoturbated peat soil in tundra. *Nature Geoscience* 2: 189-192.
- Ruppel M, Väiranta M, Virtanen T and Korhola A (2013) Postglacial spatiotemporal peatland initiation and lateral expansion dynamics in North America and northern Europe. *The Holocene*. doi:10.1177/0959683613499053.
- Rydin H and Jeglum J (2013). *The biology of peatlands*. 2nd edition, Oxford University Press, Oxford, UK, 382 p.
- Sannel ABK and Kuhry P (2008) Long-term stability of permafrost in subarctic peat plateaus, west-central Canada. *The Holocene* 18: 589-601.
- Sheng Y, Smith LC, MacDonald GM, Kremenetski KV, Frey KE, Velichko AA, Lee M, Beilman DW and Dubinin P (2004) A high-resolution GIS-based inventory of the West Siberian peat carbon pool. *Global Biogeochemical Cycles* 18, GB3004. doi: 10.1029/2003GB002190.
- Smith, L.C., MacDonald, G.M., Velichko, A.A., Beilman, D.W., Borisova, O.K., Frey, K.E., Kremenetski, K.V., Sheng, Y. 2004. Siberian peatlands a net carbon sink and global methane source since the Early Holocene. *Science*, 303, 353–356.
- Stolbovoi V (2002) Carbon in Russian soils. *Climatic Change* 55: 131-156.

1
2
3 1004 Stolbovoi V and McCallum I (2002) *Land resources of Russia*. International Institute for
4 1005 Applied Systems Analysis. www.iiasa.ac.at/Research/FOR/russia_cd/download.htm.
5 1006
6
7 1007 Stolbovoi V, Montanarella L, Medvedev V, Smeyan N, Shishov L, Ungureanu V,
8 1008 Dobrovolski G, Jamagne M, King D, Rozhkov V and Savin I (2001) Integration of data
9 1009 on the soils of Russia, Byelorussia, Moldova and Ukraine into the Soil Geographic Database
10 1010 of the European Community. *Eurasian Soil Science* 34: 687-703.
11 1011
12 1012 Stuiver M and Reimer PJ (1993) Extended ¹⁴C database and revised CALIB radiocarbon
13 1013 calibration program. *Radiocarbon* 35: 215-230.
14 1014
15 1015 Telford RJ, Heegaard E and Birks HJB (2004) All age-depth models are wrong: but how
16 1016 badly? *Quaternary Science Reviews* 23: 1-5.
17 1017
18 1018 Troels-Smith J (1955) Characterization of unconsolidated sediments. Danmarks
19 1019 Geologiske *Undersøgelser* Series 4: 1-73.
20 1020
21 1021 Turetsky MR, Wieder RK, Vitt DH, Evans RJ and Scott KD (2007) The disappearance of
22 1022 relict permafrost in boreal North America: effects on peatland carbon storage and fluxes.
23 1023 *Global Change Biology* 13: 1922-1934.
24 1024
25 1025 Turunen J, Tomppo E, Tolonen K and Reinikainen A (2002) Estimating carbon
26 1026 accumulation rates of undrained mires in Finland – application to boreal and subarctic
27 1027 regions. *The Holocene* 12: 69-80.
28 1028
29 1029 Turunen J, Roulet NT, Moore TR and Richard PJH (2004) Nitrogen deposition and
30 1030 increased carbon accumulation in ombrotrophic peatlands in eastern Canada. *Global*
31 1031 *Biogeochemical Cycles*, 18, GB3002. doi: 10.1029/2003GB002154.
32 1032
33 1033 van Bellen S, Dallaire P-L, Garneau M and Bergeron Y (2011) Quantifying spatial and
34 1034 temporal Holocene carbon accumulation in ombrotrophic peatlands of the Eastmain
35 1035 region, Quebec, Canada. *Global Biogeochemical Cycles* 25, GB2016.
36 1036 doi:10.1029/2010GB003877.
37 1037
38 1038 Vitt DH, Halsey LA, Bauer IE and Campbell C (2000) Spatial and temporal trends in
39 1039 carbon storage of peatlands of continental western Canada through the Holocene.
40 1040 *Canadian Journal of Earth Sciences* 37: 683-693.
41 1041
42 1042 Yu Z (2011) Holocene carbon flux histories of the world's peatlands: global carbon-cycle
43 1043 implications. *The Holocene* 21: 761-774.
44 1044
45 1045 Yu Z (2012) Northern peatland carbon stocks and dynamics: a review. *Biogeosciences* 9:
46 1046 4071-4085.
47 1047
48
49
50
51
52
53
54
55
56
57
58
59
60

- Yu Z, Campbell ID, Campbell C, Vitt DH, Bond GC and Apps MJ (2003) Carbon sequestration in western Canadian peat highly sensitive to Holocene wet-dry climate cycles at millennial timescales. *The Holocene* 13: 801-808.
- Yu Z, Beilman DW and Jones MC (2009) Sensitivity of northern peatland carbon dynamics to Holocene climate change. In: Baird A, Belyea L, Comas X, Reeve A, Slater L (Eds.), *Northern peatlands and carbon cycling*. American Geophysical Union Monograph Series, Washington D.C., USA, pp. 55-69.
- Yu Z, Loisel J, Brosseau DP, Beilman DW and Hunt SJ (2010) Global peatland dynamics since the Last Glacial Maximum. *Geophysical Research Letters* 37, L13402. doi:10.1029/2010GL043584.
- Yu Z, Loisel J, Turetsky MR, Cai S, Zhao Y, Frolking S, MacDonald GM and Bubier JL (2013) Evidence for elevated emissions from high-latitude wetlands contributing to high atmospheric CH₄ concentration in the early Holocene. *Global Biogeochemical Cycles* 27: 131-140. doi:10.1002/GBC.20025.
- Zoltai SC (1971) Southern limit of permafrost features in peat landforms, Manitoba and Saskatchewan. *Geological Association of Canada, Special Paper* 9, 305-310.
- Zoltai SC (1995) Permafrost distribution in peatlands of west-central Canada during the Holocene warm period 6000 years B.P. *Géographie Physique et Quaternaire* 49: 45-54.

Figure and Table Captions

Figure 1. Location of study sites. Map showing the distribution of northern peatlands (green area from Yu et al., 2010) and peatland sites included in this study (n = 215 sites, including 268 peat cores). Long-term rate of peat-carbon accumulation was estimated from 127 sites (151 peat cores; red and blue dots). The yellow dots represent cores for which only peat properties (bulk density, organic matter content, etc.) were available and synthesized. Refer to Supplementary Table 1 for details.

Figure 2. Overview of data availability for North America (black bars) and Eurasia (white bars). (A) Number of cores (total = 238) containing information on carbon/nitrogen ratio (n = 40), nitrogen content (n = 40), carbon content (n = 56), peat

1
2
3 1083 types (n = 83), organic matter bulk density (n = 184), organic matter content (n = 190),
4
5 1084 and bulk density (n = 214). (B) Number of cores (total = 151) with a dating quality better
6
7 1085 than two dates per 1000 years (n = 35), one to two dates per 1000 years (n = 52), and less
8
9 1086 than one date per 1000 years (n = 64). (C) Number of calibrated basal peat ages (median)
10
11 1087 in 500-year bins from the database (n = 199) compared to all northern hemisphere basal
12
13 1088 peat ages (median) in 200-year bins (n = 2559, MGK data from MacDonald et al., 2006,
14
15 1089 Gorham et al., 2007, Korhola et al., 2010).

19
20 1090
21
22 1091 **Figure 3.** Distribution histograms of peat properties in northern peatlands. (A) Frequency
23
24 1092 distribution of bulk density for unidentified peat type samples (white bars) and different
25
26 1093 peat types (color bars). (B) Frequency distribution of organic matter content for different
27
28 1094 peat types. (C) Frequency distribution of organic matter bulk density for different peat
29
30 1095 types. (D) Frequency distribution of carbon content for different peat types. (E)
31
32 1096 Frequency distribution of nitrogen content for different peat types. (F) Frequency
33
34 1097 distribution of carbon/nitrogen mass ratio for different peat types.

38
39 1098
40
41 1099 **Figure 4.** Relation between carbon content and organic matter content in northern
42
43 1100 peatlands. The slope of each regression line is used as a conversion factor for estimating
44
45 1101 carbon content from organic matter content.

48
49 1102
50
51 1103 **Figure 5.** Temporal patterns of peat properties (mean, standard deviation, and number of
52
53 1104 samples). (A) Bulk density. (B) Organic matter content. (C) Organic matter bulk density.

(D) Carbon content. (E) Nitrogen content. (F) Carbon/Nitrogen mass ratio. White bars represent values that were based on a limited number of samples and peat records.

Figure 6. Main differences between *Sphagnum* and non-*Sphagnum* peat samples. (A) Bulk density. (B) Organic matter bulk density. (C) Organic carbon bulk density. (D) Carbon content. (E) Nitrogen content. (F) Carbon/Nitrogen mass ratio. (G) Temporal pattern of organic C bulk density. (H) Proportional change in the number of peat records that are *Sphagnum*-dominated, presented as a percentage of the total number of records.

Figure 7. Long-term apparent rate of carbon and nitrogen accumulation from northern peatlands (n = 127 sites). (A) Summer insolation at 60°N (data from Berger and Loutre, 1991) and temperature anomaly from an 11,300-year reconstruction for the northern extra-polar region from 30 to 90°N (data from Marcott et al., 2013). Temperature anomaly was calculated based on the 1961-1990 temperature averages. (B) Mean peat-carbon accumulation rates (PCAR) and standard error in 500-year bins. The number of sites per 500-year bins is also presented. (C) Mean peat-nitrogen accumulation rates (PNAR) and standard error in 500-year bins. These values were obtained using different C/N values over time, as indicated by the line.

Table 1. Peat properties in northern peatlands. Means and standard deviations are presented, along with the number of samples (n).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1127 **Table 2.** Northern peatland peat properties by regions. Means and standard deviations are
1128 presented, along with the number of samples (n).
1129
1130 **Table 3.** Comparison of northern peatland peat properties estimates with other published
1131 values. Means and standard deviations are presented, along with the number of samples
1132 (n) when available.
1133
1134 **Supplementary Material**
1135 **Table S1.** Summary information for the study sites included in the database.

1 A ~~Circum-Arctic~~ database and synthesis of northern peatland soil properties and

2 Holocene carbon and nitrogen accumulation

4 Running title

5 ~~Circum-Arctic-Northern~~ peatland database and synthesis

7 Authors

8 Julie Loisel^{1*}, Zicheng Yu^{1*}, David W. Beilman², Philip Camill³, Jukka Alm⁴, Matthew J.

9 Amesbury⁵, David Anderson⁶, Sofia Andersson⁷, Christopher Bochicchio¹, Keith Barber⁸,

10 Lisa R. Belyea⁹, Joan Bunbury¹⁰, Frank M. Chambers¹¹, Daniel J. Charman⁵, François De

11 Vleeschouwer¹², Barbara Fiałkiewicz-Kozieł¹³, Sarah A. Finkelstein¹⁴, Mariusz Galka¹³,

12 Michelle Garneau¹⁵, Dan Hammarlund¹⁶, William Hinchcliffe⁵, James Holmquist¹⁷, Paul

13 Hughes⁸, Miriam C. Jones¹⁸, Eric S. Klein¹, Ulla Kokfelt¹⁹, Atte Korhola²⁰, Peter Kuhry⁷,

14 Alexandre Lamarre¹⁵, Mariusz Lamentowicz¹³, David Large²¹, Martin Lavoie²², Glen

15 MacDonald¹⁷, Gabriel Magnan¹⁵, Markku Mäkilä²³, Gunnar Mallon⁸, Paul Mathijssen²⁰,

16 Dmitri Mauquoy²⁴, Julia McCarroll¹¹, Tim R. Moore²⁵, Jonathan Nichols²⁶, Benjamin

17 O'Reilly¹⁴, Pirita Oksanen²⁷, Maara Packalen²⁸, Dorothy Peteet²⁶, Pierre J.H. Richard²⁹,

18 Stephen Robinson³⁰, Tiina Ronkanen²⁰, Mats Rundgren¹⁶, A. Britta K. Sannel⁷, Charles

19 Tarnocai³¹, Tim Thom³², Eeva-Stiina Tuittila⁴, Merritt Turetsky³³, Minna Väliranta²⁰,

20 Marjolein van der Linden³⁴, Bas van Geel³⁵, Simon van Bellen²³, Dale Vitt³⁶, Yan Zhao³⁷,

21 Weijian Zhou³⁸

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Revised manuscript submitted on 19 December 2013 as a Research Paper for the special issue *Holocene Peatland Carbon Dynamics in the Circum-Arctic Region*.

Affiliations

¹Department of Earth and Environmental Sciences, Lehigh University, Bethlehem, PA 18015, USA

²Department of Geography, University of Hawaii – Manoa, Honolulu, HI 96822, USA

³Department of Earth and Oceanographic Sciences, Bowdoin College, Brunswick, ME 04011, USA

⁴School of Forest Sciences, University of Eastern Finland, Joensuu, FI 80101, Finland

⁵Department of Geography, University of Exeter, Exeter, EX4 4RJ, UK

⁶Department of Geography, Eton College, Windsor, Berkshire SL4 6DW, UK

⁷Department of Physical Geography and Quaternary Geology, Stockholm University, Stockholm, 106 91, Sweden

⁸Geography and Environment, University of Southampton, Southampton, SO17 1BJ, UK

⁹School of Geography, Queen Mary University of London, London, E1 4NS, UK

¹⁰Department of Geography and Earth Science, University of Wisconsin – La Crosse, La Crosse, WI 54601, USA

¹¹Centre for Environmental Change and Quaternary Research, University of Gloucestershire, Cheltenham, GL50 4AZ, UK

¹²CNRS and Université de Toulouse, Castanet Tolosan, 31326, France

¹³Department of Biogeography and Paleoecology, Adam Mickiewicz University, Poznan, 61-680, Poland

¹⁴Department of Earth Sciences, University of Toronto, Toronto, ON M5S 3B1, Canada

¹⁵Departement de Géographie and GEOTOP, Université du Québec – Montréal, Montréal, QC H3C 3P8, Canada

¹⁶Department of Geology, Lund University, Lund, SE-223 62, Sweden

- 65
66 ¹⁷Department of Geography, University of California – Los Angeles, Los Angeles, CA
67 90095, USA
68
69 ¹⁸U.S. Geological Survey, Reston, VA 20192, USA
70
71 ¹⁹Department of Geosciences and Natural Resource Management, University of
72 Copenhagen, Copenhagen, DK-1350, Denmark
73
74 ²⁰Department of Environmental Sciences, University of Helsinki, Helsinki, FIN-00014,
75 Finland
76
77 ²¹Department of Chemical and Environmental Engineering, University of Nottingham,
78 Nottingham, NG7 2RD, UK
79
80 ²²Département de Géographie and Centre d'études nordiques, Université Laval, Québec,
81 QC G1V 0A6, Canada
82
83 ²³[Geological Survey of Finland, P.O. Box 96, Espoo, 02151, Finland](#)
84
85 ²⁴School of Geosciences, University of Aberdeen, Aberdeen, AB24 3UF, UK
86
87 ²⁵Department of Geography and Global Environmental and Climate Change Centre,
88 McGill University, Montreal, QC H3A 0B9, Canada
89
90 ²⁶Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964, USA
91
92 ²⁷Centre for Economic Development, Transport and the Environment, Vaasa, 65101,
93 Finland
94
95 ²⁸[Department of Geography, University of Toronto, Toronto, ON M5S 3G3, Canada](#)
96
97 ²⁹Département de Géographie, Université de Montréal, Montréal, QC H2V 2B8, Canada
98
99 ³⁰Champlain College – Dublin Campus, Dublin, Ireland
100
101 ³¹[Agriculture and Agri-Food Canada, Ottawa, ON K1A 0C6, Canada](#)
102
103 ³²Yorkshire Peat Partnership, Yorkshire Wildlife Trust, York, YO24 1GN, UK
104
105 ³³Department of Integrative Biology, University of Guelph, Guelph, ON N1G 2W1,
106 Canada
107
108 ³⁴BIAX Consult, Zaandam, 1506 AL, The Netherlands
109

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

³⁵Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam,
Amsterdam, 1098 XH, The Netherlands
³⁶Department of Plant Biology, Southern Illinois University, Carbondale, IL 62901, USA
³⁷Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of
Sciences, Beijing, 100101, China
³⁸Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, 710075 Shaanxi,
China

Corresponding authors

Julie Loisel email: jul208@lehigh.edu phone: 610-758-3660
Zicheng Yu email: ziy2@lehigh.edu phone: 610-758-6751

Abstract

Here we present results from the most comprehensive compilation of Holocene peat soil
properties ~~withand associated Holocene~~ carbon and nitrogen accumulation rates for
northern peatlands. Our database consists of 268 peat cores from 215 sites located north
of 45°N. It encompasses regions within which peat carbon data have only recently
become available, such as the West Siberia Lowlands, ~~and~~ the Hudson Bay Lowlands,
Kamchatka in Far East Russia, and the Tibetan Plateau. For all northern peatlands, carbon
content in organic matter was estimated at $42.3 \pm 3\%$ (S.D.) for *Sphagnum* peat, $51.3 \pm$
 2% for non-*Sphagnum* peat, and at $49.2 \pm 2\%$ overall. Dry bulk~~Bulk~~ density averaged
 $0.12 \pm 0.07 \text{ g cm}^{-3}$, organic matter bulk density averaged $0.11 \pm 0.05 \text{ g cm}^{-3}$, and total
carbon content in peat averaged $46.847 \pm 6.4\%$. In general, large differences were found
between *Sphagnum* and non-*Sphagnum* peat types in terms of peat properties. Time-
weighted peat ~~-~~carbon accumulation rates averaged 232.9 ± 2 (S.E.M.) $\text{g C m}^{-2} \text{ yr}^{-1}$

during the Holocene on the basis of 151 peat cores from 127 sites, with the highest rates of carbon accumulation ($25\text{--}28\text{ g C m}^{-2}\text{ yr}^{-1}$) recorded during the ~~warmer than today~~ early Holocene when the climate was warmer than the present. Furthermore, Finally, we ~~provide the first~~ estimate ~~for~~ the northern peatland carbon and nitrogen pools at 436 and 109.7 gigatons, respectively equivalent to 10% of the world's soil nitrogen. The database is publicly available at <https://www.peatlands.lehigh.edu>.

Keywords

Northern peatlands, Data synthesis, Climate change, Soil carbon and nitrogen, Biogeochemical cycles, Long-term ecosystem dynamics

Introduction

Of all terrestrial ecosystems, peatlands are arguably the most efficient at sequestering carbon (C) over long time scales. Northern peatlands cover approximately $4,000,000\text{ km}^2$ or 3% of the global land area (Maltby and Immirzi, 1993) and have accumulated about 500 gigatons of C (GtC) mostly during the Holocene, equivalent to $\sim 30\%$ of the present-day the global soil organic carbon (SOC) pool (Gorham, 1991; Bridgham et al., 2006; Yu et al., 2010). These ecosystems have also ~~been playing~~played a dynamic role in the Holocene global C cycle as important sinks of carbon dioxide (CO_2) and major sources of methane (CH_4) to the atmosphere (Frolking and Roulet, 2007; Korhola et al., 2010; Yu, 2011). As climate warming positively affects both plant growth and organic matter decomposition, rRecent and projected climate change could shift the balance between peat production and organic matter decomposition, potentially affecting the peatland C-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

sink capacity and modifying peat ~~C~~ fluxes to the atmosphere (Frolking et al., 2011; Yu, 2012). This ~~prediction is~~ particularly ~~the case in~~ holds true for the northern high-latitude regions, where the intensity of climate change is expected to be greatest (McGuire et al., 2009). The peatland ~~C cycle carbon~~ climate feedback remains difficult to assess, however, because of (1) limited understanding of peatland responses to climate change (Frolking et al., 2011), (2) data gaps and large uncertainties in regional peatland C stocks (Yu, 2012), and (3) non-linear peatland responses to external forcing (Belyea, 2009).

Very little is known about the nitrogen (N) budget that accompanies C accumulation in northern peatlands (but see Limpens et al., 2006 for a review). Assuming a net C sequestration of ~270 GtC (Yu, 2012) and a C/N ratio of 20-30 for fen peat (Bergner et al., 1990; Rydin and Jeglum, 2013; results therein) during the early Holocene (11-7 ka), about 10-13 GtN would have been required to build such peat deposits. It is therefore possible that northern peatlands have been playing an undocumented, dynamic role in the Holocene global N cycle as important sinks of N, potentially limiting the amount of N available for other ecosystems at the global scale (McLauchlan et al., 2013). Alternatively, if the main N input to peatlands was through N₂ fixation by cyanobacteria, these microorganisms might have been more important in driving the C cycle in peatlands than previously thought. Overall, studying the coupling between N and C cycling in northern peatlands is essential for a better understanding of how key biogeochemical processes interact in these systems and for predicting the future of peat C stocks.

Here we present the most comprehensive compilation of Holocene C and N data for
 northern peatlands. ~~Our database consists of 268 peat cores from 215 sites located north
 of 45 °N or at high elevations (e.g., Tibetan Plateau).~~ This synthesis encompasses regions
 within which peat -C and N data have only recently become available, such as the West
 Siberian Lowlands in Asian Russia, ~~a and~~ the Hudson Bay Lowlands in Canada,
Kamchatka in the Russian Far East, and the Tibetan Plateau. In addition, we present the
 most comprehensive synthesis of peat soil properties (such as bulk density, organic
 matter content, ~~and carbon-C~~ and ~~nitrogen-N~~ content) from the northern hemisphere.
Also, this new database and synthesis work represent a major expansion from Yu et al.'s
 (2009) synthesis on Holocene peat C dynamics, which was based on 33 sites (vs. 127
 sites as reported in this paper). Finally, it constitutes a natural continuation of Charman et
 al.'s (2013) recent study on peat C accumulation in northern peatlands during the last
 millennium.
In addition to filling regional data and knowledge gaps, The main objectives of this
 paper are to (1) describe a database of peat soil properties and synthesize this information
 for different peat types, time intervals, and geographic regions, and (2) produce ~~a~~ time
 series of Holocene peat -C and N accumulation rates in 500-yr ~~increments-bins~~ for
 comparison with climate history. Key differences in Holocene peat properties between
 different regions and peatland types are also discussed in light of their implications for
 long-term peat C stocks. Of particular importance and relevance are the differences
 between *Sphagnum* and non-*Sphagnum* peat types. Finally, we present new estimates for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

northern peat C and N stocks for northern peatlands on the basis of the expanded
database.

Database and ~~Database and analytical methods~~ analysis
Database

We compiled a dataset of 268 published and unpublished Holocene peat records from 215 sites located in North America and Eurasia (Figure 1, Supplementary Table 1). The difference in the number of peat cores and peatland sites is due to the fact that, in a few instances, multiple cores were collected from a single peatland. As these multiple cores were not designed as true replicates in the original publications, each of these cores was considered as an independent record of peat properties in the present study. However, only the oldest core for each site was ~~accounted used~~ for ~~when~~ estimating peat inception age. Finally, when calculating peat ~~-C~~ carbon accumulation rates, ~~estimates, multiple~~ cores from a single site were each attributed an equal fraction of the weight for that site. ~~For example, for~~ For a site with three cores, the peat ~~-C~~ accumulation history of each core only accounted for 1/3 of ~~a record~~ the site's record.

The ~~H~~ latitude of most peatland sites ranged from 45 to 69 °N. The cutoff at 45 °N represents the southern limit for defining what is considered to be the area contributing to the C cycle of the Arctic region (McGuire et al., 2009). ~~with the exception of Four high-elevation~~ three high-elevation sites found in China (the Tibetan Plateau) and Japan were also included, as they developed under similar 'northern' climatic conditions. ~~The name and coordinates of these four sites are as follows: Zoige (33.5 °N, 102.6 °E), Hongyuan~~

(32.8 °N, 102.5 °E), Hani (42.2 °N, 126.5 °E), and Utsai (42.4 °N, 140.2 °E). -A total of 155 cores originate from Eurasia, ~~comprising including~~ 112 cores from Russia. The remaining 113 cores come from North America, ~~including 12 cores from Alaska~~. Approximately 40% of all cores were collected ~~from in~~ ombrotrophic bogs (n = 110 ~~cores~~⁰⁸) and 20% were extracted from minerotrophic fens (n = 50 ~~cores~~). The remainder (40%) was collected in ~~permafrost~~ peatlands ~~currently affected by permafrost~~ (n = 108 ~~cores~~⁴⁰). Note that peatland type was identified independently for each site by the original investigators. From an ecosystem functioning perspective, distinguishing bogs from fens and permafrost peatlands is important, as these peatland types are characterized by different hydrological regimes, vegetation communities, and peat-growth trajectories, all of which impact long-term rates of peat ~~-~~C sequestration (Rydin and Jeglum, 2013~~2006~~). Bogs are ~~nutrient~~^{mineral}-poor, rain-fed peatland ecosystems with relatively low plant net primary production (NPP) and slow peat decomposition rates. In contrast, fens are hydrologically connected to surface or ground water, thereby receiving ~~more nutrients-mineral nutrients from the groundwater~~. Generally speaking, ~~they fens~~ have greater NPP but also faster peat decay rates than bogs (Blodau, 2002). Finally, in the sub-~~A~~arctic and ~~A~~arctic regions, peatland hydrology, structure, and peat ~~-~~C balance are sensitive to the underlying permafrost aggradation and degradation dynamics (Camill, 1999; Turetsky et al., 2007). For the analysis, ~~frozen~~ peat plateaus (87~~9~~ out of 108~~40~~ cores), permafrost bogs (18 out of 108~~40~~ cores), and collapse scars (3 out of 108~~40~~ cores) were grouped under the peatland type 'permafrost peatlands'. Original peatland categories can be found in Supplementary Table 1.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The database was built to include as many peat records as possible. Therefore, we included any peat core that was extracted north of 45 °N (or at high elevation) and for which bulk density or organic matter bulk density data were available. Information related to peat-core location, peatland type, peat properties, age, and data source can be found in Supplementary Table 1. Additional information related to the type of coring device used and the year of coring can be found in the original publications. Data ~~and age depth relationships~~ used in this synthesis are readily accessible from the Holocene Peatland Carbon Network website (<https://www.peatlands.lehigh.edu>). ~~The website includes all published records and will be updated as new records included in this synthesis are eventually published.~~ This database ~~should~~ will be useful in future studies of ecosystem – ~~carbon~~ C cycle – climate interactions and for modeling long-term peatland dynamics.

In the following sub-sections, we present the criteria for site selection and the protocols used to develop the ~~data~~ dataset. In an effort to only analyze and synthesize *peat* samples, inorganic-rich horizons often found at the base of the peat cores were removed from the database. When available, stratigraphic information was used to distinguish peat vs. non-peat material. For example, gyttja (organic-rich lake sediments) was excluded from the dataset, as well as marshy, clayey, and silty sediments. When stratigraphic information was not available, a bulk density value of 0.5 g cm⁻³ was used as a cut-off between peat and non-peat material. This value was chosen on the basis of stratigraphic information from peatland records where peaty sediments with bulk density values up to 0.5 g cm⁻³ were identified. We ~~do~~ acknowledge that this cut-off value is arbitrary, and that our

dataset likely contains some non-peat samples. Inorganic horizons (e.g., tephra layers)

were also excluded from the ~~peat records~~ database.

Peat properties

A ~~subset-total~~ of 232 peat cores (181 sites) ~~were-was~~ used for characterizing peat

properties, though not all cores have all types of peat properties available (Figure 2a).

This dataset contains 139 cores from Eurasia (including 109 cores from Russia) and 93

cores from North America (Figure 1). While approximately half of these cores ~~were-as~~

sampled and analyzed at high resolution (1-5 cm increments), the ~~remainder-wremainder~~

~~was-as~~ sampled at lower resolution, typically at 10 cm increments. Dry b Bulk density

(BD; g cm⁻³), organic matter content (OM%; gravimetric %), as well as elemental C and

N concentration values were compiled and synthesized. On the basis of these raw

datasets, ~~carbon-to-nitrogen-ratio (C/N mass ratio)~~ and organic matter bulk density

(OMBD; g OM cm⁻³) were calculated. These peat geochemical values are examined in

light of peat stratigraphy, peat ages, and geographic regions. The following paragraphs

briefly describe the protocols used to obtain these values.

Peat stratigraphic information was obtained for 83 peat cores ('peat types' in Figure 2a)

for which plant macrofossil analysis or detailed peat description had been performed

following standard techniques (e.g., Troels-Smith, 1955; Mauquoy and van Geel, 2007).

This peat stratigraphic information was condensed into the following five peat types:

Sphagnum, herbaceous, woody, brown moss, and humified peat. In a few cases, the

investigators only ascribed a general peat type to the samples (e.g., 'bog' vs. 'fen' peat, or

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

298 ‘*Sphagnum*’ vs. ‘herbaceous’ peat). In these cases, the uncertainty associated with
299 classifying peat samples mostly relates to the uniformity of naming convention used
300 among the investigators. For example, a peat sample that contains sizable fractions of
301 brown moss and humified peat may be classified by an investigator as ‘brown moss peat’
302 and by another one as ‘humified peat’. We recognize that, due to their nature, brown
303 moss and humified peat types might be less uniform than *Sphagnum*, ~~or~~ herbaceous, ~~or~~
304 ~~woody~~ peat types. We included as much stratigraphic information as possible in the
305 database, though ambiguous or imprecise descriptions were left out to avoid further
306 ~~uncertainties~~confusions.
307
308 Dry bulk density and organic matter concentration content were determined following
309 standard procedures; in all cases where these parameters were measured, were
310 determined following standard procedures (Dean, 1974; Chambers et al., 2011). Peat
311 samples of a known fresh volume were either freeze-dried or oven-dried at ca. 100 °C
312 until constant weight was reached and weighed to determine bulk density, then burned at
313 500-600 °C for one to four hours and weighed again to determine organic matter content.
314 The accuracy of these measurements mostly depends on sample handling (care must be
315 taken to prevent peat compaction in the field and in the laboratory) and the analytical
316 error associated with weighing. The product of bulk density (BD; g cm⁻³) and organic
317 matter content (OM%) of each peat sample was used to calculate organic matter bulk
318 density (OMBD; g OM cm⁻³), also referred to as ash-free bulk density (AFBD) or organic
319 bulk density (OBD) in the literature (Yu et al., 2003; Björck and Clemmensen, 2004). We
320 compiled a total of 21,220 bulk density measurements ~~from 214 cores~~, 18,973 organic

321 matter content values ~~from 190 cores~~, and computed 18,544 organic matter bulk density
 322 values ~~from 184 cores~~ (Figure 2a).

324 Total peat ~~carbon-C~~ and ~~nitrogen-N~~ content were directly measured by combustion and
 325 elemental analysis of dry peat samples (Chambers et al., 2011). We compiled 3741 ~~C and~~
 326 ~~3365 carbon-N~~ measurements ~~from 56 cores and 3365 nitrogen content measurements~~
 327 ~~from 40 cores~~ (Figure 2a). We also computed a total of 3362 ~~carbon to nitrogen ratio~~
 328 ~~(C/N) mass ratio~~ values ~~from 40 cores~~.

330 Finally, the regression between ~~total~~ peat ~~carbon-C~~ content (C%) and peat organic matter
 331 content (OM%) for each peat type is presented as an estimate for ~~carbon-C~~ content in
 332 organic matter (OC%). A total of 995 samples ~~from 19 cores~~ were used in this analysis.
 333 The slope of each one of these regressions is interpreted as the 'conversion factor' from
 334 OM% to OC%, such that it provides an indirect way for estimating the C% content of
 335 ash-free peat for investigators who do not perform elemental C% measurements directly.

337 Peat-core chronology

338 Peat-core chronologies were almost exclusively based on radiocarbon (^{14}C) dates that
 339 were determined mostly by accelerator mass spectrometry (AMS) on terrestrial plant
 340 macrofossils or bulk peat (Piotrowska et al., 2011). A few older chronologies were based
 341 on conventional ^{14}C dating of bulk samples. Because no systematic offset has been
 342 observed in the ^{14}C age of bulk vs. non-woody plant macrofossils ~~The use of bulk dates is~~
 343 ~~justified since the absence of a systematic offset between the ^{14}C dates of bulk peat~~

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

~~versus those of plant macrofossils it contains~~ (G.M. MacDonald, pers. comm. 2013), the
use of bulk dates is justifiable. For the purpose of this study, all ¹⁴C ~~ages-dates~~ were
calibrated to calendar years before present (cal. BP) using the program CALIB 6.1.0
(Stuiver and Reimer, 1993) with the IntCal09 calibration dataset (Reimer et al., 2009). In
this paper, ages are reported in thousands of calibrated years before present (ka), ~~such~~
~~that 1 ka = 1000 cal. B.P.~~
Age-depth relationships were established for all continuous peat cores for which at least
five ~~radiocarbon dates~~ (¹⁴C) age determinations were available (n = 151 cores). Except for
a few palynostratigraphic and tephrochronologic markers, nearly all records have
chronologies exclusively based on ¹⁴C ages (Supplementary Table 1). Chronologies were
obtained through linear interpolation of calibrated ages between dated horizons. Single-
age estimates were taken from the mid-point of each calibrated 2σ probability
distribution. This parsimonious approach captures general patterns of temporal changes
in peat accumulation and allows for analysis of peat ¹⁴C accumulation trajectories (e.g.,
Telford et al., 2004). Although more sophisticated approaches are possible (e.g.,
Charman et al., 2013), we seek to make the fewest or simplest assumptions for this
analysis because the temporal resolution target for peat ¹⁴C accumulation rate calculations
is relatively low (at 500 years). In the cases where the original investigator identified
hiatuses (e.g., peat loss caused by erosion or fire) or depositional anomalies (e.g., thick
tephra layers that interrupted peat accumulation) along their peat records, these gaps were
taken into consideration when building age-depth relationships (Glaser et al., 2012).
Otherwise, peat records were assumed to be continuous.

367

368 In an effort to assess the representativeness of our samples in terms of peat^{land} inception

369 ~~age~~estimating, peat basal ages were compiled and compared to results from large datasets

370 (MacDonald et al., 2006; Gorham et al., 2007; Korhola et al., 2010). ~~When multiple cores~~

371 ~~were collected from a single peatland, only the oldest core was accounted for when~~

372 ~~estimating peat inception age~~. Peat inception ages from 19946 sites (Supplementary

373 Table 1) were summed and binned in 500-year intervals.

374

375 Long-term rates of carbon and nitrogen accumulation

376 A ~~subset-total~~ of 151 peat cores from 127 sites ~~were~~was used for estimating rates of peat -

377 C accumulation. This dataset contains 96 cores from 78 North American sites and 55

378 cores from 49 Eurasian sites (Figure 1). Of the 33 sites presented in Yu et al.'s (2009)

379 study, 25 were used in the present study (Supplementary Table 1). The remaining 8 sites

380 did not fulfill our dating quality criterion (presented below). When multiple cores were

381 collected from a single peatland, each core was attributed a fraction of the weight for that

382 site such that, for a site with three cores, the peat C accumulation history of each core

383 only accounted for 1/3. For each core, peat chronology was constrained by at least five

384 age determinations. The dating ~~resolution-quality~~ of ~~these each~~ records was determined

385 by the quotient of the calibrated peat basal age and the number of age determinations. For

386 example, a 10,000-year-old peat core with a chronology constrained by 10 ¹⁴C dates was

387 attributed a dating ~~resolution-quality~~ of ~~1000 years per one~~ date per 1000 years. About

388 58% of ~~these our~~ 151 cores ~~we~~are characterized by an acceptable dating ~~resolution~~

389 ~~between 10 and~~quality of one to two ¹⁴C dates ~~per~~ 1000 years (Figure 2b). These

Formatted: Superscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

resolutions are well suited to capture millennial-scale variations in C accumulation.

Several peat cores with ~~high resolution (10-500 years)~~ more than two ¹⁴C dates per 1000 years-per date, (n = 35 cores) ~~we~~are available from North America and Europe. The ~~coarser lower dating resolution quality~~ cores (~~<1 date per~~ >1000 years-per date) ~~we~~are unevenly distributed and comprise 44% of the North American records (~~42 out of 96~~ ~~cores~~) and 40% of the Eurasian records (~~22 out of 55 cores~~).

The long-term rate of peat-C accumulation was calculated for each core following one of the ~~these following~~ five approaches (Supplementary Table 1): (1) ~~w~~Whenever possible, peat core chronology was combined with bulk density and C% ~~content values~~ for each depth increment (n = 47 cores); (2) ~~i~~In the cases where direct C measurements were lacking, peat core chronology was combined with ~~ash-free~~organic matter bulk density measurements and a mean organic ~~matter carbon-C content value~~ of 49% ~~in organic matter~~ (n = 57 cores; ~~see the Results section for details~~); (3) ~~f~~For cores that lacked ~~ash-free~~organic matter bulk density and direct C% ~~content measurements~~, peat core chronology was combined with bulk density measurements and a ~~mean total carbon-C~~ content of 47% ~~in total peat~~ (n = 3 cores; ~~see the Results section for details~~); (4) ~~w~~Whenever neither bulk density nor C% ~~content~~ was ~~directly~~ available ~~from the cores~~, long-term rate of peat-C accumulation was calculated for each core by combining time-dependent bulk densities (0.08 g cm⁻³ at 0-0.5 ka; 0.12 g cm⁻³ at 0.5-6 ka; 0.14 g cm⁻³ at 6-12 ka) with a mean C content of 47% ~~in total peat~~ for each dated interval (n = 32 cores; ~~see the Results section for details~~), and; (5) ~~p~~Peat-C accumulation rates for the remaining 12 cores were directly obtained from published figures and tables. For all the cores, time-

Formatted: Superscript

weighted peat $\delta^{13}\text{C}$ accumulation rates were summed and binned in 500-year intervals. It is important to note that such reconstructions are 'apparent rates' that are different from true rates of C ~~uptake-accumulation~~ in peatlands, because decomposition processes have been affecting old peat layers for thousands of years (Turunen et al., 2002). Finally, the long-term rate of peat N accumulation was calculated by combining our binned peat-C accumulation rates with time-dependent C/N values (65 at 0-6 ka, and 40 at 6-10 ka).

Results

Peat Properties

~~, the mean bulk density value for each peat type is reported in Table 1, and regional differences are presented in Table 2.~~ Descriptive statistics of dataset
The frequency distribution of each peat property is shown in Figure 3. Mean values and standard deviations for all peat properties are presented by peat type in The frequency distribution is presented in Figure 3a Table 1, and by region in Table 2.

Bulk density values ($n = 21,220$) range~~ed~~ from 0.003 to 0.498 g cm^{-3} , with a mean value of $0.118 \pm 0.069 \text{ g cm}^{-3}$ (1 standard deviation (S.D.)). A one-way analysis of variance (ANOVA) reveal~~ed~~ an significant effect of peat type on bulk density ($F(10709) = 941$, $p < 0.0001$), with all peat types significantly different from each other on the basis of post-hoc Tukey's LSD tests ($p < 0.0001$). In increasing order, mean bulk density of the peat types is *Sphagnum* < Woody < Herbaceous < Brown Moss < Humified (Table 1).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Organic matter content (n = 18,973) has a mean value of $90.7 \pm 13\%$ (1 S.D.) and a median of 95.7%. The frequency distribution is presented in Figure 3b, the mean organic matter content for each peat type is reported in Table 1, and regional differences are presented in Table 2. The ANOVA revealed an significant effect of peat type on organic matter content ($F(9512) = 349, p < 0.0001$), with all peat types significantly different from each other (Tukey's LSD: $p < 0.0001$). In decreasing order, mean organic matter content of the peat types is *Sphagnum* > Woody > Herbaceous > Brown Moss > Humified (Table 1).

Organic matter bulk density or ash-free bulk density (n = 18,544) ranges from 0.003 to $0.452 \text{ g OM cm}^{-3}$, with a mean value of $0.105 \pm 0.051 \text{ g OM cm}^{-3}$ (1 S.D.). The frequency distribution is presented in Figure 3c, the mean organic matter density for each peat type is reported in Table 1, and regional differences are presented in Table 2. The ANOVA revealed an significant effect of peat type on organic matter density ($F(9081) = 942, p < 0.0001$), with all peat types significantly different from each other (Tukey's LSD: $p < 0.0001$). In increasing order, mean organic matter density of the peat types is *Sphagnum* < Herbaceous < Woody < Brown Moss < Humified (Table 1).

Total peat carbon C content in total peat (n = 3741) ranges from 30 to 60%, with a mean value of $46.8 \pm 6.1\%$ (1 S.D.) and a median of 47.8%. While the lowest values (< 35%) are almost exclusively associated with samples from Alaska, western Canada, Fennoscandia, and eastern Russia, the highest values (> 55%) are characteristic of sites located in the western European Islands and Fennoscandia. Mean C% values for all five

peat types are reported in Table 1, and regional differences are presented in Table 2. The ANOVA revealed an insignificant effect of peat type on C% ($F(2494) = 161, p < 0.0001$), with *Sphagnum* samples significantly different from other peat types (Tukey's LSD: $p < 0.0001$). Herbaceous and woody peats were distinct from other types, but hardly indistinguishable from one another (Tukey's LSD: $p = 0.238$). Likewise, humified and brown moss peats were distinct from other types, but indistinguishable from each other (Tukey's LSD: $p = 0.448$). In increasing order, mean C% of the peat types is *Sphagnum* < Humified = Brown Moss < Herbaceous = Woody (Table 1). The frequency distribution of C% in peat is also characterized by a second, though minor, mode at 40% (Figure 3d). The latter is mostly associated with *Sphagnum* peat samples, as their average C-content (C%) is significantly lower than those for other peat types (Table 1). This difference is likely caused by the high content of complex and recalcitrant compounds found in *Sphagnum* tissues such as lipids and waxes, which have lower C% than more labile biopolymers such as cellulose (Cagnon et al., 2009).

On the basis of 995 samples for which both OM% and C% were quantified, we therefore developed conversion factors (slopes of linear regressions) for several peat types, providing ways to estimate OC% for investigators who do not perform C% measurements routinely (Figure 4). There was a noticeable difference between the slope of *Sphagnum* peat (0.423 ± 0.030 ; $n = 454$) and that of non-*Sphagnum* peat (0.514 ± 0.024 ; $n = 308$). The carbon content in organic matter (OC%) for *Sphagnum* peat was smaller than expected (at $42.3 \pm 3.0\%$ (e.g., Bauer et al., 2006; Beilman et al., 2009; Table 3)). As the majority of our *Sphagnum* samples for this specific analysis were

1
2
3
4
5
6
7
8
9 481 younger than 0.5 ka (304 out of 454 samples) and extracted from raised bogs, it is very
10 482 likely that our estimated OC% biases towards young and undecomposed *Sphagnum*. This
11
12 483 ‘young *Sphagnum* peat effect’ heavily influenced the slope of the overall relation
13
14 484 between C% and OM% (0.467 ± 0.045) due to the overrepresentation of *Sphagnum*
15
16 485 samples in the dataset (454 out of 995 samples). To minimize this bias when estimating
17
18 486 mean OC% in peat, all 304 young *Sphagnum* samples were removed from the dataset,
19
20 487 yielding an overall conversion factor of 0.492 ± 0.024 .
21
22 488
23
24 489 ~~Total peat nitrogen-N~~ content in peat (n = 3365) ~~range~~ed from 0.04 to 3.39%, with a mean
25
26 490 value of $1.2 \pm 0.7\%$ (1 S.D.). ~~T~~he frequency distribution ~~is~~is asymmetric and
27
28 491 characterized by,~~with~~ a mode at 0.65% (Figure 3e). The latter is largely due to the
29
30 492 overrepresentation of *Sphagnum* peat samples (having low N content) in our
31
32 493 samplesdatabase. The ANOVA revealed ~~an~~ -significant effect of peat type on N content
33
34 494 ($F(2504) = 666, p < 0.0001$) with *Sphagnum* and herbaceous peat types significantly
35
36 495 different from each other and from all other types (Tukey’s LSD: $p < 0.0001$). Humified
37
38 496 and brown moss peats ~~awere~~ distinct from other types, but indistinguishable from each
39
40 497 other (Tukey’s LSD: $p = 0.113$). Likewise, woody and brown moss peats ~~awere~~
41
42 498 indistinguishable from one another (Tukey’s LSD: $p = 0.240$). In increasing order, mean
43
44 499 ~~N content~~N% of the peat types is *Sphagnum* < Woody = Brown Moss = Humified <
45
46 500 Herbaceous (Table 1).
47
48 501
49 502 C/N mass ratio (n = 3362) ~~range~~ed from 12 to 217, with a mean value of 55 ± 33 (1 S.D.).
50
51 503 The frequency distribution ~~is~~is asymmetric and characterized by,~~with~~ a mode at 25
52
53
54
55
56
57
58
59
60

(Figure 3f). While the distribution mode (25) ~~is~~ associated with non-*Sphagnum* peat types, the distribution mean (55) is skewed towards *Sphagnum* peat samples owing to their overrepresentation in our ~~samples database~~ (Figures 3f). The ANOVA revealed ~~an~~ ~~significant~~ effect of peat type on C/N ratio ($F(2501) = 174, p < 0.0001$) with *Sphagnum* peat significantly different from all other types (Tukey's LSD: $p < 0.0001$). Woody peat ~~was~~ distinguishable from all peat types except for brown moss peat (Tukey's LSD: $p = 0.665$). Herbaceous and humified peat types ~~are~~ indistinguishable from one another (Tukey's LSD: $p = 0.721$). In decreasing order, the mean C/N ratio of peat types is *Sphagnum* > Woody = Brown Moss > Humified = Herbaceous (Table 1).

Temporal changes in peat properties

The frequency distribution is presented in Figure 3a

We find a decreasing trend in bulk density over the Holocene (Figure 5a), with the densest peat characterizing the oldest samples (8-10 ka) and the least-dense peat characterizing the youngest samples (0-2 ka). This trend is most likely attributable to the progressive decomposition and subsequent compaction of peat over time, as well as to higher ash content in early-stage peat (likely from fens). Conversely, a clear increasing trend in organic matter content (OM%) is found over the Holocene (Figure 5b), with the greatest OM% characterizing the youngest samples (0-2 ka) and the least OM% characterizing the oldest samples (8-10 ka). This trend is most likely attributable to higher inorganic material inputs during early-stage peatland development as well as to a greater loss of OM in the deeper portions of peat profiles.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(OMBD) remains relatively constant over the Holocene (Figure 5c) because of the opposite trends exhibited by BD and OM% (Figures 5a, 5b). The only exceptions ~~except~~ ~~for~~ are the low OMBD values characterizing the youngest samples (< 0.5 ka), probably due to the large proportion of young, undecomposed *Sphagnum* peat samples.

While OMBD values closely tracked changes in bulk density during the mid and late Holocene (0–8 ka), their trajectory diverged between 8 and 10 ka (Figure 4a and 4e) because of decreasing organic matter content in early Holocene peat samples (Figure 4b). C content in peat remains uniform over the Holocene (Figure 5d), except for slightly lower C% during the late Holocene. However, as C values for early Holocene samples are based on a very limited number of samples and peat records (white bars in Fig. 4d), we caution against analysis or interpretation of the documented trend in our data. We also find a decreasing trend in N% ~~content~~ over the Holocene, such that young peat deposits are associated with low N% ~~content~~ (Figure 5e). This trend could be explained by a progressive retention of N downcore as a result of long-term peat decomposition processes and associated C loss (e.g., Kuhry and Vitt, 1996). Peat deposits older than 6 ka are mostly associated with lower C/N ratios, whereas peat samples younger than 6 ka are characterized by high C/N values (Figure 5f).

In general, *Sphagnum* peat is characterized by lower BD, OMBD, C%, N%, and C/N ratio than samples composed of non-*Sphagnum* peat (Figure 6). Therefore, peatland development could explain much of the aforementioned temporal trends (Figure 5), as

edense peat (herbaceous, brown moss, and humified types) typically characterarly-stage
 rich fens are typically characterized by non-*Sphagnum* peat, whereas and late-stage poor
 fens and bogs are *Sphagnum*-dominated (Figure 6h). ~~AlternativeThis effect may be~~
~~partly explained by peat type differences between young and old samples, with young~~
~~samples mostly composed of *Sphagnum* peat (having low N content) and old samples~~
~~composed of non-*Sphagnum* peat (having high N content). peat type differences between~~
~~young and old samples, with young samples mostly composed of *Sphagnum* peat (having~~
~~low N content) and old samples composed of non-*Sphagnum* peat (having high N~~
~~content). This 'peat type effect' may also be combined with~~

Spatial differences in peat properties
 Significant differences in bulk density are found at the regional scale (Table 2). The
 densest peat is observed in Alaska (mean = $0.168 \pm 0.087 \text{ g cm}^{-3}$) and western Canada
 (mean = $0.166 \pm 0.076 \text{ g cm}^{-3}$), whereas the least dense peat is found from the western
 European Islands (mean = $0.055 \pm 0.027 \text{ g cm}^{-3}$). Organic matter bulk density values
 follow a similar pattern across these regions (Table 2). These differences are strongly
 correlated with peat types and sample ages, with the Alaskan and western Canadian
 samples largely constituted of herbaceous, humified, and brown moss peat types. These
 regional differences were strongly correlated with peat types and sample ages, with the
 United Kingdom samples mostly composed of young *Sphagnum* peat and the Alaskan
 samples largely constituted of old herbaceous, humified, and brown moss peat types.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

573
574 Significant differences in bulk density were found at the regional scale (Table 2 and
575 Figure 6a). The densest peat was found at permafrost sites (), whereas the least dense peat
576 was reported from non-permafrost peatlands (). There was also a marked difference
577 between Alaskan (mean = $0.168 \pm 0.087 \text{ g cm}^{-3}$) and western Canadian sites (mean =
578 $0.166 \pm 0.076 \text{ g cm}^{-3}$) versus the Western European Island sites (mean = $0.055 \pm 0.027 \text{ g$
579 cm^{-3}). These regional differences were strongly correlated with peat types and sample
580 ages, with the United Kingdom samples mostly composed of young *Sphagnum* peat and
581 the Alaskan samples largely constituted of old herbaceous, humified, and brown moss
582 peat types.
583
584 OM% does not vary much between regions (> 90% in all regions), with the notable
585 exception of Alaskan and eastern Russian/Asian peatlands that exhibit a mean values of
586 $76.6 \pm 18.8\%$ and $80.3 \pm 16.7\%$, respectively (Table 2). Aeolian dust and tephra ash
587 inputs to some peatlands in Alaska, Kamchatka, and Japan might partly explain such low
588 OM% values.
589
590 ***Peat inception ages and long-term rates of carbon and nitrogen accumulation***
591 Calibrated ages (mid-point) for peat inception ranged from 04.6 to 15 ka and the
592 frequency distribution followed a bimodal distribution with this characterized by a modes at
593 11-9 ka and at 8-6 ka (Figure 2c). The early Holocene model latter corresponds with peat
594 inception peaks in the West Siberian Lowlands and, in Alaska, and Fennoscandia. The
595 mid Holocene peak is linked with peatland inception across western Canada and the

Hudson Bay Lowlands. While the majority of sites older than 11 ka are found in Alaska and Siberia, most sites younger than 6 ka are located in northeastern Canada. Generally speaking, in general, our samples are in agreement with much larger networks of peat basal ages (Smith et al., 2004; MacDonald et al., 2006; Gorham et al., 2007; Korhola et al., 2010; Ruppel et al., 2013; Yu et al., 2013), except for our underrepresentation of peatlands with inception ages between 10 and 8 ka (Figure 2b). This discrepancy may be attributable to an underrepresentation of Siberian sites in our dataset (Smith et al., 2004).

The time-weighted long-term rate of C accumulation averages 22.9 ± 2.0 g C m⁻² yr⁻¹ (standard error of mean (S.E.M.); Figure 7b). Values exhibit an increasing trend that initially peaks during the early Holocene between 10 and 7.5 ka at 25.2 ± 2.6 g C m⁻² yr⁻¹. This peak is largely caused by rapid peat accumulation in Alaska, the Western Siberia Lowlands, and southeastern Canada. The remainder of the Holocene is characterized by a decreasing trend in C accumulation rates from 24 to 18 g C m⁻² yr⁻¹ and a time-weighted mean at 22.0 ± 1.9 g C m⁻² yr⁻¹, and millennial scale variations (Figure 7b). There is a notable minimum value between 3 and 1.5 ka at 18-19 g C m⁻² yr⁻¹. Lack of decomposition probably explains most of the apparent increase in accumulation over the past millennium (24-32 g C m⁻² yr⁻¹), as young peat appears to be accumulating more quickly than old peat simply because the former has undergone less decomposition than the latter (Clymo, 1984).

The time-weighted long-term rate of N accumulation averages 0.5 ± 0.04 g N m⁻² yr⁻¹ (S.E.M.; Figure 7c). While the mid and late Holocene (6-0 ka) are characterized by the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

lowest rates of N accumulation at 0.34 g N m⁻² yr⁻¹, the highest rates (0.61 g N m⁻² yr⁻¹) occur between 12 and 6 ka (Figure 7c). This trend mirrors that of C accumulation (Figure 7b), as C and N sequestration rates are both mainly influenced by peat density and its accumulation rate. The low rates of N accumulation over the past 6 ka might also relate to the increasing presence and persistence of *Sphagnum* (having high C/N ratio and low N concentration) in northern peatlands (Figures 6 and 7).

Discussion

~~R~~Data representativeness of the database for northern peatlands

The present database contains the most comprehensive compilation of peat properties and C accumulation records for northern peatlands. The previous large-scale synthesis (Yu et al., 2009) only contained 33 sites and lacked records from the Hudson Bay lowlands and the Russian Far East, and had limited sites from West Siberia and the western European Islands. The present database fills gaps from these regions.

However, European Russia. The database comprises 268 peat cores from 215 sites located between 45 and 69 °N, throughout the circum-Arctic peatland domain. These sites were found across broad gradients of continentality, temperature, growing season length, and precipitation (Yu et al., 2009). The majority of our study sites were found in large peatland complexes such as the Hudson Bay Lowlands or in peatland-rich regions such as the northwestern European Islands and Fennoscandia. Conversely, smaller peatland systems such as kettle bogs and other isolated features were sparsely represented. As the main objective of our dataset was to estimate the northern peatland C

Formatted: Comment Text

stock, the high proportion of sites from large peatland complexes seems justified. On another note, further studies of small depressional peatlands could potentially lead to a better understanding of peat C accumulation processes and peatland sensitivity to climate change (e.g., Buffam et al., 2010; Ireland et al., 2013). East Siberia, and the Russian Far East clearly remain poorly studied regions in terms of northern peat C stocks and accumulation histories (Figure 1). A wetland map by Stolbovoi and McCallum (2002) suggests that shallow peaty deposits (<50 cm) interspersed with few deeper peat bogs (>50 cm) dominate the Far East Russian landscape. Most of these deeper peatlands are presumably found in Kamchatka and Sakhalin (Stolbovoi, 2002). This broad portrait is, however, based on fewer than 30 soil profiles from across East Siberia and Far East Russia (Stolbovoi et al., 2001), making it difficult to evaluate the importance of this region in the northern peatland C cycle. In general, peat C stocks in Eastern Russia may not be as massive as those from West Siberia or European Russia (Stolbovoi and McCallum, 2002). Therefore, understanding how these shallow peatlands in East Siberia and the Russian Far East have developed during the Holocene would provide useful end-members of climate controls of peat C accumulation, but these peatlands do not seem to represent a large missing C stock.

659

660 *Northern peatland soil properties: key findings and uncertainties*

661 Peat-carbon stocks and density

662 Several studies have quantified the soil C density and total C pool of peatlands using
663 different approaches (e.g., Armentano and Menges, 1986; Gorham, 1991; Yu et al.,
664 2010). These different methods have led to total C pool estimates for northern peatlands

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

that vary by at least a factor of two, from 234 to 547 GtC (Lappalainen, 1996; Yu et al., 2010; see Yu, 2012 for a review). Many of these studies have combined mean peat depth, modern peatland area, and a single mean ~~peat-C bulk~~-density value (BD x C% or OMBD x OC%) in their calculations (e.g., Gorham, 1991). Applying Gorham's (1991) mean peat depth and peatland area estimates to the mean BD and C% results ~~presented in this~~from our database yields a ~~peat-C~~ pool estimate of 436 GtC (2.3 m x 3.42 Mkm² x 0.118 g cm⁻³ x 47% C). However, it is well documented that most peatlands undergo a shift from herbaceous ~~fen~~-to *Sphagnum* ~~peat~~bog during their developmental history (Hughes, 2000; Figure 6h) and that different BD, C%, and rates of peat accumulation are associated with fen and bog peats (e.g., Vitt et al., 2000; Figure 6). We also know that peat-C accumulation rates have varied asynchronously between regions throughout the Holocene as a result of regional changes in hydroclimatic conditions (e.g., Yu et al., 2009; Charman et al., 2013). Therefore, we argue that reconstructing Holocene changes in peat-C accumulation on the basis of *measured* peat C density ~~peat-C bulk density values~~ and reliable peat-core chronologies constitutes a step forward in providing the best possible peat-C stock estimates (see Yu et al., 2010~~99~~ for an example). It also allows for quantifying spatial and temporal differences in rates of peat-C accumulation, as well as the temporal trajectories of peat-C fluxes to the atmosphere (MacDonald et al., 2006; Yu et al., 2013). However, better maps of the present peatland area (and its change over time) are still needed to improve current peat C stock estimates. ~~Therefore, our database should be of great use for updating current peat C stock estimates, as it contains the most extensive set of peat-C bulk density measurements.~~

Carbon content in organic matter

For each peat layer, peat ~~C~~_{carbon bulk} density can be estimated by the product of either (1) bulk density and ~~total peat C~~_{carbon} content in total peat (BD x C%), or (2) ash-free bulk density and ~~carbon C~~_{content in organic matter} (OMBD x OC%). ~~It is important to mention that peat C stocks estimated using BD x C% or OMBD x OC% yield the same value. Whenever BD and C% measurements were available along peat cores, we used the former formula. However, when only OMBD was available, the second formula was used.~~ It could be argued that the first option is preferable when estimating peat ~~C~~_{stocks}, as it produces values that are directly comparable to routine soil ~~C~~_{measurements} from other terrestrial ecosystems. However, the present database clearly indicates that the majority of peatland scientists routinely analyze organic matter content (OM%; n = 18,973 samples) rather than C% (n = 3741 samples) along peat cores. To provide a way to estimate OC% from OM%, we developed the following conversion factors: $42.3 \pm 3.0\%$ for *Sphagnum* peat, $51.4 \pm 2.4\%$ for non-*Sphagnum* peat, and $49.2 \pm 2.4\%$ overall (Figure 4, Table 1).

While the overall peat and the non-*Sphagnum* peat conversion factors are in line with those from previous studies (e.g., Gorham, 1991; Vitt et al., 2000; Turunen et al., 2002; Beilman et al., 2009), the *Sphagnum* peat factor is lower than other estimates (e.g., Bauer et al., 2006; Beilman et al., 2009; Table 3). Indeed, our mean OC% *Sphagnum* value at 42.3% is close to that of surface *Sphagnum* tissues, suggesting that it constitutes a valid estimate for ash-free and poorly decomposed *Sphagnum* peat. As previously mentioned,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

this bias towards low OC% is due to a large number of *Sphagnum* samples younger than 0.5 ka (304 out of 454 *Sphagnum* samples).

Although each one of the three conversion factor slopes was significant ($p < 0.0001$), there is a noticeable scatter in the data (Figure 4) that cannot solely be explained by the ~1% analytical error associated with the loss-on-ignition procedure (Heiri et al., 2001). The progressive accumulation of recalcitrant C in old samples (lignin ~ 60% C vs. cellulose ~ 42% C), assuming it occurs at a greater rate than the loss of OM in the deeper portions of peat profiles, could explain why C% appears higher than our OC% conversion factors (Cagnon et al., 2009). The presence of inorganic C, particularly for the humified and brown moss peat types, could also explain these results. While the overall peat and the non-*Sphagnum* peat conversion factors are in line with those from previous studies (e.g., Gorham, 1991; Vitt et al., 2000; Turunen et al., 2002; Beilman et al., 2009), the *Sphagnum* peat factor is lower than other estimates. Indeed, our mean OC% *Sphagnum* value at 42% is close to that of surface *Sphagnum* tissues, suggesting that it constitutes a valid estimate for ash-free and poorly decomposed *Sphagnum* peat. As previously mentioned, this bias toward low OC% is due to a large number of *Sphagnum* samples younger than 0.5 ka (304 samples out of 454).

Oligotrophication and the fen-to-bog transition in northern peatlandsCarbon content in organic matter

Formatted: Font: Font color: Black

The relation between total peat carbon content (C%) and peat organic matter content (OM%) is presented in Figure 4 (n = 995).

Our averaged bulk density ($0.12 \pm 0.07 \text{ g cm}^{-3}$) and organic matter bulk density ($0.11 \pm 0.05 \text{ g OM cm}^{-3}$) values are within the range of most widely used estimates (e.g., Clymo, 1984; Yu et al., 2010; Table 3).

Sphagnum and non-*Sphagnum* peat types were characterized by very different peat properties, with *Sphagnum* peat having lower BD, OMBD, C%, N%, and C/N ratio than non-*Sphagnum* peat (Figure 6). These differences become important when estimating Holocene peat C fluxes, as the proportion of *Sphagnum*-dominated peat records increases during the late Holocene due to the fen-to-bog transition (Figure 6h). For example, much stronger CH₄ emissions are associated with fens than bogs (e.g., Pelletier et al., 2007). In terms of C sequestration rates, the systematically higher organic C density of non-*Sphagnum* peat suggests that higher accumulation rates are possible in fens than in bogs (Figure 6g), assuming optimal hydroclimatic conditions leading to rapid peat burial. In addition, as non-*Sphagnum* samples contain twice the N mass of *Sphagnum* peat (Figure 6e), early-stage fens have the ability to stock more N than late-stage bogs. Overall, further studies on the timing of the fen-to-bog transition across the northern peatland domain are needed to better our understanding of its impact on C sequestration and CH₄ emissions.

Formatted: Subscript

Formatted: Subscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

*Holocene pattern of ~~circum-Arctic peatland~~ carbon accumulation in northern
peatlands*

The overall trajectory and shape of our Holocene peat -C accumulation curve is similar to the synthesis ~~from~~ a much smaller -dataset ($n = 33$; Yu et al., 2009). As such, an that
shows an early Holocene peak during the Holocene Thermal Maximum (HTM) as well
as and an overall slowdown of carbon -C accumulation during the mid- and late-Holocene, particularly after 4 ka during the Neoglacial period and associated permafrost development, were found in both syntheses (Vitt et al., 2000; Figure 76). However, the mean Holocene value of $22.9 \pm 2.0 \text{ g C m}^{-2} \text{ yr}^{-1}$ (1 S.E.) presented here is approximately 24% higher than the estimate in Yu et al.'s 2009 study ($18.6 \text{ g C m}^{-2} \text{ yr}^{-1}$). Our larger dataset likely better represents the northern peatland -C accumulation rates~~stock~~. These results imply that current peat -C stocks might be underestimated.

While the peak value at $27 \text{ g C m}^{-2} \text{ yr}^{-1}$ is about 23% higher than the time-weighted mean
peat-C accumulation rate for the remainder of the Holocene at $22 \text{ g C m}^{-2} \text{ yr}^{-1}$, we only
found a 2% difference in organic C density values between young ($0.053 \pm 0.02 \text{ g C cm}^{-3}$
 3) and old ($0.057 \pm 0.03 \text{ g C cm}^{-3}$) peat samples. These results clearly show that the peak
value during the early Holocene cannot be mainly attributed to presumably dense peat
deposits that would be rich in recalcitrant C due to long-term decomposition and
compaction. Instead, factors influencing the rate of peat burial such as peat type
(*Sphagnum* vs. non-*Sphagnum* peat; Figure 6), growing season length, and other
environmental variables, must have been responsible for such high rates of C
sequestration during the early Holocene.

778

779 The Holocene Thermal Maximum (HTM) is a well-documented period of orbitally-

780 induced warm climate in the northern high-latitude region (Kaufman et al., 2004;

781 Renssen et al., 2012; Marcott et al., 2013) that reaches its maximum around 11 ka

782 (Berger and Loutre, 1991; Figure 7a). The peak in warm climatic conditions shows a

783 transgressional pattern across northern North America that moved eastward with the

784 waning Laurentide Ice Sheet during the early and mid Holocene (Kaufman et al., 2004).

785 This progressive increase in land availability coupled with warming summer conditions

786 have been proposed as the main controls on peatland inception and rapid C accumulation

787 across northern North America (Harden et al., 1992; Gorham et al., 2007, 2012; Yu et al.,

788 2009; Jones and Yu, 2010). As such, the highest rates of C accumulation have been

789 recorded between 11 and 8.5 ka in Alaska, but only around 7 ka in western Canada

790 (Figure 6; Vitt et al., 2000; Yu et al., 2009; Jones and Yu, 2010). In general, our results

791 confirm this trend and support the hypothesis that warm summers could promote peat

792 formation and C sequestration (Beilman et al., 2009; Yu et al., 2009; Charman et al.,

793 2013), as the highest rates of C accumulation broadly coincide with the peak in summer

794 insolation from 11 to 7 ka (Figure 7). We acknowledge that water input was necessary to

795 allow for peatland development. Furthermore, the observed temporal asymmetry in

796 peatland inception age and peaks in C accumulation rates between Alaska, western

797 Canada, and the Hudson Bay Lowlands follows the transgressional pattern of the HTM.

798 For example, peat inception and highest peat C accumulation rates occur at 11-9 ka in

799 Alaska, whereas they are delayed in western Canada with peak values around 9-7 ka.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

These findings have important implications for projecting the fate of peat-C stocks in a future warmer world.

The Neoglacial period is characterized by generally cooler and wetter conditions than the HTM (Figure 7a; Marcott et al., 2013). Particularly low C accumulation rates coincide with this time period across the northern peatland domain (Figure 7; Vitt et al., 2000; Jones and Yu, 2010). Peat accumulation processes might even have stopped in some regions (e.g., Peteet et al., 1998). The onset of permafrost aggradation in many peatlands also occurred during the Neoglacial period (Zoltai, 1971, 1995; Vitt et al., 2000; Oksanen et al., 2003; Sannel and Kuhry, 2008), reducing the peat C-sink capacity. In addition to shorter and cooler growing seasons, lower C accumulation rates in permafrost sites likely relates to a slower peat burial due to (1) more intense peat decomposition in the acrotelm due to drier surface conditions, and (2) a slower rate of peat formation and associated C inputs to soil because many peat plateaus are not *Sphagnum*-dominated. Overall, our results support the notion that climatic changes such as the HTM and the Neoglacial cooling impact C sequestration rates in peatlands.

Role of northern peatlands in the global nitrogen cycle

As relatively few downcore peat N concentrations have been reported in the literature, it was difficult to compare our mean value of 1.2% to previous estimates. ~~Percent N in new foliage of vascular species growing in subarctic peatlands average 1.8% (Schuur et al., 2007), similar to our mean value for non-*Sphagnum* peat types (1.5%).~~ Bragazza et al. (2012) reported N content values of 0.7% for *Sphagnum fuscum* litter and 1.48% for

823 *Eriophorum vaginatum* (herbaceous) litter, ~~also~~ in line with our results (Table 1).
 824 Similarly, Turunen et al. (2004) documented peat N concentrations ranging from 0.35 to
 825 2.25% (mean value of 0.8%) for the uppermost sections of 23 *Sphagnum* bogs across
 826 northeastern Canada. Overall, these values closely match our findings for *Sphagnum*
 827 (0.7%) and herbaceous (1.7%) peat types (Table 1).
 828
 829 Using our peat C pool estimate of 436 Gt and assuming a mean C/N ratio of 45 yields a
 830 peat-N pool of 9.7 Gt, roughly equivalent to 10% of the global soil N pool at 95 Gt (Post
 831 et al., 1985). This estimate is within the range proposed by Limpens et al. (2006) at 8-15
 832 GtN. We also calculated 500-year bin N accumulation rates for the past 10,000 years by
 833 combining our binned peat C accumulation rates with time-dependent C/N values.
 834 ~~Results indicate a~~ The Holocene time-weighted peat-N accumulation rate ~~of~~ 0.5 ± 0.04
 835 $\text{g N m}^{-2} \text{ yr}^{-1}$ (S.E.; Figure 76) is also in line with a previous estimate of 0.19-0.48 g N m⁻²
 836 yr⁻¹ (Limpens et al., 2006). While the mid and late Holocene (6-00-6 ka) ~~are~~
 837 characterized by the lowest rates of peat-N accumulation at $0.34 \text{ g N m}^{-2} \text{ yr}^{-1}$, the highest
 838 rates ($0.61 \text{ g N m}^{-2} \text{ yr}^{-1}$) occurred between 126 and 642 ka (Figure 7c6). ~~This trend~~
 839 ~~mirrors that of C accumulation (Figure 6), as C and N sequestration rates are both~~
 840 ~~influenced by bulk density and peat accumulation rates.~~ The low rates of N accumulation
 841 over the past 6 ka might also relate to the increasing presence and persistence of
 842 *Sphagnum* peat (having high C/N ratio and low N concentration) across the northern
 843 peatlands ~~as a result of the fen to bog transition~~ (Figures 6 and 76). Overall, given the
 844 bias toward *Sphagnum*-dominated sites in our database, N pools and N accumulation
 845 rates are probably underestimated.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

846

847 Rapid N sequestration in peatlands during the early Holocene might have contributed to

848 the global decline in reactive N availability for terrestrial ecosystems (McLaughlan et

849 al., 2013), pointing to a potentially important and undocumented role of northern

850 peatlands in the global N cycle. These results also raise the important question of N

851 provenance: in the absence of large rates of atmospheric N deposition during the early

852 Holocene, the only process that could account for such a large N pool in peatlands is N

853 fixation, either through symbiotic or asymbiotic processes (Limpens et al., 2006).

854

855 The fate of these large peat-N stocks remains largely unknown under recent and

856 projected warming. Indeed, the importance of peatlands as sources of nitrous oxide (N₂O)

857 is just emerging (e.g., Repo et al., 2009; Marushchak et al., 2011; Palmer et al., 2012),

858 and studies have suggested that reduced surface moisture or increasing temperatures

859 might significantly promote the production, transformation, and transport of dissolved N,

860 and N₂O emissions to the atmosphere through denitrification (e.g., Kane et al., 2010). On

861 the contrary, some authors have speculated that the potential increase in peatland-N₂O

862 emissions from climate change may not be significant relative to the global N₂O budget

863 (e.g., Martikainen et al., 1993; Froking et al., 2011). Overall, additional peat N cycling

864 studies are needed to address these remaining questions.

866 **Conclusions and future directions**

867 Peat core analysis has been extensively used over the past 20 years for estimating rates of

868 peat-Carbon accumulation at local, regional, and continental scales (e.g., Mäkilä, 1997;

869 Clymo et al., 1998; Vitt et al., 2000; Turunen et al., 2002; [Mäkälä and Saarnisto, 2008](#);
 870 Yu et al., 2010; van Bellen et al., 2011; Gorham et al., 2012). The present study analyzed
 871 a new database that comprises 268 peat records from 215 [northern peatland](#) sites ~~located~~
 872 ~~throughout the circum-Arctic peatlands~~. This systematic analysis of peat properties and
 873 Holocene C accumulation rates is essential for accurately addressing the following
 874 general research topics in the future: (1) describing and quantifying spatial and temporal
 875 patterns of Holocene peatland C and N accumulation; (2) assessing the sensitivity of C
 876 and N accumulation to climate ~~forcing~~[change](#); (3) estimating peatland ~~soil organic matter~~
 877 ~~(SOM)~~, soil organic carbon (SOC), and soil organic nitrogen (SON) pools at regional and
 878 hemispheric scales, (4) furthering our understanding of peatland ~~C cycle-carbon~~ --
 879 climate linkages, and (5) providing the scientific community with a large dataset for
 880 developing and testing earth system and ecological models.

882 Acknowledgements

883 ~~The authors would like to~~[We](#) acknowledge the peatland research community for sharing
 884 their datasets. The U.S. NSF supported the synthesis work through grant ARC-1107981
 885 [to Lehigh University](#). The collection and analysis of unpublished records used in this
 886 synthesis were supported by the following funding agencies and research grants: Alaska
 887 (NSF ARC-1107981, AGS-0628455, and EAR-0819717; [USGS Climate Research and](#)
 888 [Development Program](#)), Canada (NSF ARC-1107981, EAR-0223271, EAR-0843685,
 889 and AGS-0628598; NSERC CRDPJ-305605, CRDPJ-365867; Hydro-Québec),
 890 Fennoscandia and Western Siberia ([NSF OPP-9818496](#); Academy of Finland 201321 and
 891 1133515; University of Helsinki), [Kamchatka \(NSF ARC-1107981, ARC-1108116\)](#), and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

the United Kingdom (Yorkshire Peat Partnership). Lehigh University’s Library and Technology Services staff is acknowledged for its support in building the web interface for the peatland database. [Finally, comments from Paul Glaser and two other journal reviewers improved the overall quality of the manuscript.](#)

References

Armentano TV and Menges ES (1986) Patterns of change in the carbon balance of organic soil-wetlands of the temperate zone. *Journal of Ecology* 74: 755-774.

Bauer IE, Bhatti JS, Cash KJ, Tarnocai, C and Robinson SD (2006) Developing statistical models to estimate the carbon density of organic soils. *Canadian Journal of Soil Science* 86: 295-304.

Beilman DW, MacDonald GM, Smith LC and Reimer PJ (2009) Carbon accumulation in peatlands of West Siberia over the last 2000 years. *Global Biogeochemical cycles* 23, GB1012. doi: 10.1029/2007GB003112.

Belyea LR (2009) Non-linear dynamics of peatlands and potential feedbacks on the climate system. In: Baird A, Belyea L, Comas X, Reeve A. and Slater L (Eds.), *Northern peatlands and carbon cycling*, American Geophysical Union Monograph Series, Washington D.C., USA, pp. 5-18.

Berger A and Loutre M-F (1991) Insolation values for the climate of the last 10 million years. *Quaternary Science Reviews* 10: 297-317.

[Bergner K, Albano Å and Bohlin E \(1990\) The content of peat: a compilation of botanical, physical and chemical data of peat. Department of Agricultural Research, Northern Sweden, Swedish University of Agricultural Sciences.](#)

Björck S and Clemmensen LB (2004) Aeolian sediment in raised bog deposits, Halland, SW Sweden: a new proxy record of Holocene winter storminess in southern Scandinavia? *The Holocene* 14: 677-688.

Blodau C (2002) Carbon cycling in peatlands - A review of processes and controls, *Environmental Reviews* 10: 111-134.

Bragazza L, Buttler A, Habermacher J, Brancaleoni L, Gerdol R, Fritze H, Hanajik P, Laiho R and Johnson D (2012) High nitrogen deposition alters the decomposition of bog plant litter and reduces carbon accumulation. *Global Change Biology* 18: 1163-1172.

- 932 Bridgman SC, Megonigal JP, Keller JK, Bliss, NB and Trettin C (2006) The carbon
 933 balance of North American wetlands. *Wetlands* 26: 889-916.
 934
- 935 Buffam I, Carpenter SR, Yeck W, Hanson PC and Turner MG (2010) Filling holes in
 936 regional carbon budgets: Predicting peat depth in a north temperate lake district. *Journal*
 937 *of Geophysical Research – Biogeosciences* 115, G01005, doi:10.1029/ 2009JG001034.
 938
- 939 [Cagnon B, Py X, Guillot A, Stoeckli F and Chambat G \(2009\) Contributions of](#)
 940 [hemicellulose, cellulose and lignin to the mass and the porous properties of chars and](#)
 941 [steam activated carbons from various lignocellulosic precursors. *Bioresource Technology*](#)
 942 [100: 292-298.](#)
 943
- 944 Camill P (1999) Peat accumulation and succession following permafrost thaw in the
 945 boreal peatlands of Manitoba, Canada. *Ecoscience* 6: 592-602.
 946
- 947 Chambers FM, Beilman DW and Yu Z (2011) Methods for determining peat
 948 humification and for quantifying peat bulk density, organic matter and carbon content for
 949 palaeostudies of climate and peatland carbon dynamics. *Mires and Peat* 7, article 7, 10 p.
 950
- 951 Charman D, Beilman D, Blaauw M, Booth RK, Brewer S, Chambers F, Christen JA,
 952 Gallego-Sala AV, Harrison SP, Hughes PDM, Jackson S, Korhola A, Mauquoy D,
 953 Mitchell F, Prentice IC, van der Linden M, De Vleeschouwer F, Yu Z, Alm J, Bauer IE,
 954 McCorish Y, Garneau M, Hohl V, Huang Y, Karofeld E, Le Roux G, Loisel J, Moschen
 955 R, Nichols JE, Nieminen TM, MacDonald GM, Phadtare NR, Rausch N, Sillasoo Ü,
 956 Swindles GT, Tuittila E-S, Ukonmaanaho L, Välranta M, van Bellen S, van Geel B, Vitt
 957 D and Zhao Y (2013) Climate-related changes in peatland carbon accumulation during
 958 the last millennium. *Biogeosciences* 10: 929-944. doi: 10.5194/bg-10-929-2013.
 959
- 960 Clymo RS (1984) The limits to peat growth. *Philosophical Transactions of the Royal*
 961 *Society of London, Series B, Biological Sciences* 303: 605-654.
 962
- 963 Clymo RS, Turunen J and Tolonen K (1998) Carbon accumulation in peatlands. *Oikos*
 964 81: 368-388.
 965
- 966 Dean Jr WE (1974) Determination of carbonate and organic matter in calcareous
 967 sediments and sedimentary rocks by loss on ignition: comparison with other methods.
 968 *Journal of Sedimentary Petrology* 44: 242-248.
 969
- 970 Frolking S and Roulet NT (2007) Holocene radiative forcing impact of northern peatland
 971 carbon accumulation and methane emissions. *Global Change Biology* 13: 1-10.
 972
- 973 Frolking S, Talbot J, Jones MC, Treat CC, Kauffman JB, Tuittila E-S and Roulet N
 974 (2011) Peatlands in the Earth's 21st century coupled climate-carbon system.
 975 *Environmental Reviews* 19: 371-396.
 976

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

977 | [Glaser PH, Volin JC, Givnish TJ, Hansen BCS and Stricker CA \(2012\) Carbon and](#)
978 | [sediment accumulation in the Everglades \(USA\) during the past 4000 years: rates,](#)
979 | [drivers, and sources of error. *Journal of Geophysical Research-Biogeosciences* 117,](#)
980 | [G03026. doi:10.1029/2011JG001821.](#)
981 |
982 | Gorham E (1990) Biotic impoverishment in northern peatlands. In: Woodwell GM (Ed.),
983 | *The Earth in transition: Patterns and processes of biotic impoverishment*. Cambridge
984 | University Press, New York, USA, pp. 65-98.
985 |
986 | Gorham E (1991) Northern peatlands: role in the carbon cycle and probable responses to
987 | climatic warming. *Ecological Applications* 1: 182-195.
988 |
989 | Gorham E, Lehman C, Dyke A, Janssens J and Dyke L (2007) Temporal and spatial
990 | aspects of peatland initiation following deglaciation in North America. *Quaternary*
991 | *Science Reviews* 26: 300-311.
992 |
993 | Gorham E, Lehman C, Dyke A, Clymo D and Janssens J. (2012) Long-term carbon
994 | sequestration in North American peatlands. *Quaternary Science Reviews* 58: 77-82.
995 |
996 | Harden JW, Sundquist ET, Stallard RF and Mark RK (1992) Dynamics of soil carbon
997 | during deglaciation of the Laurentide ice sheet. *Science* 258: 1921-1924.
998 |
999 | [Heiri O, Lotter AF and Lemcke G \(2001\) Loss on ignition as a method for estimating](#)
1000 | [organic and carbonate content in sediments: reproducibility and comparability of results.](#)
1001 | [Journal of Paleolimnology 25: 101-110.](#)
1002 |
1003 | ~~Hughes~~ PDM (2000) A reappraisal of the mechanisms leading to ombrotrophy in
1004 | British raised mires. *Ecology Letters* 3: 7-9.
1005 |
1006 | Ireland AW, Booth RK, Hotchkiss SC and Schmitz JE (2013) A comparative study of
1007 | within-basin and regional peatland development: implications for peatland carbon
1008 | dynamics. *Quaternary Science Reviews* 61: 85-95.
1009 |
1010 | [Jones MC and Yu Z \(2010\) Rapid deglacial and early Holocene expansion of peatlands in](#)
1011 | [Alaska. *Proceedings of the National Academy of Sciences* 107: 7347-7352.](#)
1012 |
1013 | Kane ES, Turetsky MR, Harden JW, McGuire AD and Waddington JM (2010) Seasonal
1014 | ice and hydrologic controls on dissolved organic carbon and nitrogen concentrations in a
1015 | boreal-rich fen. *Journal of Geophysical Research – Biogeosciences* 115, G04012. doi:
1016 | 10.1029/2010JG001366.
1017 |
1018 | Kaufman DS, Ager TA, Anderson NJ, Anderson PM, Andrews JT, Bartlein PJ, Brubaker
1019 | LB, Coats LL, Cwynar LC, Duvall ML, Dyke AS, Edwards ME, Eisner WR, Gajewski
1020 | K, Geirsdottir A, Hu FS, Jennings AE, Kaplan MR, Kerwin MW, Lozhkin AV,
1021 | MacDonald GM, Miller GH, Mock CJ, Oswald WW, Otto-Bliesner BL, Porinchu DF,

- 1022 Ruhland KR, Smol JP, Steig EJ, Wolfey BB (2004) Holocene thermal maximum in the
 1023 western Arctic (0-180 °W). *Quaternary Science Reviews* 23: 529-560.
 1024
- 1025 Korhola A, Ruppel M, Seppä H, [Välranta M](#), Virtanen T and Weckström J (2010) The
 1026 importance of northern peatland expansion to the late-Holocene rise of atmospheric
 1027 methane. *Quaternary Science Reviews* 29: 611-617.
 1028
- 1029 Lappalainen E (1996) General review on world peatlands and peat resources. In:
 1030 Lappalainen E (Ed.), *Global Peat Resources*. International Peat Society, Jyska, pp. 53-56.
 1031
- 1032 Limpens J, Heijmans MPD and Berendse F (2006) The nitrogen cycle in boreal
 1033 peatlands. In: Wieder RK and Vitt DH (Eds.), *Boreal Peatland Ecosystems*. Ecological
 1034 Studies Vol. 188, Springer-Verlag, Berlin Heidelberg, Germany, pp. 195-230.
 1035
- 1036 MacDonald GM, Beilman DW, Kremenetski KV, Sheng Y, Smith LC and Valichko AA
 1037 (2006) Rapid early development of circum-arctic peatlands and atmospheric CH₄ and
 1038 CO₂ variations. *Science* 314: 285-288.
 1039
- 1040 Mäkilä M (1997) Holocene lateral expansion, peat growth and carbon accumulation on
 1041 Haukkasuo, a raised bog in southeastern Finland. *Boreas* 26: 1-14. doi:10.1111/j.1502-
 1042 3885.1997.tb00647.x.
 1043
- 1044 [Mäkilä M and Saarnisto M \(2008\) Carbon accumulation in boreal peatlands during the
 1045 Holocene – impacts of climate variations. In: Strack, M. \(Ed.\), *Peatlands and climate
 1046 change. International Peat Society, Jyväskylä, Finland, pp. 24-43.*](#)
 1047
- 1048 Maltby E and Immirzi P (1993) Carbon dynamics in peatlands and other wetland soils,
 1049 regional and global perspectives. *Chemosphere* 27: 999-1023.
 1050
- 1051 Marcott SA, Shakun JD, Clark PU and Mix AC (2013) A reconstruction of regional and
 1052 global temperature for the past 11,300 years. *Science* 339: 1198-1201.
 1053
- 1054 Martikainen PJ, Nykänen H, Crill P and Silvola J (1993) Effect of a lowered water-table
 1055 on nitrous-oxide fluxes from northern peatlands. *Nature* 366: 51-53.
 1056
- 1057 Marushchak ME, Pitkämäki A, Koponen H, Biasi C, Seppälä M and Martikainen PJ
 1058 (2011) Hot spots for nitrous oxide emissions found in different types of permafrost
 1059 peatlands. *Global Change Biology* 17: 2601-2614.
 1060
- 1061 Mauquoy D and van Geel B (2007) Mire and Peat Macros. In: Elias S.A. (Ed.),
 1062 *Encyclopedia of Quaternary Science*. vol. 3. Elsevier, Amsterdam, The Netherlands, pp.
 1063 2315-2336.
 1064
- 1065 McGuire AD, Anderson LG, Christensen TR, Dallimore S, Guo L, Hayes DJ, Heimann
 1066 M, Lorensen TD, MacDonald RW and Roulet N (2009) Sensitivity of the carbon cycle in
 1067 the Arctic to climate change. *Ecological Monographs* 79: 523-555.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1068
1069 McLaughlan KK, Williams JJ, Craine JM and Jeffers ES (2013) Changes in global
1070 nitrogen cycling during the Holocene epoch. *Nature* 495: 352-357.
1071
1072 Oksanen PO, Kuhry P and Alekseeva RN (2003) Holocene development and permafrost
1073 history of the Usinsk Mire, northeast European Russia. *Géographie Physique et*
1074 *Quaternaire* 57: 169-187.
1075
1076 Palmer K, Biasi C and Horn MA (2012) Contrasting denitrifier communities relate to
1077 contrasting N₂O emission patterns from acidic peat soils in arctic tundra. *The ISME*
1078 *Journal* 6: 1058-1077.
1079
1080 Pelletier L, Moore TR, Roulet NT, Garneau M and Beaulieu-Audy V (2007) Methane
1081 fluxes from three peatlands in the La Grande Rivière watershed, James Bay lowland,
1082 Canada. *Journal of Geophysical Research – Biogeosciences* 112, G01018. doi:
1083 10.1029/2006JG000216.
1084
1085 Peteet D, Andreev A, Bardeen W and Mistretta F (1998) Long-term Arctic peatland
1086 dynamics, vegetation and climate history of the Pur-Taz region, Western Siberia. *Boreas*
1087 27: 115-126.
1088
1089 Piotrowska N, Blaauw M, Mauquoy D and Chambers FM (2011) Constructing deposition
1090 chronologies for peat deposits using radiocarbon dating. *Mires and Peat* 7, article 10, 14
1091 p.
1092
1093 Post WM, Pastor J, Zinke PJ and Stangenberger AG (1985) Global patterns of soil
1094 nitrogen storage. *Nature* 317: 613-616.
1095
1096 Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Bertrand C, Blackwell PG, Buck
1097 CE, Burr G, Cutler KB, Damon PE, Edwards RL, Fairbanks RG, Friedrich M, Guilderson
1098 TM, Hughen KA, Kromer B, McCormac FG, Manning S, Bronk Ramsey C, Reimer RW,
1099 Remmele S, Southon JR, Stuiver M, Talamo S, Taylor FW, van der Plicht J,
1100 Weyhenmeyer CE (2009) IntCal09 and Marine09 radiocarbon age calibration curves, 0–
1101 50,000 years cal BP. *Radiocarbon* 51: 1111-1150.
1102
1103 Renssen H, Seppä H, Crosta X, Goose H and Roche DM (2012) Global characterization
1104 of the Holocene Thermal Maximum. *Quaternary Science Reviews* 48: 7-19.
1105
1106 Repo ME, Susiluoto S, Lind SE, Jokinen S, Elsakov V, Biasi C, Virtanen T, Martikainen
1107 PJ (2009) Large N₂O emissions from cryoturbated peat soil in tundra. *Nature Geoscience*
1108 2: 189-192.
1109
1110 Ruppel M, Välranta M, Virtanen T and Korhola A (2013) Postglacial spatiotemporal
1111 peatland initiation and lateral expansion dynamics in North America and northern
1112 Europe. *The Holocene*. doi:10.1177/0959683613499053.
1113

Formatted: Font: Italic

- 1114 Rydin H and Jeglum J (2013). *The biology of peatlands*. 2nd edition, Oxford University
 1115 Press, Oxford, UK, 382 p.
- 1116
- 1117 Sannel ABK and Kuhry P (2008) Long-term stability of permafrost in subarctic peat
 1118 plateaus, west-central Canada. *The Holocene* 18: 589-601.
- 1119
- 1120 Sheng Y, Smith LC, MacDonald GM, Kremenetski KV, Frey KE, Velichko AA, Lee M,
 1121 Beilman DW and Dubinin P (2004) A high-resolution GIS-based inventory of the West
 1122 Siberian peat carbon pool. *Global Biogeochemical Cycles* 18, GB3004. doi:
 1123 10.1029/2003GB002190.
- 1124
- 1125 Smith, L.C., MacDonald, G.M., Velichko, A.A., Beilman, D.W., Borisova, O.K., Frey,
 1126 K.E., Kremenetski, K.V., Sheng, Y. 2004. Siberian peatlands a net carbon sink and global
 1127 methane source since the Early Holocene. *Science*, 303, 353–356.
- 1128
- 1129 Stolbovoi V (2002) Carbon in Russian soils. *Climatic Change* 55: 131-156.
- 1130
- 1131 Stolbovoi V and McCallum I (2002) *Land resources of Russia*. International Institute for
 1132 Applied Systems Analysis. www.iiasa.ac.at/Research/FOR/russia_cd/download.htm.
- 1133
- 1134 Stolbovoi V, Montanarella L, Medvedev V, Smeyan N, Shishov L, Ungureanu V,
 1135 Dobrovolski G, Jamagne M, King D, Rozhkov V and Savin I (2001) Integration of data
 1136 on the soils of Russia, Byelorus, Moldova and Ukraine into the Soil Geographic Database
 1137 of the European Community. *Eurasian Soil Science* 34: 687-703.
- 1138
- 1139 Stuiver M and Reimer PJ (1993) Extended ¹⁴C database and revised CALIB radiocarbon
 1140 calibration program. *Radiocarbon* 35: 215-230.
- 1141
- 1142 Telford RJ, Heegaard E and Birks HJB (2004) All age-depth models are wrong: but how
 1143 badly? *Quaternary Science Reviews* 23: 1-5.
- 1144
- 1145 Troels-Smith J (1955) Characterization of unconsolidated sediments. Danmarks
 1146 Geologiske *Undersøgelse Series* 4: 1-73.
- 1147
- 1148 Turetsky MR, Wieder RK, Vitt DH, Evans RJ and Scott KD (2007) The disappearance of
 1149 relict permafrost in boreal North America: effects on peatland carbon storage and fluxes.
 1150 *Global Change Biology* 13: 1922-1934.
- 1151
- 1152 Turunen J, Tomppo E, Tolonen K and Reinikainen A (2002) Estimating carbon
 1153 accumulation rates of undrained mires in Finland – application to boreal and subarctic
 1154 regions. *The Holocene* 12: 69-80.
- 1155
- 1156 Turunen J, Roulet NT, Moore TR and Richard PJH (2004) Nitrogen deposition and
 1157 increased carbon accumulation in ombrotrophic peatlands in eastern Canada. *Global*
 1158 *Biogeochemical Cycles*, 18, GB3002. doi: 10.1029/2003GB002154.
- 1159

Formatted: Superscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

van Bellen S, Dallaire P-L, Garneau M and Bergeron Y (2011) Quantifying spatial and temporal Holocene carbon accumulation in ombrotrophic peatlands of the Eastmain region, Quebec, Canada. *Global Biogeochemical Cycles* 25, GB2016. doi:10.1029/2010GB003877.

Vitt DH, Halsey LA, Bauer IE and Campbell C (2000) Spatial and temporal trends in carbon storage of peatlands of continental western Canada through the Holocene. *Canadian Journal of Earth Sciences* 37: 683-693.

Yu Z (2011) Holocene carbon flux histories of the world's peatlands: global carbon-cycle implications. *The Holocene* 21: 761-774.

Yu Z (2012) Northern peatland carbon stocks and dynamics: a review. *Biogeosciences* 9: 4071-4085.

Yu Z, Campbell ID, Campbell C, Vitt DH, Bond GC and Apps MJ (2003) Carbon sequestration in western Canadian peat highly sensitive to Holocene wet-dry climate cycles at millennial timescales. *The Holocene* 13: 801-808.

Yu Z, Beilman DW and Jones MC (2009) Sensitivity of northern peatland carbon dynamics to Holocene climate change. In: Baird A, Belyea L, Comas X, Reeve A, Slater L (Eds.), *Northern peatlands and carbon cycling*. American Geophysical Union Monograph Series, Washington D.C., USA, pp. 55-69.

Yu Z, Loisel J, Brosseau DP, Beilman DW and Hunt SJ (2010) Global peatland dynamics since the Last Glacial Maximum. *Geophysical Research Letters* 37, L13402. doi:10.1029/2010GL043584.

Yu Z, Loisel J, Turetsky MR, Cai S, Zhao Y, Frolking S, MacDonald GM and Bubier JL (2013) Evidence for elevated emissions from high-latitude wetlands contributing to high atmospheric CH₄ concentration in the early Holocene. *Global Biogeochemical Cycles* 27: 131-140. doi:10.1002/GBC.20025.

Zoltai SC (1971) Southern limit of permafrost features in peat landforms, Manitoba and Saskatchewan. *Geological Association of Canada, Special Paper* 9, 305-310.

Zoltai SC (1995) Permafrost distribution in peatlands of west-central Canada during the Holocene warm period 6000 years B.P. *Géographie Physique et Quaternaire* 49: 45-54.

Figure and Table Captions

Figure 1. Location of study sites. Map showing the distribution of northern peatlands (green area from Yu et al., 2010) and peatland sites included in this study (n = 215 sites,

including 268 peat cores). Long-term rate of peat-carbon accumulation was estimated from 127 sites (151 peat cores; red and blue dots). The yellow dots represent cores for which only peat properties (bulk density, organic matter content, etc.) were available and synthesized. Refer to Supplementary Table 1 for details.

Figure 2. Overview of data availability for North America (black bars) and Eurasia (white bars). (A) Number of cores (total = 238) containing information on carbon/nitrogen ratio (n = 40), nitrogen content (n = 40), carbon content (n = 56), peat types (n = 83), organic matter bulk density (n = 184), organic matter content (n = 190), and bulk density (n = 214). (B) Number of cores (total = 151) with a ~~temporal-dating~~ resolution-quality better than ~~500-year~~two dates per 1000 years (n = 35), ~~500-1000~~one to two dates per 1000 years (n = 52), and less than one date per 1000 years (n = 64). (C) Number of calibrated basal peat ages (median) in 500-year bins from the database (~~black bars~~, n = 199) compared to all northern hemisphere basal peat ages (median) in 200-year bins (~~grey bars~~, n = 2559, MGK data from MacDonald et al., 2006, Gorham et al., 2007, Korhola et al., 2010).

Figure 3. Distribution histograms of peat properties in northern peatlands. (A) Frequency distribution of bulk density for unidentified peat type samples (white bars) and different peat types (color bars). (B) Frequency distribution of organic matter content for different peat types. (C) Frequency distribution of organic matter bulk density for different peat types. (D) Frequency distribution of carbon content for different peat types. (E)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1224 Frequency distribution of nitrogen content for different peat types. (F) Frequency
1225 distribution of carbon/nitrogen mass ratio for different peat types.
1226

1227 **Figure 4.** Relation between carbon content and organic matter content in northern
1228 peatlands. The slope of each regression line is used as a conversion factor for estimating
1229 carbon content from organic matter content.

1230
1231 **Figure 5.** Temporal patterns of peat properties (mean, standard deviation, and number of
1232 samples). (A) Bulk density. (B) Organic matter content. (C) Organic matter bulk density.
1233 (D) Carbon content. (E) Nitrogen content. (F) Carbon/Nitrogen mass ratio. White bars
1234 represent values that were based on a limited number of samples and peat records.

1235
1236 **Figure 6.** Main differences between *Sphagnum* and non-*Sphagnum* peat samples. (A)
1237 Bulk density. (B) Organic matter bulk density. (C) Organic carbon bulk density. (D)
1238 Carbon content. (E) Nitrogen content. (F) Carbon/Nitrogen mass ratio. (G) Temporal
1239 pattern of organic C bulk density. (H) Proportional change in the number of peat records
1240 that are *Sphagnum*-dominated, presented as a percentage of the total number of records.

1241
1242 **Figure 7.** Long-term apparent rate of carbon and nitrogen accumulation from northern
1243 peatlands (n = 127 sites). (A) Summer insolation at 60°N (data from Berger and Loutre,
1244 1991) and temperature anomaly from an 11,300-year reconstruction for the northern
1245 extra-polar region from 30 to 90°N (data from Marcott et al., 2013). Temperature
1246 anomaly was calculated based on the 1961-1990 temperature averages. (B) Mean peat-

1
2
3
4
5
6
7
8 1247 carbon accumulation rates (PCAR) and standard error in 500-year bins. The number of
9
10 1248 sites per 500-year bins is also presented. (C) Mean peat-nitrogen accumulation rates
11
12 1249 (PNAR) and standard error in 500-year bins. These values were obtained using different
13
14 1250 C/N values over time, as indicated by the line. ~~PCAR: peat carbon accumulation rate;~~
15
16 1251 ~~PNAR: peat N accumulation rate.~~
17
18 1252
19
20 1253 **Table 1.** Peat properties in northern peatlands. Means and standard deviations are
21
22 1254 presented, along with the number of samples (n).
23
24 1255
25
26 1256 **Table 2.** Northern peatland peat properties by regions. Means and standard deviations are
27
28 1257 presented, along with the number of samples (n).
29
30 1258
31
32 1259 **Table 3.** Comparison of northern peatland peat properties estimates with other published
33
34 1260 values. Means and standard deviations are presented, along with the number of samples
35
36 1261 (n) when available.
37
38 1262
39 1263 **Supplementary Material**
40
41 1264 **Table S1.** Summary information for the study sites included in the database.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Location of study sites. Map showing the distribution of northern peatlands (green area from Yu et al., 2010) and peatland sites included in this study (n = 215 sites, including 268 peat cores). Long-term rate of peat-carbon accumulation was estimated from 127 sites (151 peat cores; red and blue dots). The yellow dots represent cores for which only peat properties (bulk density, organic matter content, etc.) were available and synthesized. Refer to Supplementary Table 1 for details.

279x361mm (300 x 300 DPI)

Figure 2. Overview of data availability for North America (black bars) and Eurasia (white bars). (A) Number of cores (total = 238) containing information on carbon/nitrogen ratio ($n = 40$), nitrogen content ($n = 40$), carbon content ($n = 56$), peat types ($n = 83$), organic matter bulk density ($n = 184$), organic matter content ($n = 190$), and bulk density ($n = 214$). (B) Number of cores (total = 151) with a dating quality better than two dates per 1000 years ($n = 35$), one to two dates per 1000 years ($n = 52$), and less than one date per 1000 years ($n = 64$). (C) Number of calibrated basal peat ages (median) in 500-year bins from the database ($n = 199$) compared to all northern hemisphere basal peat ages (median) in 200-year bins ($n = 2559$, MGK data from MacDonald et al., 2006, Gorham et al., 2007, Korhola et al., 2010).

Figure 3. Distribution histograms of peat properties in northern peatlands. (A) Frequency distribution of bulk density for unidentified peat type samples (white bars) and different peat types (color bars). (B) Frequency distribution of organic matter content for different peat types. (C) Frequency distribution of organic matter bulk density for different peat types. (D) Frequency distribution of carbon content for different peat types. (E) Frequency distribution of nitrogen content for different peat types. (F) Frequency distribution of carbon/nitrogen mass ratio for different peat types.

279x361mm (300 x 300 DPI)

Figure 4. Relation between carbon content and organic matter content in northern peatlands. The slope of each regression line is used as a conversion factor for estimating carbon content from organic matter content.

279x361mm (300 x 300 DPI)

Figure 5. Temporal patterns of peat properties (mean, standard deviation, and number of samples). (A) Bulk density. (B) Organic matter content. (C) Organic matter bulk density. (D) Carbon content. (E) Nitrogen content. (F) Carbon/Nitrogen mass ratio. White bars represent values that were based on a limited number of samples and peat records.

Figure 6. Main differences between Sphagnum and non-Sphagnum peat samples. (A) Bulk density. (B) Organic matter bulk density. (C) Organic carbon bulk density. (D) Carbon content. (E) Nitrogen content. (F) Carbon/Nitrogen mass ratio. (G) Temporal pattern of organic C bulk density. (H) Proportional change in the number of peat records that are Sphagnum-dominated, presented as a percentage of the total number of records.

279x361mm (300 x 300 DPI)

Figure 7. Long-term apparent rate of carbon and nitrogen accumulation from northern peatlands (n = 127 sites). (A) Summer insolation at 60°N (data from Berger and Loutre, 1991) and temperature anomaly from an 11,300-year reconstruction for the northern extra-polar region from 30 to 90°N (data from Marcott et al., 2013). Temperature anomaly was calculated based on the 1961-1990 temperature averages. (B) Mean peat-carbon accumulation rates (PCAR) and standard error in 500-year bins. The number of sites per 500-year bins is also presented. (C) Mean peat-nitrogen accumulation rates (PNAR) and standard error in 500-year bins. These values were obtained using different C/N values over time, as indicated by the line.

Table 1. **Peat properties in northern peatlands.** Means and standard deviations are presented, along with the number of samples (*n*).

	<i>Sphagnum</i>	Herbaceous	Woody	Humified	Brown Moss	Overall***
Bulk density (g cm⁻³)	0.076 ± 0.038 (<i>n</i> = 4372)	0.118 ± 0.075 (3188)	0.108 ± 0.047 (1584)	0.192 ± 0.082 (452)	0.177 ± 0.076 (1114)	0.118 ± 0.069 (21,220)
Organic matter content (%)	94.3 ± 9.3 (3297)	85.6 ± 15.4 (3121)	92.0 ± 13.5 (1587)	78.4 ± 17.8 (418)	81.4 ± 15.5 (1090)	90.7 ± 13.0 (18,973)
Organic matter bulk density (g OM cm⁻³)	0.073 ± 0.031 (3332)	0.089 ± 0.036 (2854)	0.098 ± 0.032 (1388)	0.144 ± 0.036 (418)	0.136 ± 0.043 (1090)	0.105 ± 0.051 (18,544)
Carbon content in total peat (%)	46.0 ± 4.1 (1520)	50.5 ± 4.9 (519)	50.9 ± 4.0 (308)	47.4 ± 4.1 (96)	47.9 ± 2.8 (72)	46.8 ± 6.1 (3741)
Carbon content in organic matter (%)	42.3 ± 3.0* (454)	51.1 ± 1.7* (147)	51.4 ± 3.4* (59)	53.2 ± 2.6* (58)	50.0 ± 2.0* (44)	49.2 ± 2.4** (458)
Nitrogen content in peat (%)	0.7 ± 0.3 (1523)	1.7 ± 0.6 (518)	1.3 ± 0.5 (308)	1.5 ± 0.4 (96)	1.4 ± 0.7 (60)	1.2 ± 0.7 (3365)
Carbon/Nitrogen mass ratio	81.0 ± 49.2 (1520)	34.4 ± 15.0 (518)	45.3 ± 19.1 (308)	36.0 ± 17.6 (96)	42.9 ± 18.8 (60)	55 ± 33 (3362)

*Obtained from regression between carbon content and organic matter content (see the Database and analysis section).

**Includes all herbaceous, woody, humified and brown moss samples, as well as *Sphagnum* samples older than 0.5 ka (see Results section).

***Includes samples for which peat type was not ascribed.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 2. **Northern peatland peat properties by regions.** Means and standard deviations are presented, along with the number of samples in parentheses (n).

	Alaska	Western Canada	Hudson & James Bays	Eastern Canada/USA	Western European Islands	Continental Europe	Fennoscandia	Western Russia	Eastern Russia & Asia
Bulk density (g cm ⁻³)	0.168 ± 0.087 (n = 1659)	0.166 ± 0.076 (3635)	0.097 ± 0.038 (6002)	0.100 ± 0.039 (2834)	0.055 ± 0.027 (656)	0.120 ± 0.139 (410)	0.075 ± 0.043 (562)	0.118 ± 0.070 (2701)	0.116 ± 0.063 (2761)
Organic matter content (%)	76.6 ± 18.8 (1659)	91.6 ± 8.1 (3442)	94.8 ± 8.2 (5129)	97.8 ± 6.5 (1835)	97.5 ± 1.8 (227)	97.4 ± 5.43 (305)	95.6 ± 8.7 (789)	94.6 ± 10.3 (2666)	80.3 ± 16.7 (2700)
Organic matter bulk density (g OM cm ⁻³)	0.119 ± 0.049 (1659)	0.151 ± 0.062 (3441)	0.088 ± 0.029 (5129)	0.107 ± 0.028 (1750)	0.055 ± 0.035 (227)	0.056 ± 0.028 (222)	0.073 ± 0.034 (422)	0.106 ± 0.058 (2773)	0.088 ± 0.034 (2700)
Carbon content in total peat (%)	42.4 ± 3.7 (64)	45.0 ± 4.3 (382)	47.9 ± 4.5 (1026)	48.9 ± 3.7 (1084)	54.0 ± 2.5 (242)	38.9 ± 1.3 (60)	44.4 ± 5.7 (580)	49.2 ± 3.2 (74)	36.0 ± 9.2 (229)
Nitrogen content in peat (%)	1.3 ± 0.6 (64)	1.1 ± 0.8 (265)	1.6 ± 0.7 (910)	0.9 ± 0.5 (1084)	1.6 ± 0.4 (242)	0.7 ± 0.1 (60)	1.0 ± 0.5 (565)	1.6 ± 0.9 (44)	1.4 ± 0.6 (131)
Carbon/Nitrogen mass ratio	43.9 ± 32.8 (64)	62.4 ± 37.5 (265)	39.5 ± 23.7 (910)	77.2 ± 56.1 (1084)	35.7 ± 10.8 (242)	54.2 ± 7.6 (60)	57.9 ± 31.4 (562)	40.8 ± 21.7 (44)	34.2 ± 21.9 (131)

Table 3. Northern peatland peat properties estimates from published studies. Means and standard deviations are presented, along with the number of samples in parentheses (*n*) when available.

Bulk density (g cm ⁻³)	Organic matter content (%)	Organic matter bulk density (g OM cm ⁻³)	Carbon content in organic matter (%)	Carbon/Nitrogen mass ratio	Region	Reference
-	-	0.094 open fens & bogs 0.105 wooded & shrubby fens	51.8 ± 4.7 (<i>n</i> = 253)	-	Western Canada	Vitt et al., 2000
0.073 ± 0.029 <i>Sphagnum</i> 0.091 ± 0.025 brown moss 0.110 ± 0.037 sedge-moss 0.211 ± 0.061 humified 0.138 ± 0.036 wood	95.5 ± 2.6 <i>Sphagnum</i> 90.3 ± 6.6 brown moss 91.4 ± 4.4 sedge-moss 73.6 ± 13.0 humified 87.8 ± 6.3 wood	0.069 ± 0.028 <i>Sphagnum</i> 0.082 ± 0.023 brown moss 0.100 ± 0.032 sedge-moss 0.149 ± 0.023 humified 0.120 ± 0.029 wood	50.7 ± 5.0 <i>Sphagnum</i> 51.9 ± 3.4 brown moss 53.4 ± 2.9 sedge-moss 54.0 ± 3.8 humified 52.1 ± 3.5 wood	-	Western Canada	Bauer et al., 2006
-	-	0.0784 bogs	52.8 (<i>n</i> = 276)	-	Eastern Canada and USA	Gorham, 1990
-	-	0.112	51.7	-	Eastern Canada and USA	Gorham, 1991
0.128 ± 0.065	96.26 ± 3.16	0.123*	52	-	West Siberia Lowlands	Sheng et al., 2004
-	-	-	51 ± 5 <i>Sphagnum</i> ** 55 ± 3 non- <i>Sphagnum</i> ** 52 ± 3 overall**	-	West Siberia Lowlands	Beilman et al., 2009
-	-	0.074 bogs 0.081 fens	50	-	Finland	Turunen et al., 2002
0.118 ± 0.069 (<i>n</i> = 21,220)	90.7 ± 13.0 (<i>n</i> = 18,973)	0.105 ± 0.051 (<i>n</i> = 18,544)	49.2 ± 2.4 (<i>n</i> = 458)	55 ± 33 (<i>n</i> = 3362)	circum-Arctic	This study

*This value was obtained by multiplying bulk density (0.128 g cm⁻³) by organic matter content (96.26%).

**Standard errors

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Supplementary Material: Table S1. Summary information for the study sites included in the circum-Arctic peatland database.

Abbreviated reference*	Core name and ID	Peatland type	Country	Latitude	Longitude	Number of ¹⁴ C dates	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
<u>NORTH AMERICA</u>											
Beaulieu-Audy, 2009	La Grande 3	Bog	Canada	53.57	-76.13	7	6816	Y	N	Y ³	Y
Beaulieu-Audy, 2009	La Grande 2	Bog	Canada	53.65	-77.73	6	6543	Y	N	Y ³	Y
Beaulieu-Audy, 2009	La Grande 1	Intermediate fen	Canada	53.9	-78.77	8	1612	Y	N	Y ³	Y
Belyea, 1996	Rainy River Bog	Bog	Canada	48.78	-94.55	9	5310	N	N	Y ⁵	N
Bender, 1969	Porcupine	Bog	Canada	52.52	-101.25	7	7624	Y	N	Y ⁵	N
Bender, 1969	Colville Lake	Bog	Canada	67.1	-125.78	7	7650	Y	N	Y ⁵	N
Booth, 2004	South Rhody	Kettle bog	USA	46.55	-86.07	9	10,562	Y	N	Y ⁵	N
Bunbury, 2012	VC04-06	Shrub bog	Canada	52.71	-84.18	6	6599	Y	N	Y ²	Y
Camill, 2009	Joey Lake 5	Permafrost bog	Canada	55.46	-98.16	8	8100	Y ¹	N	Y ²	Y
Camill, 2009	Joey Lake 7	Permafrost bog	Canada	55.46	-98.16	10	8256	Y	N	Y ²	Y
Camill, 2009	Joey Lake 2	Permafrost bog	Canada	55.47	-98.16	11	7980	Y ¹	N	Y ²	Y
Camill, 2009	Joey Lake 12	Permafrost bog	Canada	55.47	-98.15	8	6564	Y ¹	N	Y ²	Y
Camill, 2009	Joey Lake 15	Permafrost bog	Canada	55.47	-98.15	10	7882	Y ¹	N	Y ²	Y
Camill, 2009	Joey Lake 17	Permafrost bog	Canada	55.47	-98.16	7	7632	Y ¹	N	Y ²	Y
P. Camill, unpubl	Lake 785 core 4	Permafrost bog	Canada	59.11	-97.4	9	6833	Y	N	Y ²	Y
P. Camill, unpubl	Unit Lake core 4	Permafrost bog	Canada	59.42	-97.48	9	7053	Y	N	Y ²	Y
P. Camill, unpubl	Lake 396 core 3	Permafrost bog	Canada	59.58	-98.57	6	6077	Y	N	Y ²	Y
P. Camill, unpubl	Shuttle Lake core 2	Permafrost bog	Canada	59.86	-97.64	6	6242	Y	N	Y ²	Y
Charman, 1995	Wally Creek Area	Bog	Canada	49.07	-80.6	10	6672	Y	N	Y ⁵	N
P. Charman, unpubl	Burnt Village	Raised bog	Canada	51.13	-55.93	26	8526	Y	N	Y ²	Y
P. Charman, unpubl	Petite Bog	Raised bog	Canada	45.14	-63.94	32	13,474	Y	N	Y ²	Y
P. Charman, unpubl	Sidney Bog	Raised bog	USA	44.39	-69.79	31	9311	Y	N	Y ²	Y
Elliott, 2011	Mer Bleue	Bog	Canada	45.68	-75.8	11	8463	Y	N	Y ⁵	N

1
2
3
4

	Abbreviated reference ^a	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
8	M. Garneau, unpubl	Ours 1	Fen	Canada	54.05	-72.45	6	5491	N	N	Y ³	Y
9	M. Garneau, unpubl	Ours 3	Fen	Canada	54.05	-72.46	5	3899	N	N	Y ³	Y
10	M. Garneau, unpubl	Ours 4	Fen	Canada	54.05	-72.46	6	4774	N	N	Y ³	Y
11	M. Garneau, unpubl	Aero 1	Fen	Canada	54.1	-72.52	6	5485	N	N	Y ³	Y
12	M. Garneau, unpubl	Aero 5	Fen	Canada	54.1	-72.52	6	4252	N	N	Y ³	Y
14	Glaser, 2004	Oldman Bog	Bog	Canada	51.02	-84.57	13	6728	Y	N	Y ⁵	N
15	Glaser, 2004	Albany River Bog	Bog	Canada	51.43	-83.62	6	5492	Y	N	Y ⁵	N
16	Glaser, 2004	Belec Lake Bog	Bog	Canada	51.62	-82.28	7	4480	Y	N	Y ⁵	N
18	Gorham, 2003 ^a	Miscou	Bog	Canada	47.93	-64.5	7	9000	Y	N	Y ⁶	N
19	Gorham, 2003 ^a	Fourchou	Bog	Canada	45.93	-60.27	8	11,200	Y	N	Y ⁶	N
20	Gorham, 2003	Denbigh	Fen	USA	48.22	-100.5	8	12,500	Y	N	Y ⁶	N
22	Holmquist, unpubl	JBL8	Sphagnum bog	Canada	50.47	-89.93	11	4481	Y	N	Y ³	Y
23	Holmquist, unpubl	JBL1	Sphagnum bog	Canada	51.07	-89.8	11	6034	Y	N	Y ³	Y
24	Holmquist, unpubl	JBL2	Sphagnum bog	Canada	52.02	-90.13	13	6742	Y	N	Y ³	Y
25	Holmquist, unpubl	JBL3	Sphagnum bog	Canada	52.87	-89.93	10	7708	Y	N	Y ³	Y
27	Holmquist, unpubl	JBL7	Sphagnum bog	Canada	54.4	-89.52	12	7607	Y	N	Y ³	Y
28	Holmquist, unpubl	JBL6	Permafrost bog	Canada	54.77	-89.32	8	3248	Y	N	Y ³	Y
29	Holmquist, unpubl	JBL4	Sphagnum bog	Canada	55.27	-88.93	11	6051	Y	N	Y ³	Y
30	Holmquist, unpubl	JBL5	Peat plateau	Canada	55.42	-88.95	12	5826	Y	N	Y ³	Y
32	Hu, 1994	Caribou Bog RC-2	Bog	USA	45	-69	6	9707	Y	pollen (1)	Y ⁵	Y
33	Hughes, 2006	Nordan's Pond Bog	Bog	Canada	53.6	-49.17	10	8827	N	N	Y ⁵	N
34	Hunt, 2013	Nuikluk 10-1	Peat plateau	USA (Alaska)	64.83	-163.45	5	6392	Y ¹	N	Y ³	Y
35	Hunt, 2013	Nuikluk 10-2	Collapse Scar	USA (Alaska)	64.83	-163.45	9	13,545	Y	N	Y ³	Y
37	Jones, 2010 ^a	Horse Trail Fen	Poor fen	USA (Alaska)	60.42	-150.9	11	12,695	Y	N	Y ³	Y
38	Jones, 2010 ^a	Kenai Gasfield 07-2	Poor Fen	USA (Alaska)	60.45	-151.25	17	11,448	Y	N	Y ³	Y
39	Jones, 2010 ^a	No Name Creek 07-1	Poor Fen	USA (Alaska)	60.63	-151.08	10	10,993	Y	N	Y ³	Y

41
42
43
44
45
46
47
48
49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Abbreviated reference ^a	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
Jones, 2010 ^a	Swanson Fen	Poor fen	USA (Alaska)	60.79	-150.83	10	14,065	Y	N	Y ³	Y
Klein, 2013	Kahiltna Valley Mor.	Bog	USA (Alaska)	62.37	-151.09	5	1949	N	N	Y ³	Y
E. Klein, unpubl	HERC 09-3	Bog	USA (Alaska)	62.37	-151.07	8	11,768	Y	N	Y ³	Y
Kuhry, 1996 ^a	Slave Lake Bog	Bog	Canada	55.01	-114.09	6	10,516	Y	N	Y ²	Y
Lamarre, 2012	KUJU-PD2	Permafrost bog	Canada	55.23	-77.7	8	5084	Y	N	Y ³	Y
Lavoie, 2000	Lac Malbaie MAL-2	Bog	Canada	47.6	-70.97	5	10,654	Y	N	Y ⁵	N
Lavoie, 2000	Frontenac FRON-2	Bog	Canada	45.97	-71.13	7	12,851	Y	N	Y ⁵	N
Lavoie, 2013	Covey Hill	Bog	Canada	45.00	-73.49	12	12,720	Y	N	Y ³	Y
Loisel, 2010	Lac Le Caron RiP2	Bog	Canada	52.28	-75.83	6	2731	N	N	Y ²	Y
Loisel, 2013	Petersville 08-S	Bog	USA (Alaska)	62.42	-150.68	6	2825	N	tephra (1)	Y ²	Y
J. Loisel, unpubl	Petersville 09-MC	Bog	USA (Alaska)	62.42	-150.68	12	13,881	Y	tephra (4)	Y ³	Y
MacDonald, 1983	Natla River Bog	Bog	Canada	63.02	-128.8	6	9747	Y	tephra (1)	Y ⁵	Y
Magnan, 2012	Radisson	Semi-forested bog	Canada	53.73	-77.7	6	6154	Y	N	Y ⁵	N
G. Magnan, unpubl	Lebel	Raised bog	Canada	49.1	-68.25	12	5831	Y	N	Y ³	Y
G. Magnan, unpubl	Baie	Raised bog	Canada	49.1	-68.22	9	4221	Y	N	Y ³	Y
G. Magnan, unpubl	Morts	Peat plateau	Canada	50.26	-63.67	10	3246	Y	N	Y ³	Y
G. Magnan, unpubl	Plaine	Peat plateau	Canada	50.27	-63.54	12	7451	Y	N	Y ³	Y
Muller, 2003 ^a	Mirabel bog (7 cores)	Bog	Canada	45.68	-74.03	2 to 7	10,000	Y	N	Y ⁶	N
J. Nichols, unpubl	Bear Bog	Bog	USA (Alaska)	60.53	-145.45	13	10357	Y	N	Y ³	Y
O'Donnell, 2012	Koyukuk Flats PP2	Peat plateau	USA (Alaska)	65.19	-155.36	7	12,329	Y	N	Y ⁵	N
O'Reilly, 2011	Victor Fen	Fen	Canada	52.71	-84.17	6	6405	Y	N	Y ⁵	N
M. Packalen, unpubl	HL-02	Patterned bog	Canada	54.61	-84.61	5	4494	Y	N	Y ²	Y
Robinson, 2006 ^a	Martin River	Bog	Canada	61.8	-121.4	6	7552	Y	N	Y ⁴	Y
Sannel, 2009 ^a	Selwyn Lake 1	Peat plateau	Canada	59.88	-104.2	14	6573	Y	N	Y ²	Y
C. Tarnocai, 2010	T5	Polygon bog	Canada	68.57	-133.50	6	8805	Y	N	Y ²	Y
C. Tarnocai, 2010	IN-BG-1	Polyg. peat plateau	Canada	68.32	-133.42	9	9121	Y	N	Y ²	Y

1
2
3
4

	Abbreviated reference ^a	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
8	C. Tarnocai, 2010	IN-BG-3	Polyg. peat plateau	Canada	68.32	-133.43	6	6293	Y	N	Y ²	Y
9	C. Tarnocai, 2010	NW-BG-8	Polyg. peat plateau	Canada	65.21	-127.01	5	10,818	Y	N	Y ²	Y
10	C. Tarnocai, 2010	NW-BG-10	Polyg. peat plateau	Canada	65.21	-127.00	5	10,480	Y	N	Y ²	Y
11	C. Tarnocai, unpubl	T1	Polygon bog	Canada	68.32	-133.42	7	8623	Y	N	Y ²	Y
12	C. Tarnocai, unpubl	T6	Polygon bog	Canada	69.12	-134.18	5	3014	Y	N	Y ²	Y
13	C. Tarnocai, unpubl	IN-BG-2B	Polyg. peat plateau	Canada	68.32	-133.43	6	5828	Y	N	Y ²	Y
14	C. Tarnocai, unpubl	NW-BG-2	Polyg. peat plateau	Canada	65.21	-127.01	5	10,932	Y	N	Y ²	Y
15	C. Tarnocai, unpubl	NW-BG-3	Polyg. peat plateau	Canada	65.21	-127.01	6	11,010	Y	N	Y ²	Y
16	C. Tarnocai, unpubl	NW-BG-3	Polyg. peat plateau	Canada	65.21	-127.01	6	11,010	Y	N	Y ²	Y
17	Turunen, 2003 ^a	Diana Lake bog	Slope bog	Canada	54.15	-130.25	5	8500	Y	N	Y ⁶	N
18	van Bellen, 2011	Mosaik	Bog	Canada	51.98	-75.4	10	7120	Y	N	Y ³	Y
19	van Bellen, 2011	Sterne	Bog	Canada	52.05	-75.17	11	7134	Y	N	Y ³	Y
20	van Bellen, 2011	Lac Le Caron	Bog	Canada	52.28	-75.83	12	7510	Y	N	Y ³	Y
21	Yu, 2003 ^a	Upper Pinto Fen	Rich fen	Canada	53.58	-118.02	20	7599	Y	N	Y ³	Y
22	Yu, 2006 ^a	Goldeye Lake Fen	Rich fen	Canada	52.45	-116.2	6	9207	Y	tephra (2)	Y ³	Y
23	Z. Yu, unpubl	Sundance Fen 03-2	Rich fen	Canada	53.58	-116.75	5	6719	Y ¹	N	Y ³	Y
24	Z. Yu, unpubl	Sundance Fen 03-3	Rich fen	Canada	53.58	-116.75	13	10,973	Y	N	Y ³	Y
25	Z. Yu, unpubl	Utikuma	Poor Fen	Canada	55.84	-115.09	18	5079	Y	N	Y ³	Y
26	Z. Yu, unpubl	Mariana Lake 03-1	Poor Fen	Canada	55.9	-112.09	14	7222	Y	N	Y ³	Y
27	Z. Yu, unpubl	Mariana Lake 03-2	Poor Fen	Canada	55.9	-112.09	11	6105	Y ¹	N	Y ³	Y
28	Z. Yu, unpubl	Mariana Lake 03-3	Poor Fen	Canada	56.02	-111.93	18	5872	Y ¹	N	Y ³	Y
29	Z. Yu, unpubl ^a	Patuanak	Internal lawn	Canada	55.85	-107.68	11	9017	Y	N	Y ³	Y
30	M. Garneau, unpubl	Ours 5	Fen	Canada	54.05	-72.46	3	5958	N	N	N	Y
31	M. Garneau, unpubl	Ours 2	Fen	Canada	54.05	-72.46	2	3496	N	N	N	Y
32	M. Garneau, unpubl	Aero 3	Fen	Canada	54.1	-72.52	2	3387	N	N	N	Y
33	Hu, 1994	Caribou Bog RC-1	Bog	USA	45	-69	2	9547	Y ¹	pollen (3)	N	Y
34	Lamarre, 2012	KUJU-BF2	Permafrost bog	Canada	55.23	-77.7	4	3914	Y	N	N	Y

49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Abbreviated reference ^a	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
Loisel, 2010	Mosaik RiP2	Bog	Canada	51.97	-75.4	4	2433	N	N	N	Y
M. Paackalen, unpubl	KJ2-3	Poor fen	Canada	51.59	-81.76	4	4677	Y	N	N	Y
Robinson, 2000	Peat Plateau LC	Peat Plateau	Canada	61.8	-121.4	0		N	tephra (1)	N	Y
Robinson, 2000	Peat Plateau 13	Peat Plateau	Canada	61.8	-121.4	0		N	tephra (1)	N	Y
Robinson, 2000	Poor Fen 11	Poor Fen	Canada	61.8	-121.4	0		N	tephra (1)	N	Y
Robinson, 2000	Rich Fen 12	Rich fen	Canada	61.8	-121.4	0		N	tephra (1)	N	Y
Robinson, 2000	Unfrozen Bog 10	Permafrost bog	Canada	61.8	-121.4	0		N	tephra (1)	N	Y
Robinson, 2000	Collapse Scar Fen 06	Collapse Scar	Canada	61.8	-121.4	0		N	tephra (1)	N	Y
Sannel, 2009	Ennadai Lake 1	Peat plateau	Canada	60.83	-101.55	4	5792	Y	N	N	Y
C. Tarnocai, unpubl	NW-BG-4	Polyg. peat plateau	Canada	65.21	-127.01	3	9916	Y	N	N	Y
C. Tarnocai, unpubl	NW-BG-9	Polyg. peat plateau	Canada	65.23	-127.00	3	9575	Y	N	N	Y
Z. Yu, unpubl	Hondo	Rich fen	Canada	55.08	-114.14	4	10,012	Y	N	N	Y
EURASIA											
Anderson, 1998 ^a	Glen Torridon	Olig. topogen. bog	UK	57.56	-5.37	7	9568	Y	N	Y ²	Y
Anderson, 1998 ^a	Glen Carron	Olig. topogen. bog	UK	57.53	-5.15	6	10,431	Y	N	Y ²	Y
Andersson, 2010	Lilla Backsjömyren 1	Mixed mire	Sweden	62.41	14.32	5	8527	Y	tephra (2)	Y ⁵	Y
Andersson, 2010	Lilla Backsjömyren 2	Mixed mire	Sweden	62.41	14.32	13	3804	Y ¹	tephra (2)	Y ⁵	Y
Barber, 2003	Bolton Fell Moss J,L	Bog	UK	55	-2	28	10,476	Y	N	Y ⁵	N
Barber, 2003	Mongan Bog	Bog	Ireland	53	-8	13	4607	N	N	Y ⁵	N
Barber, 2003	Abbeyknockmoy Bog	Bog	Ireland	53.5	-9	10	6707	N	N	Y ⁵	N
C. Bocchicchio, unpubl.	KAM12-C4	Bog	Russia (Far-E)	54.01	156.08	10	12,891	Y	N	Y ³	Y
Borren, 2004 ^a	Vasyugan (V21)	Bog	Russia (Siberia)	56.83	78.42	11	9709	Y	N	Y ³	Y
Borren, 2004 ^a	86-Kvartal (Zh0)	Fen	Russia (Siberia)	56.83	84.58	9	8711	Y	N	Y ³	Y
Charman, 1994	East Southerland	Fen	UK	58	-3	6	10,084	Y	N	Y ⁵	N
D. Vleeschouwer, 2009	Ślowińskie Błota	Raised bog	Poland	54.36	16.49	8	1165	N	N	Y ²	Y
D. Vleeschouwer, 2012	Misten	Raised bog	Belgium	50.56	6.16	15	1434	N	N	Y ²	Y

1
2
3
4

	Abbreviated reference ^a	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
8	Galka, 2013a	Stažki-B	Bog	Poland	54.43	18.09	9	7352	Y	N	Y ⁵	Y
9	Galka, 2013b	Kusowo	Raised bog	Poland	54	18	8	578	N	N	Y ⁵	N
10	Glebov, 2002	Ob-Vasygan	Bog	Russia (Siberia)	60.52	77.68	17	10,817	Y	N	Y ⁵	N
11	Hendon, 2001	Butternburn Flow 1	Intermed. ombrotr	UK	55.08	-2.5	7	9,213	Y	N	Y ⁵	N
12	Hughes, 2013	Utasai Bog	Oligotrophic bog	Japan	42.38	140.18	7	2954	N	tephra (4)	Y ²	Y
13	Kokfelt, 2010	Stordalen	Permafrost bog	Sweden	68.35	19.05	11	4717	Y	N	Y ²	Y
14	Lamentowicz, 2013	Stažki-F	Rich fen	Poland	54.43	18.09	8	1225	Y ¹	N	Y ⁵	Y
15	Large, 2009	Hongyuan HYLK1	Bog	China	32.77	102.52	14	10,827	Y	N	Y ²	Y
16	J. Loisel, unpubl.	KAM12-C1	Bog	Russia (Far-E)	54.9	156.6	13	11,914	Y	N	Y ³	Y
17	G. MacDonald, unpubl	N-1	Peat plateau	Russia (W Sib)	63.16	74.82	8	10,072	Y	N	Y ³	Y
18	MacDonald, unpubl	V-34	Open raised bog	Russia (W Sib)	61.47	79.46	8	8824	Y	N	Y ³	Y
19	MacDonald, unpubl	E-110	Peat plateau	Russia (W Sib)	66.47	76.99	6	9496	Y	N	Y ³	Y
20	MacDonald, unpubl	D-127	Peat plateau	Russia (W Sib)	64.31	70.29	6	10,034	Y	N	Y ³	Y
21	MacDonald, unpubl	SIB06	Pine-domin. bog	Russia (W Sib)	58.44	83.43	17	8680	Y	N	Y ³	Y
22	Mäkilä, 2007 ^a	Hanhijänkä	Palsa	Finland	68.4	23.55	7	9800	Y	N	Y ⁶	N
23	Mäkilä, 2007 ^a	Luovuoma (3 cores)	Fen	Finland	68.4	23.55	6	9800	Y	N	Y ⁶	N
24	Mäkilä, 2001 ^a	Ruosuo (P8)	Aapa	Finland	65.65	27.32	7	9500	Y	N	Y ²	Y
25	Mäkilä, 2001 ^a	Ruosuo (P20)	Aapa	Finland	65.65	27.32	9	9500	Y ¹	N	Y ²	Y
26	Mäkilä, 2001 ^a	Saarisuo (B800)	Fen	Finland	65.65	27.32	11	9600	Y	N	Y ²	Y
27	Mäkilä, 1997 ^a	Haukkasuo (3 cores)	Bog	Finland	60.82	26.95	13	9500	Y	N	Y ⁶	N
28	P. Mathijssen, unpubl	Lompolojänkä	Fen	Finland	68	24.22	10	9969	Y	N	Y ³	Y
29	Mathijssen, unpubl	Siikaneva	Bog	Finland	61.84	24.17	6	9622	Y	N	Y ⁴	Y
30	Mauquoy, 2002	Walton Moss 21	Raised bog	UK	54.98	-2.77	21	1120	N	N	Y ²	Y
31	Mauquoy, 2002	Walton Moss 20	Raised bog	UK	54.98	-2.77	23	1048	N	N	Y ²	Y
32	Mauquoy, 2002	Walton Moss 19	Raised bog	UK	54.98	-2.77	30	925	N	N	Y ²	Y
33	Mauquoy, 2002	Lille Vildmose	Raised bog	Denmark	56.83	10.25	19	609	N	N	Y ²	Y

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Abbreviated reference [*]	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
Oksanen, 2001	Rogovaya River 2	Peat plateau	Russia (E Eur.)	67.27	62.14	5	10,413	Y ¹	N	Y ²	Y
Oksanen, 2001	Rogovaya River 3	Peat plateau	Russia (E Eur.)	67.25	62.07	6	10,641	Y	N	Y ³	Y
Oksanen, 2003	Usinsk Mire 1	Peat plateau	Russia (E. Eur.)	57.42	65.67	6	13,236	Y	N	Y ²	Y
Ronkainen, unpubl	Seida	Peat plateau	Russia (E. Eur.)	67.05	62.92	6	8469	Y	N	Y ⁴	Y
Ruhland, 2000	Lena River Valley	Wet fen	Russia (E Sib)	69.38	125.13	6	8022	Y	N	Y ⁵	N
Tuittila, 2007	Lakkasuo (hummock)	Bog	Finland	61.78	24.3	12	6567	Y ¹	N	Y ⁵	Y
Tuittila, 2007	Lakkasuo (lawn)	Bog	Finland	61.78	24.3	7	6803	Y	N	Y ⁵	Y
Turunen, 2001 ^a	Salym-Gyugan Mire 3	Bog	Russia (W Sib)	60.17	72.83	6	10,500	Y	N	Y ⁶	N
Välianta, 2007	Kontolanrahka	Bog	Finland	60.78	22.78	40	4937	Y	¹³⁷ Cs	Y ²	Y
van der Linden, 2006	Saxnäs Mosse	Raised bog	Sweden	56.86	13.46	36	1068	N	N	Y ²	Y
van der Linden, 2007	Barschpfuhl	Kettle hole	Germany	53.05	13.83	32	134	N	N	Y ²	Y
van der Linden, 2008	Lappmyran	String & flark mire	Sweden	64.16	19.58	40	1712	N	N	Y ²	Y
van der Linden, 2008	Åkerlänna Römossa	Raised bog	Sweden	60.02	17.36	36	392	N	N	Y ²	Y
Y. Zhao, unpubl.	Altay	Sedge-dom rich fen	China	48.12	88.35	18	11,308	Y	N	Y ³	Y
Zhao, 2011	Zoige	Sedge-dom rich fen	China	33.45	102.63	7	9996	Y	N	Y ³	Y
Zhou, 2010	Hani Peat Bog	Bog	China	42.22	126.52	6	15,014	Y	N	Y ⁵	N
Anderson, 1998	Eilean Subhainn	Olig. topogen. bog	UK	57.69	-5.48	4	8700	Y	N	N	Y
Beilman, unpubl.	KAM12-C10	Fen	Russia (Far-E)	55.5	159.87	1	7500	Y	N	N	Y
Juutinen, 2013	Kiposuo III	Fen	Finland	69.18	27.28	3	9510	Y	pollen (1)	N	Y
Juutinen, 2013	Kiposuo IV	Fen	Finland	69.18	27.28	2	8574	Y ¹	N	N	Y
McCarroll, unpubl	Mossdale Moor 2	Blanket bog	UK	49.85	-7.46	3	1429	N	N	N	Y
Smith, 2004, 2012	N-2	Peat plateau	Russia (W Sib)	63.88	75.02	1	3600	Y	N	N	Y
Smith, 2004, 2012	S-4	Non-permafrost	Russia (W Sib)	61.55	72.71	1	6285	Y	N	N	Y
Smith, 2004, 2012	S-5	Non-permafrost	Russia (W Sib)	61.98	72.18	1	3885	Y	N	N	Y
Smith, 2004, 2012	S-6	Non-permafrost	Russia (W Sib)	61.62	73.98	1	11,120	Y	N	N	Y
Smith, 2004, 2012	S-7	Non-permafrost	Russia (W Sib)	61.49	74.32	1	8675	Y	N	N	Y

1
2
3
4

	Abbreviated reference ^a	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
5	Smith, 2004, 2012	S-8	Pine-domin. bog	Russia (W Sib)	61.75	73.39	1	9860	Y	N	N	Y
6	Smith, 2004, 2012	S-9	Non-permafrost	Russia (W Sib)	62.12	73.84	2	8725	Y	N	N	Y
7	Smith, 2004, 2012	N-10	Peat plateau	Russia (W Sib)	63.14	76.54	1	4720	Y	N	N	Y
8	Smith, 2004, 2012	N-11	Non-permafrost	Russia (W Sib)	62.66	76.77	1	5090	Y	N	N	Y
9	Smith, 2004, 2012	N-12	Peat plateau	Russia (W Sib)	63.50	76.82	1	10,080	Y	N	N	Y
10	Smith, 2004, 2012	N-13	Peat plateau	Russia (W Sib)	63.77	76.64	1	9465	Y	N	N	Y
11	Smith, 2004, 2012	N-14	Peat plateau	Russia (W Sib)	63.77	75.51	1	9035	Y	N	N	Y
12	Smith, 2004, 2012	N-15	Peat plateau	Russia (W Sib)	63.65	74.27	2	9630	Y	N	N	Y
13	Smith, 2004, 2012	N-16	Peat plateau	Russia (W Sib)	64.50	75.53	1	3540	Y	N	N	Y
14	Smith, 2004, 2012	N-17	Peat plateau	Russia (W Sib)	64.07	74.99	1	11,330	Y	N	N	Y
15	Smith, 2004, 2012	N-18	Peat plateau	Russia (W Sib)	62.85	75.22	1	1005	Y	N	N	Y
16	Smith, 2004, 2012	N-19	Peat plateau	Russia (W Sib)	62.96	74.26	1	8290	Y ¹	N	N	Y
17	Smith, 2004, 2012	N-19-1	Peat plateau	Russia (W Sib)	62.96	74.26	1	8675	Y	N	N	Y
18	Smith, 2004, 2012	S-20	Pine-domin. bog	Russia (W Sib)	62.55	71.72	1	3395	Y	N	N	Y
19	Smith, 2004, 2012	S-21	Pine-domin. bog	Russia (W Sib)	62.40	72.87	1	9905	Y	N	N	Y
20	Smith, 2004, 2012	S-22	Pine-domin. bog	Russia (W Sib)	60.84	71.26	2	7125	Y	N	N	Y
21	Smith, 2004, 2012	S-23	Pine-domin. bog	Russia (W Sib)	60.65	73.08	1	6665	Y	N	N	Y
22	Smith, 2004, 2012	S-24	Open raised bog	Russia (W Sib)	61.32	73.24	1	2305	Y	N	N	Y
23	Smith, 2004, 2012	S-25	Pine-domin. bog	Russia (W Sib)	62.25	74.78	1	9910	Y	N	N	Y
24	Smith, 2004, 2012	V-26	Open raised bog	Russia (W Sib)	61.03	76.47	2	9700	Y	N	N	Y
25	Smith, 2004, 2012	V-27	Open raised bog	Russia (W Sib)	61.32	76.73	1	4540	Y	N	N	Y
26	Smith, 2004, 2012	V-28	Open raised bog	Russia (W Sib)	61.81	77.50	1	7750	Y	N	N	Y
27	Smith, 2004, 2012	V-29	Open raised bog	Russia (W Sib)	61.23	75.31	1	9750	Y	N	N	Y
28	Smith, 2004, 2012	V-30	Open raised bog	Russia (W Sib)	61.74	75.20	1	5455	Y	N	N	Y
29	Smith, 2004, 2012	V-31	Open raised bog	Russia (W Sib)	62.37	75.79	1	5600	Y	N	N	Y
30	Smith, 2004, 2012	V-32	Open raised bog	Russia (W Sib)	62.36	77.48	1	2140	Y	N	N	Y
31												
32												
33												
34												
35												
36												
37												
38												
39												
40												
41												
42												
43												
44												
45												
46												
47												
48												
49												

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Abbreviated reference	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
Smith, 2004, 2012	V-33	Open raised bog	Russia (W Sib)	62.00	76.71	1	10,975	Y	N	N	Y
Smith, 2004, 2012	V-35	Open raised bog	Russia (W Sib)	60.80	77.62	1	10,350	Y	N	N	Y
Smith, 2004, 2012	V-36	Open raised bog	Russia (W Sib)	60.81	78.58	1	4400	Y	N	N	Y
Smith, 2004, 2012	V-37	Betula & Salix fen	Russia (W Sib)	61.25	74.73	1	2425	Y	N	N	Y
Smith, 2004, 2012	V-38	Open raised bog	Russia (W Sib)	60.80	74.54	2	7525	Y	N	N	Y
Smith, 2004, 2012	V-39	Open raised bog	Russia (W Sib)	61.09	79.38	2	10,925	Y	N	N	Y
Smith, 2004, 2012	V-40	Open raised bog	Russia (W Sib)	61.20	77.84	1	7850	Y	N	N	Y
Smith, 2004, 2012	E-101	Peat plateau	Russia (W Sib)	66.46	76.68	1	10,970	Y	N	N	Y
Smith, 2004, 2012	E-102	Peat plateau	Russia (W Sib)	66.04	76.59	1	8065	Y	N	N	Y
Smith, 2004, 2012	E-103	Peat plateau	Russia (W Sib)	66.74	76.48	1	10,395	Y	N	N	Y
Smith, 2004, 2012	E-104	Peat plateau	Russia (W Sib)	65.97	77.99	1	4240	Y	N	N	Y
Smith, 2004, 2012	E-105	Peat plateau	Russia (W Sib)	65.98	77.61	1	735	Y	N	N	Y
Smith, 2004, 2012	E-106	Peat plateau	Russia (W Sib)	66.00	77.35	1	9175	Y	N	N	Y
Smith, 2004, 2012	E-107	Peat plateau	Russia (W Sib)	66.01	75.86	1	6650	Y	N	N	Y
Smith, 2004, 2012	E-108	Peat plateau	Russia (W Sib)	65.86	75.29	1	10,685	Y	N	N	Y
Smith, 2004, 2012	E-111	Peat plateau	Russia (W Sib)	66.20	79.14	1	8630	Y	N	N	Y
Smith, 2004, 2012	E-112	Peat plateau	Russia (W Sib)	66.20	79.14	1	8765	Y	N	N	Y
Smith, 2004, 2012	E-113	Peat plateau	Russia (W Sib)	66.45	79.32	4	8305	Y	N	N	Y
Smith, 2004, 2012	E-114	Peat plateau	Russia (W Sib)	66.44	76.32	1	605	Y	N	N	Y
Smith, 2004, 2012	E-115	Peat plateau	Russia (W Sib)	67.81	75.43	2	9120	Y	N	N	Y
Smith, 2004, 2012	E-116	Peat plateau	Russia (W Sib)	67.46	76.42	1	3050	Y	N	N	Y
Smith, 2004, 2012	E-118	Peat plateau	Russia (W Sib)	66.60	77.41	1	2540	Y	N	N	Y
Smith, 2004, 2012	E-118M	Peat plateau	Russia (W Sib)	66.60	77.41	0		N	N	N	Y
Smith, 2004, 2012	E-119	Peat plateau	Russia (W Sib)	65.50	75.50	2	9750	Y	N	N	Y
Smith, 2004, 2012	E-120	Peat plateau	Russia (W Sib)	65.61	77.96	1	2585	Y	N	N	Y
Smith, 2004, 2012	E-120M	Peat plateau	Russia (W Sib)	65.61	77.96	0		N	N	N	Y

1
2
3
4

	Abbreviated reference*	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
5												
6												
7												
8	Smith, 2004, 2012	E-121	Peat plateau	Russia (W Sib)	65.87	78.81	1	2190	Y	N	N	Y
9	Smith, 2004, 2012	E-121M	Peat plateau	Russia (W Sib)	65.87	78.81	0		N	N	N	Y
10	Smith, 2004, 2012	D-122	Peat plateau	Russia (W Sib)	65.58	73.01	2	8495	Y	N	N	Y
11	Smith, 2004, 2012	D-123	Peat plateau	Russia (W Sib)	64.42	71.03	1	10,080	Y	N	N	Y
12	Smith, 2004, 2012	D-123M	Peat plateau	Russia (W Sib)	64.42	71.03	0		N	N	N	Y
13	Smith, 2004, 2012	D-124	Peat plateau	Russia (W Sib)	65.08	72.97	1	6475	Y	N	N	Y
14	Smith, 2004, 2012	D-124M	Peat plateau	Russia (W Sib)	65.08	72.97	0		N	N	N	Y
15	Smith, 2004, 2012	D-125	Peat plateau	Russia (W Sib)	64.52	72.16	1	9600	Y	N	N	Y
16	Smith, 2004, 2012	D-125M	Peat plateau	Russia (W Sib)	64.52	72.16	1	9735	Y	N	N	Y
17	Smith, 2004, 2012	D-126	Peat plateau	Russia (W Sib)	64.33	71.20	1	9140	Y	N	N	Y
18	Smith, 2004, 2012	D-126M	Peat plateau	Russia (W Sib)	64.33	71.20	0		N	N	N	Y
19	Smith, 2004, 2012	D-127M	Peat plateau	Russia (W Sib)	64.31	70.29	1	10,420	Y	N	N	Y
20	Smith, 2004, 2012	D-128	Peat plateau	Russia (W Sib)	65.55	72.46	1	9180	Y	N	N	Y
21	Smith, 2004, 2012	P-129	Peat plateau	Russia (W Sib)	66.61	73.75	1	9635	Y	N	N	Y
22	Smith, 2004, 2012	P-130	Peat plateau	Russia (W Sib)	66.87	74.53	1	8815	Y	N	N	Y
23	Smith, 2004, 2012	P-131	Peat plateau	Russia (W Sib)	66.17	73.99	2	9940	Y	N	N	Y
24	Smith, 2004, 2012	P-132	Peat plateau	Russia (W Sib)	66.50	73.95	1	10,065	Y	N	N	Y
25	Smith, 2004, 2012	P-133	Peat plateau	Russia (W Sib)	65.79	74.35	1	6515	Y	N	N	Y
26	Smith, 2004, 2012	G-134	Peat plateau	Russia (W Sib)	64.43	77.18	1	8285	Y	N	N	Y
27	Smith, 2004, 2012	G-135	Peat plateau	Russia (W Sib)	64.83	77.67	1	9450	Y	N	N	Y
28	Smith, 2004, 2012	G-136	Peat plateau	Russia (W Sib)	64.15	75.36	2	7820	Y	N	N	Y
29	Smith, 2004, 2012	G-136M	Peat plateau	Russia (W Sib)	64.15	75.36	1	6385	Y	N	N	Y
30	Smith, 2004, 2012	G-137	Peat plateau	Russia (W Sib)	63.75	75.77	4	9360	Y	N	N	Y
31	Smith, 2004, 2012	G-138	Peat plateau	Russia (W Sib)	64.52	76.67	1	9915	Y	N	N	Y
32	Smith, 2004, 2012	G-139	Peat plateau	Russia (W Sib)	64.89	76.73	1	6240	Y	N	N	Y
33	Smith, 2004, 2012	G-139M	Peat plateau	Russia (W Sib)	64.89	76.73	0		N	N	N	Y
34												
35												
36												
37												
38												
39												
40												
41												
42												
43												
44												
45												
46												
47												
48												
49												

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Abbreviated reference ^a	Site name	Peatland type	Country	Latitude	Longitude	¹⁴ C dates (number)	Oldest age (cal BP)	Basal age (cal BP)	Other dates	Carbon rate site	Peat properties site
Smith, 2004, 2012	G-140	Peat plateau	Russia (W Sib)	64.27	79.55	1	10,365	Y	N	N	Y
Smith, 2004, 2012	G-140M	Peat plateau	Russia (W Sib)	64.27	79.55	0		N	N	N	Y
Smith, 2004, 2012	G-141	Peat plateau	Russia (W Sib)	64.69	75.40	1	10,410	Y	N	N	Y
Smith, 2004, 2012	G-142	Peat plateau	Russia (W Sib)	64.09	78.60	0		N	N	N	Y
Smith, 2004, 2012	G-142M	Peat plateau	Russia (W Sib)	64.09	78.60	1	8675	Y	N	N	Y
Smith, 2004, 2012	SIB01	Pine-domin. bog	Russia (W Sib)	59.36	68.98	3	6970	Y	N	N	Y
Smith, 2004, 2012	SIB02	Pine-domin. bog	Russia (W Sib)	61.06	70.06	2	8500	Y	N	N	Y
Smith, 2004, 2012	SIB03	Pine-domin. bog	Russia (W Sib)	56.36	79.07	3	2770	Y	N	N	Y
Smith, 2004, 2012	SIB04	Pine-domin. bog	Russia (W Sib)	56.80	78.74	3	3770	Y	N	N	Y
Smith, 2004, 2012	SIB05	Pine-domin. bog	Russia (W Sib)	57.35	81.16	3	4240	Y	N	N	Y
Välranta, 2003	Ortino 1	Peat plateau	Russia (E. Eur.)	68	54	4	10,374	Y	N	N	Y
Välranta, 2003	Ortino 2	Peat plateau	Russia (E. Eur.)	68	54	3	8786	Y ¹	N	N	Y

^aA list of detailed references is presented below the table.

^aSite used in Yu et al.'s (2009) synthesis.

¹Basal age not considered in the peatland inception age database because older cores were collected from the same site.

²Measured bulk density was multiplied by measured C content (elemental analyzer) for each layer to estimate C bulk density (g C cm⁻³).

³Measured ash-free bulk density was multiplied by inferred C content (ash-free bulk density x 49%) for each layer to estimate C bulk density.

⁴Measured bulk density was multiplied by assumed C content (47%) for each layer to estimate C bulk density.

⁵Assumed time-dependent bulk density was multiplied by assumed C content (47%) for each dated interval to estimate peat-C density.

⁶Peat-C accumulation rates directly obtained from published figures and tables.

References:

- Anderson, D.E. 1998. A reconstruction of Holocene climatic changes from peat bogs in north-west Scotland. *Boreas*, 27, 208-224.
- Andersson, S., Schoning, K. 2010. Surface wetness and mire development during the late Holocene in central Sweden. *Boreas*, 39, 749-760.
- Barber, K.E., Chambers, F.M., Maddy, D. 2003. Holocene palaeoclimates from peat stratigraphy: macrofossil proxy climate records from three oceanic raised bogs in England and Ireland. *Quaternary Science Reviews*, 22, 521-539.
- Beaulieu-Audy, V., Garneau, M., Richard, P.J.H., Asnong, H. 2009. Holocene palaeoecological reconstruction of three boreal peatlands in the La Grande Rivière region, Québec, Canada. *The Holocene*, 19(3), 459-476.
- Belyea, L.R., Warner, B.G. 1996. Temporal scale and the accumulation of peat in a *Sphagnum* bog. *Canadian Journal of Botany*, 74, 366-377.
- Bender, M.M., Bryson, R.A., Baerreis, D.A. 1969. University of Wisconsin radiocarbon dates VI. *Radiocarbon*, 11(1), 228-235.
- Booth, R.K., Jackson, S.T., Gray, C.E.D. 2004. Paleoecology and high-resolution paleohydrology of a kettle peatland in upper Michigan. *Quaternary Research*, 61, 1-13.
- Borren, W., Bleuten, W., Lapshina, E.D. 2004. Holocene peat and carbon accumulation rates in the southern taiga of western Siberia. *Quaternary Research*, 61, 42-51.
- Bunbury, J., Finkelstein, S.A., Bollman, J. 2012. Holocene hydro-climatic change and effects on carbon accumulation inferred from a peat bog in the Attawapiskat River watershed, Hudson Bay Lowlands, Canada. *Quaternary Research*, 78, 275-284.
- Camill, P., Barry, A., Williams, E., Andreassi, C., Limmer, J., Solick, D. 2009. Climate-vegetation-fire interactions and their impact on long-term carbon dynamics in a boreal peatland landscape in northern Manitoba, Canada. *Journal of Geophysical Research*, 114, G04017, doi:10.1029/2009JG001071.
- Charman, D.J. 1994. Patterned fen development in northern Scotland: developing a hypothesis from palaeoecological data. *Journal of Quaternary Science*, 9(3), 285-297.
- Charman, D.J., Aravena, R., Warner, B.G. 1995. Carbon dynamics in a forested peatland in North-Eastern Ontario, Canada. *Journal of Ecology*, 82(1), 55-62.
- De Vleeschouwer, F., Piotrowska, N., Sikorski, J., Pawlyta, J., Cheburkin, A., Le Roux, G., Lamentowicz, M., Fagel, N., Mauquoy, D. 2009. Multiproxy evidence of 'Little Ice

Age' palaeoenvironmental changes in a peat bog from northern Poland. *The Holocene*, 19(4), 625-637.

De Vleeschouwer, F., Pazdur, A., Luthers, C., Streel, M., Mauquoy, D., Wastiaux, C., Le Roux, G., Moschen, R., Blaauw, M., Pawlyta, J., Sikorski, J., Piotrowska, N. 2012. A millennial record of environmental change in peat deposits from the Misten bog (East Belgium). *Quaternary International*, 268, 44-57.

Elliott, S.M., Roe, H.M., Patterson, T. 2011. Testate amoebae as indicators of hydroseral change: An 8500 years record from Mer Bleue Bog, eastern Ontario, Canada. *Quaternary International*, 268, 128-144.

Gałka, M., Miotk-Szpiganowicz, G., Goslar, T., Ješko, M., van der Knaap, W., Lamentowicz, M. 2013a. Paleohydrology, fires and vegetation succession in the southern Blatic during the last 7500 years reconstructed from a raised bog based on multi-proxy data. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 370, 209-221.

Gałka, M., Tobolski, K., Górski, A., Milecka, K., Fiałkiewicz-Kozieł, B., Lamentowicz, M. 2013b. Disentangling the drivers of a Baltic bog development during the Little Ice Age in northern Poland. *Quaternary International*, doi.org/10.1016/j.quaint.2013.02.026.

Glaser, P.H., Hansen, B.C.S., Siegel, D.I., Reeve, A.S., Morin, P.J. 2004. Rates, pathways and drivers for peatland development in the Hudson Bay Lowlands, northern Ontario, Canada. *Journal of Ecology*, 92, 1036-1053.

Glebov, F.Z., Karpenko, L.V., Dashkovskaya, I.S. 2002. Climatic changes, successions of peatlands and zonal vegetation, and peat accumulation dynamics in the Holocene (the West-Siberia peat profile 'Vodorasdel'). *Climatic Change*, 55, 175-181.

Gorham, E., Janssens, J.A., Glaser, P.H. 2003. Rates of peat accumulation during the postglacial period in 32 sites from Alaska to Newfoundland, with special emphasis on northern Minnesota. *Canadian Journal of Botany*, 81, 429-438.

Hendon, D., Charman, D.J., Kent, M. 2001. Palaeohydrological records derived from testate amoebae analysis from peatlands in northern England: within-site variability, between-site comparability and palaeoclimatic implications. *The Holocene*, 11(2), 127-148.

Hu, F.S., Davis, R.B. 1994. Postglacial development of a Maine bog and paleoenvironmental implications. *Canadian Journal of Botany*, 73, 638-649.

Hughes, P.D.M., Blundell, A., Charman, D.J., Bartlett, S., Daniell, J.R.G., Wojatschke, A., Chambers, F.M. 2006. An 8500 cal. year multi-proxy climate record from a bog in eastern Newfoundland: contributions of meltwater discharge and solar forcing. *Quaternary Science Reviews*, 25, 1208-1227.

Hughes, P.D.M., Mallon, G., Brown, A., Esssex, H.J., Stanford, J.D., Hotes, S. 2013. The impact of high tephra loading on late-Holocene carbon accumulation and vegetation succession in peatland communities. *Quaternary Science Reviews*, 67, 160-175.

Hunt, S., Yu, Z., Jones, M. 2013. Lateglacial and Holocene climate, disturbance and permafrost peatland dynamics on the Seward Peninsula, western Alaska. *Quaternary Science Reviews*, 63, 42-58.

Jones, M.C., Yu, Z. 2010. Rapid deglacial and early Holocene expansion of peatlands in Alaska. *Proceedings of the National Academy of Sciences*, 107(15), 7347-7352.

Juutinen, S., Väiliranta, M., Kuutti, V., Laine, A.M., Virtanen, T., Seppä, H., Weckström, Tuittila, E.-S. 2013. Short-term and long-term carbon dynamics in a northern peatland-stream-lake continuum: A catchment approach. *Journal of Geophysical Research*, 118, 1-13, doi:10.1002/JGRG20028.

Klein, E.S., Booth, R.K., Yu, Z., Mark, B.G., Stansell, N.D. 2013. Hydrology-mediated differential response of carbon accumulation to late Holocene climate change at two peatlands in Southcentral Alaska. *Quaternary Science Reviews*, 64, 61-75.

Kokfelt, U., Reuss, N., Struyf, E., Sonesson, M., Rundgren, M., Skog, G., Rosén, P., Hammarlund, D. 2010. Wetland development, permafrost history and nutrient cycling inferred from late Holocene peat and lake sediment records in subarctic Sweden. *Journal of Paleolimnology*, 44, 327-342.

Kuhry, P., Vitt, D.H. 1996. Fossil carbon/nitrogen ratios as a measure of peat decomposition, *Ecology*, 77(1), 271-275.

Lamentowicz, M., Gałka, M., Milecka, K., Tobolski, K., Lamentowicz, L., Fiałkiewicz-Kozieł, B., Blaauw, M. 2013. A 1300-year multi-proxy, high-resolution record from a rich fen in northern Poland: reconstructing hydrology, land use and climate change. *Journal of Quaternary Science*, 28(6): 582-594.

Lamarre, A., Garneau, M., Asnong, H. 2012. Holocene paleohydrological reconstruction and carbon accumulation of a permafrost peatland using testate amoeba and macrofossil analyses, Kuujjuarapik, subarctic Québec, Canada. *Review of Palaeobotany and Palynology*, 186, 131-141.

Large, D.J., Spiro, B., Ferrat, M., Shopland, M., Kylander, M., Gallagher, K., Li, X., Shen, C., Possnert, G., Zhang, G., Darling, W.G., Weiss, D. 2009. The influence of climate, hydrology and permafrost on Holocene peat accumulation at 3500 m on the eastern Qinghai-Tibetan Plateau. *Quaternary Science Reviews*, 3303-3314.

Lavoie, M., Richard, P.J.H. 2000. The role of climate on the developmental history of Frontenac Peatland, southern Quebec. *Canadian Journal of Botany*, 78, 668-684.

Lavoie, M., Richard, P.J.H. 2000. Paléoécologie de la tourbière du lac Malbaie, dans le massif des Laurentides (Québec): évaluation du rôle du climat sur l'accumulation de la tourbe. *Géographie Physique et Quaternaire*, 54(2), 169-185.

Lavoie, M., Pellerin, S., Larocque, M. 2013. Examining the role of allogenuous and autogenous factors in the long-term dynamics of a temperate headwater peatland (southern Québec, Canada). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 386, 336-348.

Loisel, J., Garneau, M. 2010. Late Holocene paleoecohydrology and carbon accumulation estimates from two boreal peat bogs in eastern Canada: Potential and limits of multi-proxy archives. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 291, 493-533.

Loisel, J., Yu, Z. 2013. Recent acceleration of carbon accumulation in a boreal peatland, south central Alaska. *Journal of Geophysical Research*, 118, 1-13, doi:10.1029/2012JG001978.

MacDonald, G.M. 1983. Holocene vegetation history of the Upper Natla River Area, Northwest Territories, Canada. *Arctic and Alpine Research*, 15(2), 169-180.

Magnan, G., Lavoie, M., Payette, S. 2012. Impact of fire on long-term vegetation dynamics of ombrotrophic peatlands in northwestern Québec, Canada. *Quaternary Research*, 77, 110-121.

Mäkilä, M. 1997. Holocene lateral expansion, peat growth and carbon accumulation on Haukkasuo, a raised bog in southeastern Finland, *Boreas*, 26 (1), 1-14, doi:10.1111/j.1502-3885.1997.tb00647.x.

Mäkilä, M., Saarnisto, M., Kankainen, T. 2001. Aapa mires as a carbon sink and source during the Holocene. *Journal of Ecology*, 89(4), 589-599.

Mäkilä, M., Moisanen, M. 2007. Holocene lateral expansion and carbon accumulation of Luovuoma, a northern fen in Finnish Lapland. *Boreas*, 36(2), 198-210.

Mauquoy, D., Engelkes, T., Groot, M.H.M., Markesteijn, F., Oudejans, M.G., van der Plicht, J., van Geel, B. 2002. High-resolution records of late-Holocene climate change and carbon accumulation in two north-west European ombrotrophic peat bogs. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 186, 275-310.

Muller, S.D., Richard, P.J.H., Larouche, A.C. 2003. Holocene development of a peatland (southern Québec): a spatio-temporal reconstruction based on pachymetry, sedimentology, microfossils and macrofossils. *The Holocene*, 13(5), 649-664.

O'Donnell, J.A., Jorgenson, M.T., Harden, J.W., McGuire, A.D., Kanevskiy, M.Z., Wickland, K.P. 2012. The effects of permafrost thaw on soil hydrologic, thermal, and carbon dynamics in an Alaskan peatland. *Ecosystems*, 15, 213-229.

O'Reilly, B.C. 2011. Paleoecological and carbon accumulation dynamics of a peatland in the Hudson Bay Lowlands, Northern Ontario, from the mid-Holocene to present. M.Sc thesis, University of Toronto, Canada.

Oksanen, P.O., Kuhry, P., Alekseeva, R.N. 2001. Holocene development of the Rogovaya River peat plateau, European Russian Arctic. *The Holocene*, 11(1): 25-40.

Oksanen, P.O., Kuhry, P., Alekseeva, R.N. 2003. Holocene development and permafrost history of the Usinsk Mire, Northeast European Russia. *Géographie Physique et Quaternaire*, 57(2-3), 169-187.

Robinson, S.D. 2000. Carbon accumulation in discontinuously frozen peatlands, Southwestern Northwest Territories, Canada. Ph.D. thesis, McGill University, Canada.

Robinson, S.D. 2006. Carbon accumulation in peatlands, southwestern Northwest Territories, Canada. *Canadian Journal of Soil Science*, 86, 305-319.

Ruhland, K., Smol, J.P., Jasinski, J.P.P., Warner, B.G. 2000. Response of diatoms and other siliceous indicators to the developmental history of a peatland in the Tiksi Forest, Siberia, Russia. *Arctic, Antarctic, and Alpine Research*, 32(2), 167-178.

Sannel, A.B.K., Kuhry, P. 2009. Holocene peat growth and decay dynamics in sub-arctic peat plateaus, west-central Canada. *Boreas*, 38: 13-24.

Smith, L.C., Beilman, D.W., Kremenetski, K.V., Sheng, Y., MacDonald, G.M., Lammers, R.B., Shiklomanov, A.I., Lapshina, E.D. 2012. Influence of permafrost on water storage in West Siberian peatlands revealed from a new database of soil properties. *Permafrost and Periglacial Processes*, 23, 69-79.

Smith, L.C., MacDonald, G.M., Velichko, A.A., Beilman, D.W., Borisova, O.K., Frey, K.E., Kremenetski, K.V., Sheng, Y. 2004. Siberian peatlands a net carbon sink and global methane source since the early Holocene. *Science*, 303(5656), 353-356.

Tarnocai, C. 2010. Carbon sequestration dynamics and climate change in Subarctic and Low Arctic organic cryosols in Canada. *Proceedings of the 19th World Congress of Soil Science*, p. 5-8.

Tuittila, E.-S., Välranta, M., Laine, J., Korhola, A. 2007. Quantifying patterns and controls of mire vegetation succession in a southern boreal bog in Finland using partial ordinations. *Journal of Vegetation Science*, 18, 891-902.

Turunen, J., Tahvanainen, T., Tolonen, K. 2001. Carbon accumulation in West Siberian mires, Russia. *Global Biogeochemical Cycles*, 15(2), 285-296.

Turunen, C., Turunen, J. 2003. Development history and carbon accumulation of a slope bog in oceanic British Columbia, Canada. *The Holocene*, 13(2), 225-238.

Väliranta, M., Kaakinen, A., Kuhry, P. 2003. Holocene climate and landscape evolution East of the Pechora delta, East-European Russian Arctic. *Quaternary Research*, 59, 335-344.

Väliranta, M., Korhola, A., Seppä, H., Tuittila, E.-S., Sarmaja-Korjonen, K., Laine, J., Alm, J. 2007. High-resolution reconstruction of wetness dynamics in a southern boreal raised bog, Finland, during the late Holocene: a quantitative approach. *The Holocene*, 17(8), 1093-1107.

van der Linden, M. 2007. Effects of climate change and human impact on late-Holocene species composition and carbon accumulation in bog ecosystems. Ph.D. thesis, Universiteit van Amsterdam, The Netherlands.

van der Linden, M., van Geel, B. 2006. Late-Holocene climate change and human impact recorded in a South Swedish ombrotrophic peat bog. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 240(3-4), 649-667.

van der Linden, M., Vickery, E., Charman, D.J., Broekens, P., van Geel, B. 2008. Vegetation history and human impact during the last 300 years recorded in a German peat deposit. *Review of Palaeobotany and Palynology*, 152, 158-175.

van Bellen, S., Dallaire, P.-L., Garneau, M., Bergeron, Y. 2011. Quantifying spatial and temporal Holocene carbon accumulation in ombrotrophic peatlands of the Eastmain region, Quebec, Canada, *Global Biogeochemical Cycles*, 25, GB2016, doi:10.1029/2010GB003877.

Yu, Z., Campbell, I.D., Campbell, C., Vitt, D.H., Bond, G.C., Apps, M.J. 2003. Carbon sequestration in western Canadian peat highly sensitive to Holocene wet-dry climate cycles at millennial timescales. *The Holocene*, 13(6), 801-808.

Yu, Z., Beilman, D.W., Jones, M.C. 2009. Sensitivity of northern peatland carbon dynamics to Holocene climate change. In: Baird, A., Belyea, L., Comas, X., Reeve, A., Slater, L. (Eds.), *Northern peatlands and carbon cycling*, American Geophysical Union Monograph Series, Washington D.C., USA, pp. 55-69.

Yu, Z. 2006. Holocene carbon accumulation of fen peatlands in boreal Western Canada: A complex ecosystem response to climate variation and disturbance. *Ecosystems*, 9(8), 1278-1288.

Zhao, Y., Yu, Z., Zhao, W. 2011. Holocene vegetation and climate histories in the eastern Tibetan Plateau: controls by insolation-driven temperature or monsoon-derived precipitation changes? *Quaternary Science Reviews*, 30, 1173-1184.

Zhou, W., Zheng, Y., Meyers, P.A., Jull, A.J.T., Xie, S. 2010. Postglacial climate-change record in biomarker lipid compositions of the Hani peat sequence, Northeastern China. *Earth and Planetary Science Letters*, 294, 37-46.

For Peer Review