



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under All Rights Reserved license:

**Chambers, Frank M ORCID logoORCID: <https://orcid.org/0000-0002-0998-2093> (2016) The 'Little Ice Age': the first virtual issue of The Holocene. *Holocene*, 26 (3). pp. 335-337.
doi:10.1177/0959683615593688**

Official URL: <http://dx.doi.org/10.1177/0959683615593688>

DOI: <http://dx.doi.org/10.1177/0959683615593688>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/2647>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

The Little Ice Age

Prof. Frank M. Chambers

Centre for Environmental Change and Quaternary Research, School of Natural and Social Sciences,
University of Gloucestershire, Francis Close Hall, Swindon Rd, Cheltenham, GL50 4AZ, UK
Tel +44 1242 714677

Introduction

The so-called Little Ice Age of the 15th to 19th Centuries is a fascinating period of time, for many reasons. Extensive reading of the literature on the topic can reveal the following: (1) in many (but not all) proxy-climate reconstructions, it is shown as having a fast and strong onset (O'Brien et al., 1995), exceeded in the Holocene perhaps only by the 8.2ka event (Mayewski et al., 2004); (2) it includes evidence for glacier re-advance—in northern Europe, particularly, to positions not otherwise (or seldom) reached within the Mid–Late Holocene (Matthews and Shakesby, 1984; McCarroll, 1991); (3) it follows the Medieval Climate Anomaly and precedes the period of recent 'Global Warming', and therefore it post-dates the Medieval Solar Maximum, encompasses up to three solar minima (Spörer, Maunder and Dalton) (Grove 1988), and precedes the 'Contemporary' (viz. Late-20th Century) Solar Maximum (Hoyt and Schatten, 1997; Pan and Yau, 2002); (4) there are multiple hypotheses as to the cause of its onset (cf. Miller et al., 2012), although it is widely considered that reduced solar activity is the cause of at least its most intense phases (cf. Mauquoy et al., 2002); (5) there are differing views as to the magnitude of the depression of global temperature (e.g. IPCC, 1995, Fig. 3.20; Mann, 2002; Soon and Baliunas, 2003; IPCC, 2013; Figs. 5.7, 5.8); indeed (6) comparison of individual reconstructions of Northern Hemisphere temperatures with the Intergovernmental Panel on Climate Change's earlier somewhat muted successive summary curves shows considerable difference (compare IPCC, 2001: Figs 2.20, 2.21; IPCC, 2007: Fig. 6.10, also Box 6.4, Fig 1); (7) it had previously been thought that its expression was *not* influenced by human activity, whereas the 'plague' claim in the 'early anthropogenic' hypothesis of Ruddiman (2003) implies otherwise; (8) it largely precedes what some have viewed and attempted to define as a new epoch, representing evidence of widespread human influence on global systems: the so-called Anthropocene (Crutzen and Stoermer, 2000; Zalasiewicz et al., 2015)); (9) its effects upon some human societies appear to have been profound in particular regions of the world, notably in Greenland (Ribeiro et al., 2012), Norway (Lamb, 1995) and the Alps (Le Roy Ladurie, 1971); and yet (11) its very existence as a coherent, globally climatically defined period has been questioned (compare Mann et al., 1999 with Goosse et al. 2005); although (12) recent work implies an in-phase relationship between the Southern and Northern Hemispheres (Simms et al., 2012; Chambers et al., 2014).

The Little Ice Age is a period for which, at the start, documentary and observational information is relatively sparse and localised, but by the close there is increasing availability of documentary, observational and direct instrumental meteorological records from many parts (though not all) of the globe.

From a scientific and cultural viewpoint, the Little Ice Age is of particular interest because, temporally, it sits between the so-called Medieval Climate Anomaly and the contemporary 'Anthropocene'; it includes the commencement both of continuous instrumental meteorological records and of reliable scientific observation and recording of sunspots; and (in its later part) encompasses the initial period of industrialization in parts of the northern hemisphere, with a concomitant commencement of a sustained increase in emissions of carbon dioxide and methane to atmosphere.

Selected papers

It is entirely appropriate that *The Holocene* journal, which focuses on recent environmental change, has published a large number of papers that either focus on or refer specifically to the Little Ice Age. In this virtual issue, a selection has been made that demonstrates geographical spread, a diversity of proxy-climate archives, and a range of techniques that can be used to extract a climate signal from these archives.

This selection of papers includes proxy-climate data from five continents: North America, from Alaska (Wiles et al., 1999), Canada (Johnston et al., 2010); South America, from Chile (Araneda et al. 2009), Argentina (Chambers et al., 2014); Europe, from Britain (Harrison et al., 2014) and Spain (Garcia-Ruiz et al., 2014); Asia, from China (Chen et al., 2005; Liu et al., 2011); Australasia, from New Zealand (Winkler, 2004); and the ocean, from the Alboran Sea (Nieto-Moreno et al., 2013)

It includes proxy-climate data from a range of archives: aeolian sand (Liu et al., 2011); documentary sources (Araneda et al., 2009); lake (Chen et al., 2005; Johnston et al., 2010); marine records (Nieto-Moreno et al., 2013); mire (Chambers et al., 2014); tree-rings (Wiles et al., 1999). It deals with LIA temperatures (Nesje and Dahl, 2003); hydrology (Nesje and Dahl, 2003; Johnston et al., 2010; Liu et al., 2011; Nieto-Moreno et al., 2013; Chambers et al., 2014); glacier activity (Wiles et al., 1999; Winkler, 2004; Araneda et al. 2009; Garcia-Ruiz et al., 2014; Harrison et al., 2014); and with inter-hemispheric comparison (Chambers et al., 2014).

Proxy-climate evidence is provided from analysis of plant macrofossils and peat humification (Chambers et al., 2014), dendroclimatology (Wiles et al., 1999), lake-sediment chemistry (Chen et al., 2005) and marine sediments (Nieto-Moreno et al., 2013); lake-level variation is derived from palaeolimnology and geophysical analyses (Johnston et al., 2010); glacier limits are reconstructed from geomorphology (Garcia-Ruiz et al., 2014), tree-ring data (Wiles et al., 1999), lichenometry (Winkler, 2004) and inferred from documentary records (Araneda et al. 2009) and glacier modelling (Harrison et al., 2014).

Summaries of the individual papers

In the most recently published of the selections, Chambers et al. (2014) compare proxy-climate records from a mire in Tierra del Fuego with those derived using identical methods in continental north-central Europe, and find that the hydrological response of the South American mire in the most extreme phases of the Little Ice Age is similar to that at 2800 cal. BP when the mire became unusually dry—an opposite response to that recorded from Europe. The timing of the dry phases in South America appears to match that of the most extreme phases of the LIA in Europe, and is

attributed to equatorward movement of moisture-bearing winds, possibly caused by reduced solar activity.

Garcia-Ruiz et al. (2014) investigated Holocene and 'Little Ice Age' glacial activity in the Marboré Cirque in the Central Spanish Pyrenees. They detected two separate glacial pulses within the LIA, the first probably being in the Maunder Minimum (late 17th or early 18th century), whereas the second took place between AD 1790 and 1830 (close to the Dalton Minimum); these pulses followed melting during the Medieval Climate Anomaly, and the two were separated by a glacial retreat.

Harrison et al. (2014) used a glacier-climate model, fuelled by data from local weather stations, to argue that the *last* glacier ice in Cairngorm, Scotland was in the LIA, and not over ten thousand years earlier in the Younger Dryas, as hitherto widely assumed. Evidence from boulder moraines is adduced in support of the contention. It is argued that the last glacier ice that existed elsewhere in upland Britain may also relate to the LIA.

Nieto-Moreno et al. (2013) analysed two deep-sea marine cores from the Mediterranean to investigate the time period from the Medieval Climate Anomaly, through the LIA and up to the late 20th Century. While inferred dry periods characterised the MCA and late 20th century, the LIA presented as a more humid phase, but "developed as a sequence of successive short and abrupt dry-humid phase alternation" (Nieto-Moreno et al., 2013: 1227).

Liu et al. (2011) analysed carbon isotopes from plant leaves in a 10.5 m aeolian section in northwestern China, finding large negative isotope excursions during the LIA, implying a wetter climate. This is interpreted as indicating regional hydrological changes associated with "possible changes in the trajectory or strength of the westerlies and/or the orographic effect in this region" (Liu et al., 2011: 409).

Johnston et al. (2010) examined a barrier-beach complex using multi-proxy palaeolimnological analyses combined with geophysical examination using ground-penetrating radar to investigate the former lake levels of Lake Athabasca, Canada. Interpretation of the data suggested that the lake level was, on average, some 2.3 m higher than present during the LIA, implying that the surrounding landscape of the Peace-Athabasca delta had, until recently, been flooded frequently.

Araneda et al. (2009) used documentary sources, including written records, maps, photographs and iconography to reconstruct the limits of the Cipreses Glacier, Chile. The authors infer that the last advance of the glacier in the LIA was c. AD 1842, and was in retreat from AD 1858 and subsequently. These data are compared with evidence from the San Rafael glacier, which reached its most recent maximum extent between AD 1857 and 1875. The 30-year discrepancy between the responses of the two glaciers is attributed to temperature and precipitation changes thought mainly associated with fluctuations in the Westerlies.

Chen et al. (2005) used principal components analysis on 21 elements in their examination of the sediment chemistry of Lake Erhai, southwest China, and found three controlling factors: (1) physical erosion in the catchment; (2) autochthonous precipitation of calcite; (3) early diagenesis in sediment. The LIA was characterised by low factor (3) and high (1), implying a cool-wet climate from AD 1550 to 1890, which the authors linked to the timing of the LIA in Europe; this contrasted with high factor

(3) and low (1) around Lake Erhai from AD 1340–1550 and AD 1890–1950, implying warm-dry episodes.

Winkler (2004) used lichenometric dating of moraines of four glaciers in the Mt Cook National Park, New Zealand, to detect the maximum extent of glaciers in the LIA, which was revealed as c. AD 1725–1740. Subsequent retreat of the four glaciers has been followed by readvances in the late 19th and early 20th centuries, and one of the glaciers (Tasman) has since reached its 18th century LIA maximum.

Nesje and Dahl (2003) challenged the simplistic view that the LIA was primarily about temperature changes by presenting data from southern Norway to show that rapid glacier advance there in the early 18th century was mainly a result of increased winter precipitation in mild, wet winters and not just lower summer temperatures. They compared LIA glacier fluctuations in southern Norway with those of the European Alps, and suggested that asynchronous 'Little Ice Age' maxima between the two regions may be related to trends in the North Atlantic Oscillation dipole pattern.

Wiles et al. (1999) conducted tree-ring studies on 13 glacier forefields in western Prince Edward Sound, Alaska. Cross-dated sequences from eight sites indicated synchronous glacial advances (on decadal timescales) in the early LIA of late 12th to 13th centuries), and again in the mid-LIA of the 17th to early 18th centuries, while 9 sites suggested a further advance in the late 19th century. The data compared well with studies of other glaciers in the region, allowing the structure of LIA glacier fluctuations to be discerned.

References

Araneda A, Torrejón F, Aguayo N et al. (2009) Historical records of Cipreses glacier (34°S): combining documentary-inferred 'Little Ice Age' evidence from Southern and Central Chile *The Holocene* 19: 1173–1183.

Chambers FM, Brain SA, Mauquoy D et al. (2014) The 'Little Ice Age' in the Southern Hemisphere in the context of the last 3000 years: peat-based proxy-climate data from Tierra del Fuego. *The Holocene* 24: 1649–1656.

Chen J, Wan G, Zhang DD et al. (2005) The 'Little Ice Age' recorded by sediment chemistry in Lake Erhai, southwest China. *The Holocene* 15: 925–931.

Crutzen P and Stoermer EF (2000) The 'Anthropocene'. *Global Change Newsletter* 41: 17–18.

Garcia-Ruiz JM, Palacios D, de Andrés N et al. (2014) Holocene and 'Little Ice Age' glacial activity in the Marboré Cirque, Monte Perdido Massif, Central Spanish Pyrenees. *The Holocene* 24: 1439–1452.

Goosse H, Renssen H, Timmermann A et al. (2005) Internal and forced climate variability during the last millennium: a model-data comparison using ensemble simulations. *Quaternary Science Reviews* 24: 1345–1360.

Grove JM (1988) *The Little Ice Age*. London: Methuen.

Harrison S, Rowan AG, Glasser NF et al. (2014) Little Ice Age glaciers in Britain: Glacier–climate modelling in the Cairngorm Mountains. *The Holocene* 24: 135–140.

Hoyt DV and Schatten KH (1997) *The Role of the Sun in Climate Change*. Oxford: Oxford University Press.

IPCC (1995) Climate Change 1995: The Science of Climate Change. *Contribution of WG I to the Second Assessment Report of the Intergovernmental Panel on Climate Change*. [Houghton JT, Meira Filho LG, Callander BA et al. (Eds)] Cambridge: Cambridge University Press.

IPCC (2001) *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, 2001*. [Houghton JT, Ding Y, Griggs DJ et al. (Eds)] Cambridge: Cambridge University Press.

IPCC (2007) Climate Change 2007: *Climate Change Working Group 1: the Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, 2007*. [Solomon S, Qin D, Manning, M. et al. (eds.)] Cambridge: Cambridge University Press.

IPCC (2013) Climate Change 2013: *Climate Change Working Group 1: the Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. [Stocker TF, Qin D, Plattner G-K et al. (Eds)] Cambridge: Cambridge University Press.

Johnston JW, Köster D, Wolfe BB et al. (2010) Quantifying Lake Athabasca (Canada) water level during the ‘Little Ice Age’ highstand from palaeolimnological and geophysical analyses of a transgressive barrier-beach complex *The Holocene* 20: 801–811.

Lamb HH (1995) *Climate, History and the Modern World, 2nd ed*. London: Routledge.

Le Roy Ladurie E (1971) *Histoire du climat depuis l’an mil*. Paris: Flammarion, 1967. Translated by Barbara Bray as *Times of Feast, Times of Famine: A History of Climate since the year 1000*. Revised and updated. Garden City: Doubleday.

Liu W, Liu Z, An Z et al. (2011) Wet climate during the ‘Little Ice Age’ in the arid Tarim Basin, northwestern China. *The Holocene* 21: 409–416.

Mann ME (2002) Little Ice Age. In: MacCracken, MC and Perry JS (eds) *Encyclopedia of Global Environmental Change*. Chichester: Wiley.

Mann ME, Bradley RS and Hughes MK (1999) Northern hemisphere temperatures during the past millennium: inferences, uncertainties, and limitations. *Geophysical Research Letters* 26: 759–762.

Matthews JA and Shakesby RA (1984) The status of the Little Ice Age in southern Norway: relative-age dating of Neoglacial moraines with Schmidt hammer and lichenometry. *Boreas* 13: 333–346.

Mauquoy D, Van Geel B, Blaauw M et al. (2002) Evidence from North-West European bogs shows ‘Little Ice Age’ climatic changes driven by variations in solar activity. *The Holocene* 12: 1–6.

Mayewski PA, Rohling EE, Stager JC et al. (2004) Holocene climate variability. *Quaternary Research* 62: 243–255.

McCarroll D (1991) The age and origin of Neoglacial moraines in Jotenheimen, southern Norway: new evidence from weathering-based data. *Boreas* 20: 283–295.

Miller GH, Geirsdóttir A, Zhong Y et al. (2012) Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks. *Geophysical Research Letters* 39: L02708.

Nesje A (2009) Latest Pleistocene and Holocene alpine glacier fluctuations in Scandinavia. *Quaternary Science Reviews* 28: 2119–2136.

Nesje, A and Dahl SO (2003) The ‘Little Ice Age’ – only temperature? *The Holocene* 13: 139–145.

Nieto-Moreno V, Martinez-Ruiz F, Giralt S et al. (2013) Climate imprints during the ‘Medieval Climate Anomaly’ and the ‘Little Ice Age’ in marine records from the Alboran Sea basin. *The Holocene* 23: 1227–1237.

O'Brien SR, Mayewski PA, Meeker LD et al. (1995) Complexity of Holocene climate as reconstructed from a Greenland ice core. *Science* 270: 1962–1964.

Pan KD and Yau KK (2002) Ancient observations link changes in sun's brightness and earth's climate. *EOS, Transactions, American Geophysical Union* 83: 481, 489–490.

Ribeiro S, Moros M, Ellegaard M et al. (2012) Climate variability in West Greenland during the past 1500 years: evidence from a high-resolution marine palynological record from Disko Bay. *Boreas* 41: 68–83.

Ruddiman WF (2003) The anthropogenic greenhouse gas era began thousands of years ago. *Climate Change* 61: 261–293.

Simms, AR, Ivins ER, DeWitt R et al. (2012) Timing of the most recent Neoglacial advance and retreat in the South Shetland Islands, Antarctic Peninsula: insights from raised beaches and Holocene uplift. *Quaternary Science Reviews* 47: 41–55.

Soon W and Baliunas S (2003) Proxy climatic and environmental changes of the past 1000 years. *Climate Research* 23: 89–110.

Wiles GC, Barclay DJ and Calkin PE (1999) Tree-ring-dated ‘Little Ice Age’ histories of maritime glaciers from western Prince William Sound, Alaska. *The Holocene* 9: 163–173.

Winkler S (2004) Lichenometric dating of the ‘Little Ice Age’ maximum in Mt Cook National Park, Southern Alps, New Zealand. *The Holocene* 14: 991–920.

Zalasiewicz J, Waters CN, Williams M et al. (2015) When did the Anthropocene begin? A mid-twentieth century boundary level is stratigraphically optimal. *Quaternary International* (in press)
DOI: 10.1016/j.quaint.2014.11.045