



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under Creative Commons: Attribution-Noncommercial-Share Alike 4.0 license:

Fryer, Simon M ORCID logoORCID: <https://orcid.org/0000-0003-0376-0104>, Cornwall, J, Stone, Keeron J ORCID logoORCID: <https://orcid.org/0000-0001-6572-7874> and Stoner, Lee (2015) Should flow-mediated dilation be conducted on the legs of patients with peripheral arterial disease? *Journal of the American Heart Association*, 4. pp. 1-8.

Official URL: <http://jaha.ahajournals.org/content/by/year>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/3090>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document. This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-nc-sa/4.0/):

Fryer, Simon M and Cornwall, J and Stone, Keeron J and Stoner, Lee (2015). *Should flow-mediated dilation be conducted on the legs of patients with peripheral arterial disease?* Journal of the American Heart Association, 4, 1-8.

Published in Journal of the American Heart Association, and available online at:

<http://jaha.ahajournals.org/content/by/year>

We recommend you cite the published (post-print) version.

The URL for the published version is <http://jaha.ahajournals.org/content/by/year>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Should flow-mediated dilation be conducted on the legs of patients with peripheral arterial disease?

- [Simon Fryer](#), Senior Lecturer, Exercise Physiology,
- Jon Cornwall, Keeron Stone, and Lee Stoner

University of Gloucestershire

The recent study "Local Association Between Endothelial Dysfunction and Intimal Hyperplasia: Relevance in Peripheral Artery Disease" assessed endothelial function, using flow-mediated dilation (FMD), in the legs (superficial femoral artery) and arms (brachial artery) of patients with peripheral arterial disease (PAD).¹ The authors concluded that endothelial function was not impaired in the arms, but was reduced in the legs of PAD patients compared to age-matched controls. While the article supports previous research suggesting that the brachial artery may not provide a global index of endothelial health for all populations, including those with spinal cord injury,² it is contrary to two previous PAD studies reporting reduced brachial FMD.^{3, 4} This publication is particularly timely as the clinical relevance of FMD has recently come under attack for reasons such as poor measurement reliability, lack of appropriate covariate analysis, concerns over expressing FMD as a ratio⁵, and improper normalization to the FMD stimulus.⁵ We suggest each of these concerns are highlighted in this study.

First, with the between-day coefficient of variation for the standard brachial FMD test being as high as 50%,⁶ no mention of power analysis or variance estimates and small group sizes ($n=10$), the analyses potentially suffer from a lack of statistical power. Second, FMD was expressed as ratio, which can result in mathematical error propagation and allometric scaling issues.⁵ FMD should be expressed as the absolute change score, using baseline diameter and shear stress as covariates.^{5, 7} Last and perhaps most important, FMD can only be assessed if reactive hyperemia induces a sufficient shear stress stimulus to activate the endothelial cells, which may be prevented by an obstructed arterial lumen in PAD. In addition, an obstructed arterial lumen would induce a non-laminar flow profile, thereby invalidating one of the assumptions of Poiseuille's Law-based shear stress calculations.

Because of these issues, the published findings should be interpreted with caution. While we commend the authors on conducting important work in this field, we suggest additional studies are required to elucidate whether FMD in the lower limbs is a true marker of endothelial function in patients with PAD.

References

1. Heinen Y, Stegemann E, Sansone R, Benedens K, Wagstaff R, Balzer J, Rassaf T, Lauer T, Kelm M, Heiss C. Local association between endothelial dysfunction and intimal hyperplasia: relevance in peripheral artery disease. *J Am Heart Assoc.* 2015;4: 1–8.

2. Stoner L, Sabatier M, VanhHiel L, Groves D, Ripley D, Palardy G, McCully K. Upper vs lower extremity arterial function after spinal cord injury. *The Journal of Spinal Cord Medicine*. 2006;29: 138–146.
3. Brevetti G, Silvestro A, Di Giacomo S, Bucur R, Di Donato A, Schiano V, Scopacasa F. Endothelial dysfunction in peripheral arterial disease is related to increase in plasma markers of inflammation and severity of peripheral circulatory impairment but not to classic risk factors and atherosclerotic burden. *J Vasc Surg*. 2003;38: 374–9.
4. Brevetti G, Silvestro A, Schiano V, Chiariello M. Endothelial dysfunction and cardiovascular risk prediction in peripheral arterial disease: additive value of flow-mediated dilation to ankle-brachial pressure index. *Circulation*. 2003;108: 2093–8.
5. Atkinson G, Batterham AM. The clinical relevance of the percentage flow-mediated dilation index. *Curr Hypertens Rep*. 2015;17: 4.
6. De Roos NM, Bots ML, Schouten EG, Katan MB. Within-subject variability of flow-mediated vasodilation of the brachial artery in healthy men and women: implications for experimental studies. *Ultrasound Med Biol*. 2003;29: 401–6.
7. Stoner L, Tarrant MA, Fryer S, Faulkner J. How should flow-mediated dilation be normalized to its stimulus? *Clin Physiol Funct Imaging*. 2013;33: 75–8.

Conflict of Interest:

None declared

Published July 16, 2015