



This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0 license:

**Alexanders, Jenny, Anderson, Anna and Henderson, Sarah L  
ORCID logoORCID: <https://orcid.org/0000-0003-0723-8057>  
(2015) Musculoskeletal physiotherapists' use of psychological  
interventions A Systematic Review. *Physiotherapy*, 101. pp.  
95-102. doi:10.1016/j.physio.2014.03.008**

Official URL: <http://dx.doi.org/10.1016/j.physio.2014.03.008>

DOI: <http://dx.doi.org/10.1016/j.physio.2014.03.008>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/4211>

#### **Disclaimer**

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

**Musculoskeletal physiotherapists' use of psychological interventions: A  
systematic review of therapists' perceptions and practice.**

**Jenny Alexanders<sup>a</sup>, Anna Anderson<sup>b</sup>, Sarah Henderson<sup>a</sup>**

*<sup>a</sup>Sport, Health and Sciences Department. The University of Hull, Don Building, Cottingham  
Road, Hull, HU67RX, United Kingdom*

*<sup>b</sup>Leeds Teaching Hospitals, Beckett Street, Leeds, LS9 7TF, United Kingdom*

## **Abstract**

**Background:** Research has demonstrated that incorporating psychological interventions within physiotherapy practice has numerous potential benefits. Despite this physiotherapists have reported feeling inadequately trained to confidently use such interventions in their day-to-day practice.

**Objective:** To systematically review musculoskeletal physiotherapists' perceptions regarding the use of psychological interventions within physiotherapy practice.

**Data sources:** Eligible studies were identified through a rigorous search of AMED, CINAHL, EMBASE, MEDLINE and PsychINFO from January 2002 until August 2013.

**Study eligibility criteria:** Full text qualitative, quantitative and mixed methodology studies published in English language investigating musculoskeletal physiotherapists' perceptions regarding their use of psychological interventions within physiotherapy practice.

**Study synthesis and appraisal:** Included studies were appraised for risk of bias using the Critical Appraisal Skills Programme qualitative checklist. Meta-analysis was not possible due to study heterogeneity.

**Results** Six studies, all with a low risk of bias, met the inclusion criteria. These studies highlighted that physiotherapists appreciate the importance of using psychological interventions within their practice, but report inadequate understanding and consequent underutilisation of these interventions.

**Limitations** These results should be noted with some degree of caution due to various limitations associated with the included studies and with this review, including the use of a qualitative appraisal tool for mixed methodology/quantitative studies.

## **Conclusion**

These findings suggest that musculoskeletal physiotherapists are aware of the potential benefits of incorporating psychological interventions within their practice but feel insufficiently trained to optimise their use of such interventions; hence highlighting a need for further research in this area and a review of physiotherapist training.

*Keywords: Rehabilitation, Psychological Interventions, Qualitative design, Physiotherapy Education, Sports.*

## **Introduction**

Research has demonstrated that individuals undergoing rehabilitation following an injury or surgery experience not only physical but also psychological disturbances (1). Diverse symptoms such as anger, depression and low self-esteem have been observed (2) and are reported to occur in both the general population (3) and in sports people (4). The importance of recognising such psychological disturbances is widely recognised amongst physiotherapists, with the Chartered Society of Physiotherapy (CSP) describing physiotherapy as a profession which “takes a ‘whole person’ approach to health and wellbeing” (5). This is reflected in a broad range of documents which guide physiotherapy practice.

For example the post outlines of physiotherapists working within the National Health Service are based on the Knowledge and Skills Framework. Dimensions HWB6 and HWB7 of this framework emphasise the importance of considering both physiological and psychological factors throughout assessment and treatment planning and implementation (6). Furthermore all physiotherapists are legally obliged to adhere to the Health and Care Professions Council’s standards of proficiency for physiotherapists (7), with standard 13.9 indicating that physiotherapists must understand how psychological aspects influence individuals’ responses to their health status and physiotherapy interventions

Correspondingly physiotherapy degrees must now incorporate training on psychology included topics such as self-efficacy in order to achieve CSP accreditation (8). Despite this a recent mixed methods study of 17 United Kingdom universities demonstrated that although some psychology training is included in physiotherapy programmes significant disparities exist in the extent of training provided and how it is delivered (9). Additionally this study highlighted the vast array of different topic areas that are considered under the umbrella term

“psychology”, including communication skills, personality theories, effective team working and the psychological impact of terminal illness.

One area of psychology of particular importance to physiotherapists is the use of psychological interventions such as relaxation, imagery, positive self-talk and goal setting, as all these techniques are used by physiotherapists in clinical practice (2), (10). Furthermore the use of psychological interventions appears to be growing in certain disciplines of physiotherapy; namely chronic pain (11), anterior cruciate ligament rehabilitation (10) neurology (12, 13) and sport (2). Techniques such as cognitive behavioural therapy (CBT), neuro-linguistic programming (NLP) are also other psychological techniques widely being used by physiotherapists within these areas (14, 15). Despite this, undergraduate physiotherapy degree programmes are not specifically required to include training on psychological interventions in order to achieve CSP accreditation (8) and evidence suggests that physiotherapists frequently show a lack of insight into the psychological tools that are within the scope of physiotherapy practice (16). Investigating physiotherapists’ perceptions and use of psychological interventions could therefore provide valuable information about the training needs of qualified physiotherapists and inform the development of future physiotherapy programme curricula.

Musculoskeletal disorders are the commonest presenting complaint encountered by physiotherapists and promoting self-management techniques is recognised as a vital component of the physiotherapy management of these disorders (17). Self-management techniques may include both physical interventions, such as exercise, and psychological interventions, such as mindfulness based stress reduction (18). Investigating the use of psychological interventions in musculoskeletal physiotherapy is therefore an important area for research and will be the focus of this review.

The aim of this review was to investigate musculoskeletal physiotherapists' perceptions and use of psychological interventions, including whether they feel sufficiently equipped to use such interventions effectively within their daily practice.

## **Methods**

This systematic review was conducted using a predetermined protocol in accordance with the PRISMA statement (19).

### *Protocol and registration*

No prior protocol was published.

### *Data sources and search strategy*

Comprehensive electronic searches were undertaken of the following electronic databases: AMED, CINAHL, EMBASE, MEDLINE and PsychINFO. A record of the number of articles that were retrieved from each database is displayed in table 1. In addition, hand searches of journals and citation tracking of reference lists related to the research title were performed. The search terms used for the databases are displayed in table 2. All the above searches were performed by one investigator (JA).

### *Trial selection*

All articles identified by the searches were assessed for eligibility using the criteria described below. Full text copies of any potentially relevant articles were obtained to confirm eligibility. The trial selection was performed by one investigator (JA).

### *Study selection*

The electronic search results were considered for inclusion if they were empirical qualitative, quantitative or mixed methodology studies focussing on Chartered physiotherapists' perceptions regarding their use psychological interventions within musculoskeletal outpatients physiotherapy departments. Only full text papers published in English language dated from 2002 to 2013 were considered. Abstracts alone and systematic reviews were excluded; however the reference lists of any relevant systematic reviews were screened for potentially relevant studies. Sifting was performed using a recommended three stage process (20). Papers were first reviewed by title, then by abstract and finally by full text, excluding those at each stage that did not fit the inclusion and exclusion criteria (20).

### *Data extraction*

Data regarding the characteristics, participants, interventions, outcome measures and results of each study selected for inclusion were extracted using a standardised form (table 3). This form was completed by one reviewer (JA) and verified by two independent reviewers (AA/SH).



### *Study appraisal and synthesis methods*

The included studies were appraised using the Critical Appraisal Skills Programme (CASP) Qualitative Research checklist (21). This checklist comprises a series of 10 questions whose completion provides a systematic and comprehensive way of appraising qualitative studies to determine whether their findings are valid and/or meaningful at a local level. The included studies were appraised by three independent reviewers (JA/AA/SH). Any discrepancies between the reviewers' completed checklists were resolved by discussion, with a fourth independent reviewer being available to help resolve any disagreements if necessary.

## **Results**

The PRISMA flow chart demonstrates the study selection process (figure 1). The electronic searches retrieved a total of 178 articles and one further article was retrieved through hand searching. Removal of duplicates left a total of 178 articles. Screening of the titles and abstracts of these articles resulted in 10 studies being identified as appropriate for full text review. After obtaining full text copies of each of these articles a total of 6 were accepted for inclusion in this systematic review (22-27).

### *Risk of bias assessment*

The results of the risk of bias assessments are presented in table 4. All studies were found to present a low risk of bias in accordance with the CASP guidelines (21). All reviewers (JA/AA/SH) used the same method to review the selected articles and were satisfied that the overall risk of bias was low. There was only a single disagreement between

the reviewers' completed risk of bias checklists. This was settled by discussion without requiring input from the fourth independent reviewer.

### *Study characteristics*

The results of the included studies, along with the characteristics of each study are presented in table 5. All studies involved exploration of musculoskeletal physiotherapists' perceptions regarding the use of psychological interventions in physiotherapy practice (22-27). Three studies focused on physiotherapists working predominately in a sports environment (22, 23, 25). The other three studies focused on physiotherapists working in general musculoskeletal outpatient settings (24, 26, 27). Three studies used qualitative semi-structured interviews (22, 26, 27), two studies used quantitative surveys (24, 25) and one study used a mixed methodology (23). Five studies sought to gain an insight into physiotherapists' perceived learning needs regarding psychological training (22-26). All the studies investigated common psychological symptoms that physiotherapists encounter within clinical practice (22-27).

### *Interventions*

No interventions were used.

## *Findings*

According to the physiotherapists in the included studies, stress and anxiety were the most frequently encountered psychological symptoms amongst individuals undergoing rehabilitation (23-25). Additional reported psychological symptoms included exercise addiction (24) and fear of re-injury (26). The findings of the included studies suggest that physiotherapists commonly use a number of psychological interventions in clinical practice. In particular goal setting, positive self-talk, effective communication and variation in rehabilitation exercises all appear to be relatively widely used (22-27). From a learning needs perspective, the included studies suggest that physiotherapists would like to improve their ability to implement realistic goal setting (24, 25). Furthermore physiotherapists from four of the included studies reported feeling that they received insufficient psychological training during their undergraduate physiotherapy degree programmes (22, 23, 26, 27). For example in the study of Arvinen-Barrow et al. (2010) it was reported that one respondent had only “a couple of lectures of psychology as part of his physiotherapy degree”, while another respondent reported having “no training in psychological interventions at all”.

## **Discussion**

This systematic review evaluated six studies that investigated musculoskeletal physiotherapists' perceptions regarding the use of psychological interventions within physiotherapy practice. The results present some interesting but concerning findings. It was reported that physiotherapists working in a musculoskeletal environment commonly encounter psychological symptoms when working with individuals undergoing rehabilitation. A wide variety of psychological symptoms were identified amongst individuals undergoing rehabilitation, with stress and anxiety appearing to be particularly common (23-25). This

latter finding is supported by a number of additional studies (28-30). Furthermore a prospective cohort study reported that patients who suffer psychological symptoms following surgery/ injuries appear to experience an increase in non-specific musculoskeletal complaints, such as increased muscular tension, development of trigger points and reduced function (31). The above findings suggest that the occurrence of psychological symptoms following injury/surgery is a frequent and significant issue. One potentially valuable approach in tackling this issue is to integrate psychological interventions within physiotherapy practice.

The results of this systematic review imply that a number of psychological interventions are widely used by musculoskeletal physiotherapists. These interventions include: goal setting, positive self-talk and effective communication, with goal setting being identified as the most common psychological intervention used by the physiotherapists in the included studies (22-27). This is a particularly encouraging finding given that the use of goal setting during rehabilitation appears to improve patients' confidence, self-esteem and self-efficacy (32).

Correspondingly goal setting was also highlighted as an intervention that physiotherapists would like to learn more about (23, 24). The study of Schoeb (2009) provides a particularly detailed insight into physiotherapists' use of goal setting in clinical practice (33). This study indicated that although this intervention is frequently employed by physiotherapists the approaches used for setting goals are rarely standardised. This study also highlighted that the effectiveness of the goal setting process appears to vary significantly between different physiotherapists (33). In addition another study, involving various different healthcare professionals, indicated that goal setting within healthcare is associated with numerous difficulties, such as poor patient involvement throughout the goal setting process (34).

Arvinen-Barrow et al.'s (2010) study implied that goal setting is rarely athlete-focussed, instead being largely physiotherapist-mandated with minimal consideration being given to its psychological implications (22). This finding is particularly significant given that patient participation appears to be of paramount importance in the success of the goal setting process, largely due to the numerous beneficial effects associated with such patient participation (35). These include increasing patient participation during rehabilitation, empowering of patients to take responsibility for their recovery and making the rehabilitation process more meaningful to patients (35). Overall the articles included in this systematic review indicate a disparity between physiotherapists' reported use of goal setting and the training provided in this area during undergraduate physiotherapy programmes. Future research using observational methods to investigate physiotherapists' implementation of goal setting in clinical practice would provide further insight into this area and be of value in identifying physiotherapists' training needs.

The results of this systematic review imply that at least some physiotherapy degree programmes do not provide a sufficient level of formal training in the use of psychological interventions. Until the 1980's minimal training in psychology was provided within physiotherapy curricula (36). In the early 1980's the Scientific Affairs Board of the British Psychological Society (BPS) set up a working party aiming to teach different aspects of psychology, including basic theory and the implementation of psychological principles to individuals from other professions, physiotherapy being one of the named professions. Despite this recent studies indicate that the psychology content of UK physiotherapy degree programmes remains inconsistent, with many universities failing to have integrated psychology training within their physiotherapy curricula (37). In support of this a number of studies have indicated that physiotherapists report feeling inadequately trained to deliver psychological interventions and would welcome further training in this field (22, 24, 38). In

light of the high prevalence of psychological issues amongst individuals undergoing rehabilitation and the apparent value of incorporating psychological interventions within rehabilitation, these findings highlight a clear need to review the design of physiotherapy degree programmes to ensure that appropriate training in the use of psychological interventions becomes an integral part of physiotherapy curricula.

### *Limitations of the included studies*

The included studies had a number of limitations. Firstly some of the included studies were largely focused on independent physiotherapists working in sport (22, 23, 25); the results cannot therefore be extrapolated to National Health Service physiotherapists working in hospital-based settings. This is especially relevant given that physiotherapists working in sports settings may have completed a sports psychology module as part of their training (39). Sample sizes ranged from 7 to 90 participants (22-27). Whilst the small sample sizes of the qualitative studies do not limit the utility of their findings, it could be argued that the sample sizes of the quantitative studies were inadequate; hence not providing a true representation of chartered physiotherapists currently practising in the UK.

Another limitation of the included studies is that they did not investigate whether respondents' answers varied according to how long they had been qualified as a physiotherapist. This is significant given that the time since qualification is likely to have affected the extent to which the physiotherapists relied on formal psychological training as opposed to experiential learning. This in turn may have directly influenced the physiotherapists' use and perceptions of psychological interventions. Furthermore the actual content of the physiotherapists' formal training in psychological interventions was not verified with the organisations providing the training; hence the results does not account for

possible disparities between the physiotherapists' perceived past training and the physiotherapists' actual past training. Finally, minimal information was provided on where the respondents completed their physiotherapy training; which again is likely to have had a profound influence on their responses.

### *Implications for practice*

Despite the aforementioned paucity of training in psychological interventions within physiotherapy degree programmes, the use of psychological interventions does appear to be an integral part of musculoskeletal physiotherapy practice [10]. It should therefore be recognised that there is a need to provide physiotherapists with the skills to effectively incorporate psychological interventions within rehabilitation. This would help physiotherapists to maximise the potential of each individual patient, for example by ensuring that when goals are set they are both patient-led and not just therapist led. This therefore indicates a need to review the curricula of physiotherapy undergraduate/pre-registration degrees to ensure that training in the use of psychological interventions is standardised.

### *Implications for future research*

This systematic review highlights that research into musculoskeletal physiotherapists' perceptions of using psychology interventions during rehabilitation and into their practical use of such interventions are both limited. Given the frequent occurrence of psychological disturbances post-injury/surgery and the powerful influence an individual's psychological state appears to have on the rehabilitation process further research in this area is clearly warranted. Possible aims of such research could include identifying how physiotherapists

actually implement goal setting strategies and investigating physiotherapists understanding of the theory behind the psychological interventions they use in practice.

Future research should include both qualitative studies and quantitative studies. For example randomised controlled trials investigating the effectiveness of a range of psychological interventions could provide highly valuable information about the worth and relative effectiveness of such interventions. It is also essential that studies are conducted in both sporting and non-sporting populations given that their psychological responses to injury/surgery and to physiotherapy interventions are likely to differ. Furthermore a particularly useful area for future research would be to investigate which psychological theories and interventions should be prioritised for inclusion in physiotherapy undergraduate/pre-registration degrees and which would be better suited for inclusion in post-graduate training courses.

#### *Limitations of this review*

This study did not accept any unpublished papers; therefore potentially relevant information from studies which have been conducted but not published may have been missed. Furthermore this review only included articles published between 2002 and 2013; hence relevant information from papers published before 2002 may have been omitted. However papers published before 2002 are unlikely to provide representative information about the current perspectives of physiotherapists given people's perspectives are fluid and are likely to be influenced by the current society in which they live. Additionally physiotherapy degree curricula and training methods are also very likely to have changed since 2002.



This systematic review did not exclude studies which investigated the perceptions of other rehabilitation professionals in addition to Chartered physiotherapists. This limits the validity of the findings given that the conclusions drawn may have been significantly influenced by the views of respondents who were not physiotherapists. This systematic review is also limited by its use of a critical appraisal tool which was designed for use in qualitative studies, despite this systematic review including qualitative, quantitative and mixed methodology studies. In addition the CASP qualitative checklist used appears to be inferior to other qualitative appraisal tools in terms of its sensitivity to certain aspects of validity (40). Finally this review was specifically focused on physiotherapists working in musculoskeletal outpatient setting; hence the results cannot be applied to physiotherapists working in alternative disciplines.

## **Conclusion**

The results of this systematic review highlight that musculoskeletal physiotherapists recognise the potential value of incorporating psychological interventions into their rehabilitation programmes for individuals following injury/surgery. Furthermore evidence suggests that physiotherapists currently use various psychological interventions such as goal setting in clinical practice. Despite this physiotherapists do not feel adequately trained to effectively utilise either the psychological interventions that they are currently using, or additional psychological interventions that they do not use at present, but which are likely improve rehabilitation outcomes. These findings indicate a clear need for further research in this area and imply that a review of the inclusion of training in psychological interventions within physiotherapy degree programmes is certainly warranted.

**Ethical approval**

Not applicable

**Funding**

None

**Conflict of interest**

None

## References

1. Walker N, Thatcher J, Lavalley D. Psychological responses to injury in competitive sport: a critical review. *The Journal of the Royal Society for the Promotion of Health*. 2007 Jul;127(4):174-80. PubMed PMID: 17711063. Epub 2007/08/23. eng.
2. Schwab Reese LM, Pittsinger R, Yang J. Effectiveness of psychological intervention following sport injury. *Journal of Sport and Health Science*. 2012 9//;1(2):71-9.
3. Pearson L, Jones G. Emotional Effects of Sports Injuries: Implications for Physiotherapists. *Physiotherapy*. 1992 10/10//;78(10):762-70.
4. Paulsen MG, Dowsey MM, Castle D, Choong PFM. Preoperative psychological distress and functional outcome after knee replacement. *ANZ Journal of Surgery*. 2011;81(10):681-7.
5. Physiotherapy CSo. What do physiotherapists do? 2014 [cited 2014 2014]. Available from: <http://www.csp.org.uk/your-health/what-physiotherapy>.
6. Health Df. The NHS Knowledge Skills Framework and the development review process. In: NHS, editor. 2004. p. 23.
7. Council HaCP. Standards of proficiency for physiotherapists. In: NHS, editor. 2013. p. 20.
8. Physiotherapy CSo. CSP learning and development principles. In: CSP, editor. London2002. p. 18.
9. Heaney C, Alison G, Rostron C, Walker N. A qualitative and quantitative investigation of the psychology content of UK physiotherapy education. *Journal of Physical Therapy Education*. 2012;26(3):1-27.
10. Cupal D, D., Brewer B, W. Effects of Relaxation and Guided Imagery on Knee Strength, Reinjury Anxiety, and Pain Following Anterior Cruciate Ligament Reconstruction. *Rehabilitation Psychology*. 2001;46(1):28-43.
11. Harland N, Lavalley D. Biopsychosocial Management of Chronic Low Back Pain Patients with Psychological Assessment and Management Tools: Overview. *Physiotherapy*. 2003 5//;89(5):305-12.
12. Scobbie L, McLean D, Dixon D, Duncan E, Wyke S. Implementing a framework for goal setting in community based stroke rehabilitation: a process evaluation. *BMC Health Services Research*. 2013;13(1):190. PubMed PMID: doi:10.1186/1472-6963-13-190.
13. Scobbie L, Dixon D, Wyke S. Identifying and applying psychological theory to setting and achieving rehabilitation goals. *Clinical rehabilitation*. 2009;23:231 - 333.
14. Brunner E, De Hart A, Minguet P, Baldew S, Probst M. Can cognitive behavioural therapy based strategies be integrated into physiotherapy for the prevention of chronic low back pain? A systematic review

Disability and Rehabilitation. 2013;35(1):1-10.

15. Karunaratne M. Neuro-linguistic programming and application in treatment of phobias. *Complementary Therapies in Clinical Practice*. 2010;16(4):203-7.
16. Arvinen-Barrow M, Penny G, Hemmings B, Corr S. UK chartered physiotherapists' personal experiences in using psychological interventions with injured athletes: An Interpretative Phenomenological Analysis. *Psychology of Sport and Exercise*. 2010;11:58-66.
17. Physiotherapy CSo. Physiotherapy works musculoskeletal disorders. In: CSP, editor. 2014. p. 1-2.
18. Dufour S, Graham S, Friesen J, Rosenblat M, Rous C, Richardson J. Physiotherapists supporting self-management through health coaching: a mixed methods program evaluation. *Physiotherapy theory and practice*. 2014;1:1-10.
19. Moher D LA, Tetzlaff J, Douglas, Altman D. . Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Physical Therapy in Sport*. 2009;89(9):873-80.

20. Jones ML. Application of systematic review methods to qualitative research: practical issues. *Journal of advanced nursing*. 2004 Nov;48(3):271-8. PubMed PMID: 15488041. Epub 2004/10/19. eng.
21. Anon. Critical Appraisal Skills Programme (CASP) <http://www.casp-uk.net2012> [cited 2012 22/6/2012].
22. Arvinen-Barrow M, Penny G, Hemmings B, Corr S. UK chartered physiotherapists' personal experiences in using psychological interventions with injured athletes: An Interpretative Phenomenological Analysis. *Psychology of Sport and Exercise*. 2010 1//;11(1):58-66.
23. Heaney C. Physiotherapists' perceptions of sport psychology intervention in professional soccer. *International Journal of Sport and Exercise Psychology*. 2006 2006/01/01;4(1):73-86.
24. Hemmings B, Povey L. Views of chartered physiotherapists on the psychological content of their practice: a preliminary study in the United Kingdom. *British journal of sports medicine*. 2002 Feb;36(1):61-4. PubMed PMID: 11867495. Pubmed Central PMCID: PMC1724459. Epub 2002/02/28. eng.
25. Lafferty M, Kenyon, R, Wright, C. Club-based and non-club based physiotherapist's views on their psychological content of thier practice when treating sports injuries. *Research in Sports Medicine*. 2008;16(4):295-306.
26. Tracey J. Inside the clinic: health professionals' role in their clients' psychological rehabilitation. *Journal of sport rehabilitation*. 2008 Nov;17(4):413-31. PubMed PMID: 19160915. Epub 2009/01/24. eng.
27. Jevon SM, Johnston LH. The perceived knowledge and attitudes of governing body chartered physiotherapists towards the psychological aspects of rehabilitation. *Physical Therapy in Sport*. 2003 5//;4(2):74-81.
28. Nicholas MK, Linton SJ, Watson PJ, Main CJ, Group tDotFW. Early Identification and Management of Psychological Risk Factors ("Yellow Flags") in Patients With Low Back Pain: A Reappraisal. *Physical therapy*. 2011 May 1, 2011;91(5):737-53.
29. Tjong VK, Murnaghan ML, Nyhof-Young JM, Ogilvie-Harris DJ. A Qualitative Investigation of the Decision to Return to Sport After Anterior Cruciate Ligament Reconstruction: To Play or Not to Play. *The American journal of sports medicine*. 2013 November 6, 2013.
30. Campbell Burton CA, Murray J, Holmes J, Astin F, Greenwood D, Knapp P. Frequency of anxiety after stroke: a systematic review and meta-analysis of observational studies. *International Journal of Stroke*. 2013;8(7):545-59.
31. Mirza SK, Deyo RA, Heagerty PJ, Turner JA, Martin BI, Comstock BA. One-year outcomes of surgical versus nonsurgical treatments for discogenic back pain: a community-based prospective cohort study. *The Spine Journal*. 2013 11//;13(11):1421-33.
32. Farouk A EA. Impact of psychological rehabilitation program on self confidence level and competition anxiety for soccer players of anterior cruciate ligament injury. *World Journal of Sport Sciences*. 2010;3:138-43.
33. Schoeb V. "The goal is to be more flexible"--detailed analysis of goal setting in physiotherapy using a conversation analytic approach. *Manual therapy* 2009 Dec;14(6):665-70. PubMed PMID: 19427808. Epub 2009/05/12. eng.
34. Playford ED, Dawson L, Limbert V, Smith M, Ward CD, Wells R. Goal-setting in rehabilitation: report of a workshop to explore professionals' perceptions of goal-setting. *Clinical rehabilitation*. 2000 Oct;14(5):491-6. PubMed PMID: 11043874. Epub 2000/10/24. eng.
35. Filoramo MA. Improving goal setting and goal attainment in patients with chronic noncancer pain. *Pain management nursing : official journal of the American Society of Pain Management Nurses*. 2007 Jun;8(2):96-101. PubMed PMID: 17544129. Epub 2007/06/05. eng.
36. Baddeley H, Bithell C. Psychology in the Physiotherapy Curriculum: A Survey. *Physiotherapy*. 1989;75(1):17-21.

37. Heaney C, Green A, Rostron C, Walker N. A qualitative and quantitative investigation of the psychology content of UK physiotherapy education. *Journal of Physical Therapy Education*. 2012;26(3):1-26.
38. Ford FI GS. Perspectives of sport physiotherapist on the frequency and significant of psychological factors in professional practice: implications for curriculum design in professional training. *The Australian Journal of Science and Medicine in Sport*. 1997;2(34):34-40.
39. Heaney C. The impact of sport psychology education on the practice of physiotherapists. *International Sports Science and Sports Medicine Conference; Newcastle Upon Tyne* 2013.
40. Hannes K LC, Pearson A. A comparative analysis of three online appraisal instruments' ability to assess validity in qualitative research. *Qualitative health research*. 2010;12(1736-43).

Table 1

Tracking of articles

MEDLINE	CINHAL	EMBASE	PyschINFO	AMED
34	22	27	47	48

Table 2

## Database search strategy

MEDLINE	CINAL	EMBASE	PsychINFO	AMED
rehabilitation\$ or Musculoskeletal Physiotherapy\$ or injuries\$ psychology\$ or intervention\$ or symptoms\$ or physiotherapist/ education\$ or perception\$	(MH"rehabilitation/ surgery") OR (MH"musculoskeletal physiotherapy") OR (MH"injuries") OR (MH"psychology") OR (MH"symptoms") OR (MH"physiotherapist") OR (MH"education") OR (MH"perception") OR	'rehabilitation'/ surgery OR 'musculoskeletal physiotherapy'/exp OR 'injuries'/exp OR 'psychological symptoms'/exp OR 'physiotherapist education'/exp OR 'perception'	rehabilitation/ surgery or musculoskeletal physiotherapy/ injuries/ psychology/ or symptoms/ or physiotherapist education/ or perception	rehabilitation AND injuries AND musculoskeletal physiotherapy AND psychology AND symptoms AND physiotherapist AND education AND perception

Table 3

Data extraction form

Study characteristics	Participant characteristics	Intervention & setting	Outcome data/ results
Arvinen-Barrow (2010) (22)			
Heaney (2006) (23)			
Hemmings, Povey (2002)			
Lafferty, Kenyon & Wright (2008) (25)			
Tracey (2008) (26)			
Jevon & Johnson (2003) (27)			

Table 4

Completed qualitative risk of bias checklist tool (CASP)

Was there a clear statement	Is a qualitative methodology	Was the research design	Was the recruitment strategy	Was the data collected in a way that	Has the relationship between the	Have ethical issues been taken in to	Was the data analysis	Is there a clear statement of	How valuable is the research?
-----------------------------	------------------------------	-------------------------	------------------------------	--------------------------------------	----------------------------------	--------------------------------------	-----------------------	-------------------------------	-------------------------------



	of the research aim?	appropriate for the research?	appropriate to address the aims of the research?	appropriate to the aims of the research?	addressed the research issue?	researcher and participants been adequately considered?	consideration?	sufficiently rigorous?	findings?	
Arvinen- Barrow et al. (2010) (22)	Y	Y	Y	N	Y	Y	Y	Y	Y	Very
Heaney (2006) (23)	Y	Y	Y	Y	Y	Y	N	N	Y	Moderately
Hemmings (2002) (24)	Y	Y	Y	N	Y	Y	N	N	Y	Moderately
Lafferty, Kenyon, Wright (2008) (25)	Y	N	Y	Can't tell	Y	Y	Y	Y	Y	Moderately
Tracey (2008) (26)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Moderately
Jevon, Johnson (2003) (27)	Y	Y	Y	Can't tell	Y	Can't tell	N	Y	Y	Very

Table 5

## Characteristics of included studies

Study characteristics	Participant characteristics	Intervention & setting	Outcome data/ results
Arvinen-Barrow (2010) (22) Qualitative semi structured interviews using interpretative phenomenological analysis. Aim of study was to investigate the views of physiotherapists using psychological interventions during rehabilitation of injured individuals	7 full time Chartered physiotherapists working in sport medicine. (4 females, 3 males).	Sport/clinic setting. Semi structured interview questions investigating physiotherapists' experiences using psychological techniques in rehabilitation, process of psychological rehabilitation of the injured athlete.	Goal setting, imagery and positive-self talk were the most common psychological interventions used in this study. However goal setting was only physiotherapist-mandated and not athlete focused or psychologically focused. Lack of formal training was highlighted as a barrier to physiotherapists fully understanding psychological interventions.
Heaney (2006) (23) Mixed methods questionnaires using the physiotherapist and sport psychology questionnaire (PSPQ). The participants then underwent semi structured interviews based on the questionnaire. Aim of study was to investigate perceptions of physiotherapists using psychology in professional soccer.	39 head physiotherapists working in sport (2 females, 37 males).	Sport/clinic setting. Semi structured interview questions investigating the psychology content of physiotherapy practice, sport psychology referrals and delivery of sports psychology.	Stress and anxiety were the most common psychological symptoms reported amongst injured athletes. Goal setting, positive self-talk and ensuring variety in the rehabilitation programme were the most common psychological techniques used by physiotherapists. Realistic goal setting was highlighted as one of the most important skills respondents believed physiotherapists should learn.
Hemmings, Povey (2002) (24) Quantitative survey using the physiotherapist and sports psychology questionnaire (PSPQ). Aim of study was to investigate physiotherapists' views of using psychology as part of their practice.	90 musculoskeletal (MSK) physiotherapists working in a clinical environment. (67 females, 23 males).	Clinic setting, survey investigated psychological responses encountered by physiotherapists, psychological techniques used by physiotherapists, psychological techniques that physiotherapists would like to learn more about.	Stress, anxiety, exercise addiction and depression and were common problems encountered by physiotherapists. Most common intervention techniques used by physiotherapists were goal setting, positive self-talk, and variation in rehabilitation and communication skills. Techniques physiotherapists would like to improve their understanding of were: setting realistic goals and motivation techniques.

Table 5 continued

Study characteristics	Participant characteristics	Intervention & setting	Outcome data/ results
Lafferty, Kenyon & Wright (2008) (25) Quantitative surveys using the athletic training and sport psychology questionnaire (ATSPQ). Aim of the study was to investigate whether there were any differences in the psychological content of physiotherapy practice between non-club and club based physiotherapists.	87 physiotherapists (42 non club and 45 club contracted).	Sport/clinic setting. Survey investigated psychological techniques used, psychological techniques physiotherapists would like to learn and common psychological symptoms physiotherapists encounter.	Stress and anxiety were common symptoms among both non club and club physiotherapists. Common techniques used by both groups were creating variation in rehabilitation, goal setting, positive self-talk, increasing confidence. Both groups felt that goal setting was the most important psychological intervention to use with patients.
Tracey (2008) (26) Qualitative semi structured interviews using interpretational analyses. Aim of the study was to investigate therapists' views on their roles in the psychological recovery of injured athletes.	18 participants (1 athletic trainer and 17 physical therapists) (12 females, 6 males).	Clinic and sports environment. Semi structured interviews investigating: perceptions of the roles and strategies they use regarding patient recovery, perceptions regarding influences on patient recovery during rehabilitation.	Common symptoms to address were fear of re-injury, reduced confidence. The main psychological techniques that respondents reported focussing on were building a rapport with clients, educating clients appropriately and communicating with clients effectively. The study concluded that providing health professionals with more specific psychological training would enable them to be more effective when rehabilitating clients
Jevon & Johnson (2003) (27) Qualitative semi structured interviews. Aim of the study was to investigate the perceived attitudes of physiotherapists regarding their use of psychological interventions in rehabilitation, their need for further training in this area and their access to appropriate psychologists.	19 physiotherapists working in MSK.	Out patients setting. Semi-structured interviews included questions regarding common psychological responses to injury, factors affecting rehabilitation, role of physiotherapists in the psychological rehabilitation of individuals following injury/surgery	Stress and anxiety were reported to be problems that physiotherapists felt they needed to address during rehabilitation. Common techniques used were goal setting, effective communication and relaxation. Most physiotherapists felt they lacked any theoretical understanding of psychology due to not being provided with this at undergraduate level.

Figure 1. PRISMA flow diagram of search strategy

(Adapted from Moher D, Liberati A, Tetzlaff J, Douglas, Altman D. Preferred Reporting Items for Systematic Reviews and Meta Analyses: The PRISMA Statement. Physical Therapy 2009;89(9):873-80)

