



UNIVERSITY OF
GLOUCESTERSHIRE

This is additional metadata of the following published document and is licensed under All Rights Reserved license:

**Harun, Sahana, Baker, Andy, Bradley, Chris, Pinay, Gilles, Boomer, Ian and Hamilton, R. Liz ORCID logoORCID:
<https://orcid.org/0000-0003-2532-2236> (2015)
Characterisation of dissolved organic matter in the Lower
Kinabatangan River, Sabah, Malaysia. Hydrology Research, 46
(3). pp. 411-428. doi:10.2166/nh.2014.196**

Official URL: <http://dx.doi.org/10.2166/nh.2014.196>

DOI: <http://dx.doi.org/10.2166/nh.2014.196>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/4464>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Abstract

Spatial patterns and trends in the concentration and quality of dissolved organic matter (DOM) are characterised across a tropical agricultural catchment using UV-visible absorbance, and fluorescence spectroscopy. Related determination of the environmental isotopes ^{18}O and ^2H clarify the dynamics of catchment water movement. Water samples were collected from the Kinabatangan River, Borneo, and selected tributaries in August and September 2008 in four regions with oil palm plantations (KB1, KB2, KB3 and KB4). The isotopic compositions of surface waters suggest that canals sampled in the vicinity of oil palm plantation were characterised by a strong evaporative effect than tributaries and streams with more natural, forested vegetation. DOM was characterised by variations in UV absorbance and spectral slope. Individual fluorescence excitation-emission matrices (EEMs) were decomposed by Parallel Factor Analysis (PARAFAC) and three components extracted (C1, C2 and C3). Components C2 and C3 both appear to be derived from microbial sources and/or photo-degradation. The PARAFAC components indicate a clear trend of increasing DOM degradation as waters pass through the catchment. It is hypothesised that upstream DOM is rapidly photo- and microbially- degraded to less fluorescent DOM, whilst DOM concentration and character of DOM downstream is controlled by the hydrology, specifically by variations in the rate of water movement.