

This is a peer-reviewed, final published version of the following document, © Michal Lehnert and is licensed under Creative Commons: Attribution 4.0 license:

Lehnert, Michal, Hůlka, Karel, De Ste Croix, Mark B ORCID: 0000-0001-9911-4355 and Horutová, Krystýna (2018) Acute effect of basketball-specific exercise on lower limb injury risk mechanisms in male basketball players U16 and U18. Research and Investigations in Sports Medicine, 2 (3). pp. 1-6. ISSN 2577-1914

Official URL: <https://crimsonpublishers.com/rism/fulltext/RISM.000539.php>

DOI: 10.31031/RISM.2018.02.000539

EPrint URI: <http://eprints.glos.ac.uk/id/eprint/5610>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.



Acute Effect of Basketball-Specific Exercise on Lower Limb Injury Risk Mechanisms in Male Basketball Players U16 and U18



Michal Lehnert^{1*}, Karel Hůlka¹, Mark De Ste Croix² and Kristýna Horutová¹

¹Faculty of Physical Culture, Palacký University in Olomouc, Czech Republic

²Exercise and Sport Research Centre, University of Gloucestershire, United Kingdom

*Corresponding author: Michal Lehnert, PhD, Faculty of Physical Culture, Palacký University, Tr Miru 117, 771-11 Olomouc, Czech Republic

Submission: 📅 March 19, 2018; Published: 📅 April 05, 2018

Abstract

The aim of this study was to examine and compare the acute effects of basketball-specific fatigue on muscular stiffness and reactive strength in male adolescent basketball players of two age categories. Reactive strength, expressed by means of the reactive strength index (RSI), and leg stiffness (LS) as mechanisms associated with the risk of ACL injury were measured in eleven players of the U16 category (age 16.1 ± 0.4 years; body height 185 ± 6.5 cm; body mass 74.3 ± 9.9 kg) and ten players of the U18 category (age 17.7 ± 0.4 years; body height 187 ± 5.7 cm; body mass 79.7 ± 7.4 kg) pre and post simulated basketball match play (SBFP₂₈). RSI was determined by a drop jump test, LS by a 20 sub-maximal two-legged hopping test. No significant effects of SBFP₂₈ on the monitored parameters were observed with the exception of RSI in U16 ($p=0.013$, $r=0.53$). Similarly, there were no significant differences in the fatigue-related responses to SBFP₂₈ for RSI and LS between the age groups. These results indicated that irrespective of age, the stabilization function of the knee muscles was not impaired and consequently the risk of ACL injury was not increased.

Keywords: Youth; Fatigue; Neuromuscular; Reactive strength; Stiffness

Introduction

Basketball is a physically demanding sport, which imposes a highly variable load on the players [1]. Basketball performance combines maximum intensity activities and medium or low intensity multidirectional movements such as acceleration and deceleration, pivoting, dribbling, shuffling, jumping and landing, often performed unilaterally [2,3]. For these reasons, youth basketball belongs among risky sports with frequent lower limb injuries including injuries of the knee joint and hamstrings [4]. In 70 to 80 percent of cases, the injury is of a non-contact nature [5]. In particular, anterior cruciate ligament (ACL) injuries are associated with the longest interruption of training and competition, and also predispose to recurrence and other difficulties including earlier degenerative changes [5,6]. Non-contact ACL injuries appear to predominantly occur in a rapid change of direction initiated on one leg, landing from a jump on one leg, and pivoting [6-8]. It was proved that injury risk increases with chronological age and may also be related to important periods during growth and maturation [9]. It was recognized in youth athletes that 16 to 18-year olds represent a group at greatest risk where most injuries are likely to occur [10], which was also confirmed in sports such as basketball

and soccer [11]. Other recent studies of youth reported the highest incidence of injury in the group of 13 to 15-year olds, followed by 16 to 18 year olds [12], and the highest injury incidence around the time of Peak Height Velocity [13].

It was also documented that muscle fatigue is one of the modifiable risk factors of ACL injury [14,15]. Injury incidence data show an increased risk of injury towards the end of matches, when fatigue is most likely to be present [16]. There are a number of mechanisms that relate to an increased risk of knee injury and it was documented that muscular and neuromuscular control is important in reducing load on ACL and thus reducing the risk of ACL injury [17,18]. Fatigue decreases the neuromuscular component of the knee dynamic stability and thus increases the risk of ACL injury [19,20]. In fatigue-related prevention of ACL injuries, a significant indicator is muscle stiffness [21]. Information about lower limb stiffness is particularly important in terms of risk of injury resulting from a jump or change of direction impact [17]. Currently, there are only few studies focusing on the effect of fatigue on lower limb stiffness, mostly in soccer. Some studies claim that lower limb stiffness is compromised as a result of fatigue [22,23], while others

report individualized changes in lower limb stiffness after a fatigue protocol [24]. It was also suggested that the influence of age on the ability to perform eccentric-concentric movements exists and that adults are capable of greater pre-activation and forward control compared with children and adolescents [25-27]. This suggests that children and adolescents represent a risk group in terms of injury [17].

Fatigue may also lead to changes in reactive strength of leg muscles [28]. The level of reactive strength in individuals is assessed by means of the reactive strength index (RSI). RSI represents an individual ability of transition from eccentric to concentric muscle contraction [29] and is determined for monitoring stress exerted on the muscle-tendon complex during plyometric exercise. RSI is also a significant indicator in the prevention of ACL injuries [30]. Studies dealing with the effect of age on the results of RSI point to gradually improving results during adolescence [26,27]. However, there are only few studies dealing with RSI as a result of fatigue in youth athletes, and on top of that with conflicting findings [23,31]. The above-mentioned findings on the occurrence of ACL injuries in team sports indicate that this is a serious phenomenon in which an important role plays the etiology of muscular and neuromuscular control, fatigue, and age. Nevertheless, there is a paucity of research on the influence of acute fatigue from match-play on physiological mechanisms in adolescent populations. Therefore, this study aims to examine and compare the acute effects of basketball-specific fatigue on muscular stiffness and reactive strength in male adolescent basketball players of two age categories.

Methods

Participants

The study involved a group of youth basketball players of the U16 category (N=11; age 16.1±0.4 years; body height 185±6.5; body mass 74.3±9.9kg) and U18 category (N=10; age 17.7±0.4 years; body height 187±5.7cm; body mass 79.7±7.4kg). The players trained on average six times per week and played one competitive match per week during the competitive season. All players tested were fully informed about the aim of the study and the testing procedures. The day before testing, the players were not exposed to any training load. The leg preferred for kicking was considered the dominant leg (DL) and the contra lateral leg as the non-dominant leg (NL). The study was approved by the institution's ethics committee and conformed to the Declaration of Helsinki regarding the use of human subjects. All players tested were free of any musculoskeletal lower-extremity injuries and were fully informed about the aim of the study and the testing procedures. A written informed consent to the testing procedures and the use of the data for further research was obtained from the players' parents.

Procedures

The measurements took place during the 2015 pre-season. During the first session, a week before testing, the players

underwent anthropometric measurements, familiarization with the fatigue protocol, and the tests applied. During the second session, before the testing procedure, the players completed non-specific warm-up exercises, which included cycling on a stationary bicycle ergometer for 5 minutes at 1.5W/kg, dynamic stretching exercises for six minutes, which targeted the main muscle groups involved during the test, and finally fifteen squats. The warm-up routine was performed under the researcher's supervision. The subsequent testing procedure consisted of measuring reactive strength, leg stiffness, and isokinetic strength. Then the players performed the fatigue protocol. Immediately after the fatigue protocol, the tests were repeated in the same order.

Reactive strength

The reactive strength index (RSI) was determined by means of a drop jump test with hands on the hips. The vertical jump performance was assessed using a portable optical timing system Opto jump Next (Microgate, Bolzano, Italy) with manufacturer-declared accuracy of 0.001s. The players jumped from a 30cm high box in order to perform a maximum jump for the shortest period of take-off [32]. To familiarize the players with the testing procedure, two trials were performed followed by three measured attempts with a 30s rest interval between the sets. The greatest value of the three attempts was used in further analyses. RSI was calculated as the ratio between jump height and contact time on the floor [33].

Leg stiffness

To determine stiffness of the lower limbs, all players performed one trial set and two measured sets of 20 sub-maximal two-legged hopping using the Opto jump Next system (Microgate, Bolzano, Italy) with hands on the hips and the torso in an upright position. The frequency of hopping was maintained at 2.5Hz by using a mechanical metronome (Wittner GmbH & Co. KG, Isny, Germany). Of the two sets performed, the set in which the frequency of jumps was closest to the frequency determined by the metronome was selected. From this set, the ten consecutive jumps that were closest to the determined frequency of hopping were used for further analyses [34]. There was a 1-minute rest interval between the sets. Both absolute leg stiffness (ABSLS; kN·m⁻¹) and relative leg stiffness (RELLS; kN·m⁻¹), i.e. normalization of absolute leg stiffness according to leg length and body mass, were calculated [32].

Specific basketball fatigue protocol (SBFP₂₈)

The data are collected over four separate periods as in a match. There is a two-minute break following the first and third periods and a fifteen-minute break following the second period [35]. Each period consists of seventeen activity sections involving maximal activity (sprints), submaximal activity (running at 3.5m·s⁻¹), low 3-5 intensity activity (jogging and walking at 1.5-3m·s⁻¹) and standing (for 15s). The first maximum intensity section starts when a whistle is blown and the time is measured by electronic timing gates (PR1aW, Alge-Timing GmbH, Austria) with one-hundredth of a second accuracy. The scheme of the test is depicted in Figure 1.




Figure 1: Measured sections of SBFP₂₈.

◇ : Photocells; **→** : Backpedalling; - - - - - : Sprinting; **→** : Shuffling; ····· : Running; 1: Maximal Intensity Activity; 2: Medium Intensity Activity; 3: Low Intensity Activity

Statistical analysis

The distributions of the data sets were checked for homogeneity and skewness using the Kolmogorov-Smirnov test. Descriptive statistics (means, medians and standard deviations) were calculated for all variables. Wilcoxon nonparametric tests were used to examine whether the basketball specific fatigue protocol led to significant changes in the variables observed. Mann-

Whitney tests were used to compare the differences between the results of the U16 and U18 groups. The effect size was determined by the Cohen's *d* [36] and classified as small ($0.00 \leq d \leq 0.49$), medium ($0.50 \leq d \leq 0.79$), and large ($d \geq 0.80$). The significance level was set at $P < 0.05$. A statistical analysis was performed using the data analysis software system Statistica, version 12 (Stat Soft, Inc., Tulsa, USA).

Results

Table 1: Monitored parameters of pre and post basketball specific fatigue protocol SBFP₂₈ in the group of U16 basketball players.

Parameter	N	Pre Test M±SD	Post Test M±SD	% Change	Z	p
ABSLS	11	26.65±5.04	25.84±4.48	96.9	0.533	0.594
RELLS	11	32.63±7.28	31.35±4.97	96.1	0.622	0.534
RSI	11	0.49±0.08	0.54±0.13	110.7	2.49	0.013*

Table Abbreviations: N: Number of Players; M: Mean; SD: Standard Deviation; ABSLS: Absolute Leg Stiffness; RELLS: Relative Leg Stiffness; RSI: Reactive Strength Index; DL: Dominant Leg; NL: Non-Dominant Leg; H/Q: Hamstring Eccentric-to-Quadriceps Concentric Functional Ratio; 60: Angular Velocity of 60°/s; 180: Angular Velocity of 180°/s; Z: Wilcoxon Test Criterion; *: Significant Difference Between Pre and Post Test ($p < 0.05$)

Table 2: Monitored parameters of pre and post basketball specific fatigue protocol SBFP₂₈ in the group of U18 basketball players.

Parameter	N	Pre Test M±SD	Post Test M±SD	% change	Z	p
ABSLS	10	30.96±6.68	29.56±6.18	95.5	1.172	0.241
RELLS	10	35.29±6.42	33.79±6.58	95.8	1.07	0.284
RSI	10	0.41±0.07	0.41±0.07	100	0.05	0.959

Table Abbreviations: N: Number of Players; M: Mean; SD: Standard Deviation; ABSLS: Absolute Leg Stiffness; RELLS: Relative Leg Stiffness; RSI: Reactive Strength Index; DL: Dominant Leg; NL: Non-Dominant Leg; H/Q: Hamstring Eccentric-to-Quadriceps Concentric Functional Ratio; 60: Angular Velocity of 60°/s; 180: Angular Velocity of 180°/s; Z: Wilcoxon Test Criterion

The values of measured variables of the pre- and post- SBFP₂₈ and their changes for U16 and U18 groups of basketball players are reported in Table 1 & 2. No significant effects of SBFP₂₈ on the monitored parameters were evident with the exception of RSI

for U16 ($p = 0.013$, $r = 0.53$). The between-group differences in the changes of the monitored parameters are reported in Table 3. There were no significant differences in the fatigue-related responses to SBFP₂₈ for any parameters between the groups.

Table 3: Differences in changes of monitored parameters of pre and post basketball specific fatigue protocol SBFP₂₈ between the U16 and U18 groups.

Parameter	U16 M	Mdn	SD	U18 M	Mdn	SD	p
Δ ABSLS 1-2	0.81	0.73	3.92	1.46	0.7	4.65	0.972
Δ RELLS 1-2	1.29	0.71	5.03	1.55	0.74	5.4	0.86
Δ RSI 1-2	-0.05	-0.03	0.06	-0.03	-0.03	0.03	0.503

Table Abbreviations: N: Number of Players; M: Mean; SD: Standard Deviation; ABSLS: Absolute Leg Stiffness; RELLS: Relative Leg Stiffness; RSI: Reactive Strength Index; DL: Dominant Leg; NL: Non-Dominant Leg; H/Q: Hamstring Eccentric-to-Quadriceps Concentric Functional Ratio; 60: Angular Velocity of 60°/s; 180: Angular Velocity of 180°/s; Δ: Absolute Change

Discussion

The main finding of the current study is that there were no significant age differences in the fatigue-related responses to SBFP₂₈ for any of the parameters. Another finding is that SBFP₂₈, which was designed as a basketball-specific match simulation protocol, did not significantly compromise any of the observed parameters in any of the groups of youth basketball players. Oppositely, RSI values significantly increased in the U16 group. These results point out that after the basketball-specific fatigue protocol, the players' ability to effectively utilize neuromuscular mechanisms to control joint movement and reduce load on the ligaments was not compromised, which suggests that injury risk did not increase in the players of both groups due to simulated match-play. The results of this study did not show fatigue-related changes in ABSLS or RELLS in the U16 and U18 categories. These results indicate that the fatigue protocol did not compromise neuromuscular pre-activation and neuromuscular feed-forward control (anticipation of known stimuli characterized by preventive muscle activity) as an indicator of dynamic stability of the knee. Thus it can be assumed that the ability of the muscles to resist movement within the tibiofemoral joint (tibiofemoral shear movements), which prevents ACL from being under strain [21] did not decrease.

Studies examining the effects of simulated match-play on muscular and neuromuscular capability in youth basketball players are missing in the literature. Nevertheless, there are a few studies examining the effects of simulated match-play on muscular and neuromuscular capability in male youth soccer players and the results of these studies are ambiguous. In some cases leg stiffness was compromised [23,31]; some authors reported a non-significant decrease in ABSLS [24]. In the latter study, the authors also identified that altered stiffness due to simulated match play is individualised (in half of the pro-bands the value increased, in the other half the value decreased).

A comparison of the results of the above mentioned studies suggests that the differences may be influenced by the specifics of fatigue protocols for individual sports (but also within a particular sport), different levels of training of the monitored groups, gender differences, individual responses to the fatigue exercise, etc. Similarly to youth, the results of studies in adult athletes did not bring coherent results. While leg stiffness decreased after sport-specific fatigue protocol in soccer [22], highly individual changes were reported in runners [37]. Also in the case of reactive strength

evaluated by RSI, the results did not show a negative effect of SBFP₂₈. In the U16 group, the authors of the present study even recorded a significant improvement in RSI (p=0.013). The significance of the changes after completing SBFP₂₈ is also confirmed by the "effect size" value (r=0.53; large effect). The results suggest that the ability to change eccentric muscular action to concentric in a stretch-shortening cycle was not compromised in the players. This indicates that the players demonstrated a good ability of the muscle-tendon unit to cope with the stress induced by SBFP₂₈. Considering that jumping and landing are important aspects of basketball-specific fitness and that RSI was determined by means of a drop jump test, it could be expected, that these findings are also positive from the game performance perspective, specifically in terms of the quality of repeated jumping and landing actions. The findings of the current study correspond to the results of a recent study on elite adolescent soccer players (age 15.7±0.5 years), in which RSI increased significantly following a simulated soccer match play [31]. However, in a similar study on elite youth soccer players (age 14.4±0.5 years), RSI decreased significantly.

In the current study, the authors did not observe significant age differences in the fatigue-related responses to SBFP₂₈, either in leg stiffness (ABSLS; RELLS) or RSI. As mentioned above, these results cannot be compared with studies on youth basketball players. Nevertheless, in a later study on youth soccer players aged 13 to 16 years, the authors did not report any significant differences in fatigue-related responses to competitive match play for the parameters observed including ABSLS, RELLS, and RSI between the two age subgroups; 13.6±0.3 years and 15.8±0.1 years. Oppositely, a recent study on female youth soccer players indicated that leg stiffness decreased as a result of post simulated soccer in U13, but did not change in the U15, and increased in the U17 [38]. The data obtained in the present study also indicate that the values of both leg stiffness and RSI were higher in the U18 category compared with U16, in both pre-and post-test. This supports the suggestion that the ability to perform eccentric-concentric movements is positively influenced by age and that the older category has better dynamic stability of the knee joint and consequently a lower risk of ACL injury in sports activities [26,27]. The observed tendency also corresponds to the findings of a later study [25], in which the authors found out a significant increase in leg stiffness between 11 and 20 years of age in boys, with the highest increments between 17 and 20 years. One of the limitations of the study is that the indicators of neuromuscular fatigue were measured only after the

fatigue protocol. Another limitation is a low number of players who were involved in the study.

Conclusion

The SBFP₂₈ did not induce a significant reduction either in leg stiffness or reactive strength in the observed group of male adolescent basketball players. The results indicated that irrespective of age, the stabilization function of the knee muscles was not impaired and consequently the risk of ACL injury was not increased. However, a small number of players, and a relatively small age difference and other aspects of the methodological design of the study need to be considered. Due to the lack of similar studies on young basketball players and the results of a small number of previously conducted studies on youth in other sports, further research into the impact of the acute effect of match simulation protocol or competitive match-play on the mechanisms associated with the risk of ACL injury in youth basketball players is needed.

Acknowledgement

This study was supported by research grant from the Czech Science Foundation NO. 16-13750S.

References

- Hůlka K, Cuberek R, Bělka J (2013) Heart rate and time-motion analyses in top junior players during basketball matches. *Acta Univ Palacki Olomuc, Gymn* 43(3): 27-35.
- Apostolidis N, Nassis GP, Bolatoglou T, Geladas ND (2004) Physiological and technical characteristics of elite young basketball players. *J Sports Med Phys Fitness* 44(2): 157-163.
- Narazaki K, Berg K, Stergiou N, Chen B (2009) Physiological demands of competitive basketball. *Scand J Med Sci Sports* 19(3): 425-432.
- Schwebel DC, Brezaussek CM (2014) Child development and pediatric sport and recreational injuries by age. *J Athl Train* 49(6): 780-785.
- Alentorn-Geli E, Myer GD, Silvers HJ, Samitier G, Romero DR, et al. (2009) Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surg Sports Traumatol Arthrosc* 17(7): 705-729.
- Thacker SB, Stroup DF, Branche CM, Gilchrist J, Goodman RA, et al. (2003) Prevention of knee injuries in sports: a systematic review of the literature. *J Sports Med Phys Fitness* 43(2): 165-179.
- Cochrane JL, Lloyd DG, Buttfield A, Seward H, McGivern J (2007) Characteristics of anterior cruciate ligament injuries in Australian football. *J Sci Med Sport* 10(2): 96-104.
- Silvers HJ, Mandelbaum BR (2007) Prevention of anterior cruciate ligament injury in the female athlete. *Br J Sports Med* 41(1): 152-159.
- Parkkari J, Pasanen K, Mattila VM, Kannus P, Rimpelä A (2008) The risk for a cruciate ligament injury of the knee in adolescents and young adults: a population-based cohort study of 46 500 people with a 9 year follow-up. *Br J Sports Med* 42(6): 422-426.
- Shea KG, Pfeiffer R, Wang JH, Curtin M, Apel PJ (2004) Anterior cruciate ligament injury in pediatric and adolescent soccer players: An analysis of insurance data. *J Pediatr Orthop* 24(6): 623-628.
- Lohmander LS, Englund PM, Dahl LL, Roos EM (2007) The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *AM J Sports Med* 35(10): 1756-1769.
- Rumpf MC, Cronin J (2012) Injury incidence, body site, and severity in soccer players aged 6-18 years: Implications for injury prevention. *Stren Cond J* 34(1): 20-31.
- Van der Sluis A, Elferink-Gemser M, Coelho-e-Silva MJ, Nijboer JA, Brink MS, et al. (2014) Sport injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med* 35(4): 351-355.
- Dai B, Mao D, Garrett WE, Yu B (2014) Anterior cruciate ligament injuries in soccer: Loading mechanisms, risk factors, and prevention programs. *J Sport Health Sci* 3(4): 299-306.
- Sangnier S, Tourny-Chollet C (2007) Comparison of the decrease in strength between hamstrings and quadriceps during isokinetic fatigue testing in semiprofessional soccer players. *Int J Sports Med* 28(11): 952-957.
- Ekstrand J, Hägglund M, Waldén M (2009) Injury incidence and injury patterns in professional football: The UEFA injury study. *Br J Sports Med* 45(7): 553-558.
- Oliver J, Smith P (2010) Neural control of leg stiffness during hopping in boys and men. *J Electromyogr Kinesiol* 20(5): 973-979.
- Hawkins RD, Hulse MA, Wilkinson C, Hodson A, Gibson M (2001) The association football medical research programme: an audit of injuries in professional football. *Br J Sports Med* 35(1): 43-47.
- Rozzi SL, Lephart SM, Fu FH (1999) Effects of muscular fatigue on knee joint laxity and neuromuscular characteristics of male and female athletes. *J Athl Train* 34(2): 106-114.
- Smith HC, Vacek P, Johnson RJ, Slaughterbeck JR, Hashemi J, et al. (2012) Risk factors for anterior cruciate ligament injury: A review of the literature-Part 1: Neuromuscular and anatomic risk. *Sports Health* 4(1): 69-78.
- Hughes G, Watkins J (2006) A risk-factor model for anterior cruciate ligament injury. *Sports Med* 36(5): 411-428.
- Dutto DJ, Smith GA (2002) Changes in spring-mass characteristics during treadmill running to exhaustion. *Med Sci Sports Exerc* 34(8): 1324-1331.
- Lehnert M, De SteCroix M, Zaatar A (2017) Muscular and neuromuscular control following soccer-specific exercise in youth male footballers: Changes in injury risk mechanisms. *Scand J Med Sci Sports* 27(9): 975-982.
- Oliver JL, De SteCroix, MBA, Lloyd RS (2014) Altered neuromuscular control of leg stiffness following soccer-specific exercise. *Eur J Appl Physiol* 114(11): 2241-2249.
- Laffaye G, Choukou MA, Benguigui N, Padulo J (2017) Age and gender-related development of stretch-shortening cycle during a sub-maximal hopping task. *Biol Sport* 34(1): 29-35.
- Lloyd RS, Oliver JL, Hughes MG, Williams CA (2011) The influence of chronological age on periods of accelerated adaptation of stretch-shortening cycle performance in pre and postpubescent boys. *J Stren Cond Res* 25(7): 1889-1897.
- Lloyd R, Oliver J, Hughes M, Williams C (2012) Age-related differences in the neural regulation of stretch-shortening cycle activities in male youths during maximal and sub-maximal hopping. *J Electromyogr Kinesiol* 22(1): 37-43.
- Toumi H, Poumarat G, Best TM, Martin A, Fairclough J, et al. (2006) Fatigue and muscle-tendon stiffness after stretch-shortening cycle and isometric exercise. *Appl Physiol Nutr Metab* 31(5): 565-572.
- Young W (1995) Laboratory strength assessment of athletes. *New Stud Athl* 10(1): 88-96.

30. Raschner C, Platzer HP, Patterson C, Werner I, Huber R, et al. (2012) The relationship between ACL injuries and physical fitness in young competitive ski racers: a 10-year longitudinal study. *Br J Sports Med* 46(15): 1065-1071.
31. Lehnert M, De SteCroix M, Xaverova Z, Botek M, Varekova R, et al. (2018) Changes in injury risk mechanisms after soccer specific fatigue in male youth soccer players. *J Hum Kinet* 62: 1-10.
32. Dalleau G, Belli A, Viale F, Lacour JR, Bourdin M (2004) A simple method for field measurements of leg stiffness in hopping. *Int J Sports Med* 25(3): 170-176.
33. Flanagan EP, Comyns TM (2008) The use of contact time and the reactive strength index to optimize fast SSC training. *Stren Cond J* 30(5): 32-38.
34. Lloyd RS, Oliver JL, Hughes MG, Williams CA (2009) Reliability and validity of field-based measures of leg stiffness and reactive strength in youths. *J Sports Sci* 27(14): 1565-1573.
35. Hůlka K, Lehnert M, Bělka J (2017) Reliability and validity of a basketball-specific fatigue protocol simulating match load. *Acta Gymnica* 47(2): 92-98.
36. Hopkins WG, Marshall SW, Batterham AM, Hanin J (2009) Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc* 41(1): 3-12.
37. Hunter I, Smith GA (2007) Preferred and optimal stride frequency, stiffness and economy: changes with fatigue during a 1h high-intensity run. *Eur J Appl Physiol* 100(6): 653-661.
38. De Ste Croix MBA, Hughes JD, Lloyd RS, Oliver JL, Read PJ (2017) Leg stiffness in female soccer players: inter-session reliability and the fatiguing effects of soccer-specific exercise. *J Stren Cond Res* 31(11): 3052-3058.



Creative Commons Attribution 4.0
International License

For possible submissions Click Here

[Submit Article](#)



Research & Investigations in Sports Medicine

Benefits of Publishing with us

- High-level peer review and editorial services
- Freely accessible online immediately upon publication
- Authors retain the copyright to their work
- Licensing it under a Creative Commons license
- Visibility through different online platforms