



This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under All Rights Reserved license:

**Ayala, Francisco, López-Valenciano, Alejandro, Jose, Antonio, De Ste Croix, Mark B ORCID: 0000-0001-9911-4355, Vera-García, Francisco, García-Vaquero, Maria, Ruiz-Pérez, Iñaki and Myer, Gregory (2019) A preventive model for hamstring injuries in professional soccer: Learning algorithms. International Journal of Sports Medicine, 40 (5). pp. 344-353. doi:10.1055/a-0826-1955**

Official URL: <https://www.thieme-connect.com/products/ejournals/abstract/10.1055/a-0826-1955>

DOI: <http://dx.doi.org/10.1055/a-0826-1955>

EPrint URI: <http://eprints.glos.ac.uk/id/eprint/6383>

#### **Disclaimer**

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

**SDC 3:** Description of the dynamic postural control testing manoeuvre and measures obtained from it.

#### Dynamic postural control

Dynamic postural control was evaluated using the Y-Balance device® and following the guidelines described by Shaffer et al. [1]. The distance reached in each direction (anterior, posteromedial and posterolateral) was normalized by dividing by the previously measured leg length to standardize the maximum reach distance ([excursion distance/leg length] x100 = % maximum reach distance). The bilateral ratio (dominant / non-dominant score) of each direction was also calculated. Finally, to obtain a global measure of the balance test for each leg, data from each direction were averaged to calculate a composite score.

Measures obtained from the dynamic postural control test.

Name	Labels	
	Dominant Leg	Non-Dominant Leg
YBalance-Anterior	<57.825, 57.825-63.035 or >63.035	<58.515, 58.515- 63.51 or >63.51
YBalance-PosteroMedial	<101.215, 101.215- 107.865 or >107.865	<102.42, 102.42- 108.49 or >108.49
YBalance-PosteroLateral	<96.395, 96.395-104.93 or >104.93	<96.19, 96.19-103.71 or >103.71
BilaRatio-YBalance-Anterior	<0.965, 0.965-1.015 or >1.015	
BilaRatio-YBalance-PosteroMedial	<0.975, 0.975-1.005 or >1.005	
BilaRatio-YBalance-PosteroLateral	<0.985, 0.985-1.035 or >1.035	
YBalance-Composite	<85.44, 85.44-91.71 or >91.71	<86.73, 86.73-91.4 or >91.4-

Bila: bilateral

#### Reference

1. Shaffer SW, Teyhen DS, Lorenson CL, Warren RL, Koreerat, CM, Straseske CA, Childs JD. Y-balance test: a reliability study involving multiple raters. *Mil Med* 2013;178:1264-1270.