



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under All Rights Reserved license:

Ayala, Francisco ORCID: 0000-0003-2210-7389, López-Valenciano, Alejandro, Jose, Antonio, De Ste Croix, Mark B ORCID: 0000-0001-9911-4355, Vera-García, Francisco, García-Vaquero, Maria, Ruiz-Pérez, Iñaki and Myer, Gregory (2019) A preventive model for hamstring injuries in professional soccer: Learning algorithms. International Journal of Sports Medicine, 40 (5). pp. 344-353. doi:10.1055/a-0826-1955

Official URL: <https://doi.org/10.1055/a-0826-1955>

DOI: <http://dx.doi.org/10.1055/a-0826-1955>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/6383>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

SDC 4: Description of the isometric hip abduction and adduction strength testing manoeuvre and list of measures obtained from it.

Isometric hip abduction and adduction strength test

Isometric hip abduction and adduction peak torques of the dominant and non-dominant limb were assessed with a portable handheld dynamometer (Nicholas Manual Muscle Tester, Lafayette Indiana Instruments) in a supine lying position on a plinth with the participant's legs extended and following the methodology described by Thorborg et al. [1]. Briefly, participants performed five trials of 5-second isometric maximal voluntary contraction for each hip movement. The mean of the three most closely related trials were used for the subsequent statistical analyses. Unilateral hip abductor/adductor peak torque ratio defined as the hip adductor peak torque divided by hip abductor peak torque was calculated for each leg. Furthermore, the hip abduction and adduction bilateral ratios were also determined as the quotient of the dominant hip mean isometric peak value by the non-dominant hip mean isometric peak value. A side-to-side difference higher than 10% was defined as bilateral asymmetry.

Measures obtained from the isometric hip abduction and adduction strength test

Name	Labels	
	Dominant Leg	Non-Dominant Leg
PT _{ISOM} -HipAbd	<190.64, 190.64- 217.625 or >217.625	<194.025, 194.025-222 or >222
PT _{ISOM} -HipAbd-Normalised	<2.555, 2.555-2.91 or >2.91	<2.655, 2.655-2.92 or >2.92
PT _{ISOM} -HipAdd	<191.575, 191.575- 219.625 or >219.625	<187.75, 187.75-215.5 or >215.5
PT _{ISOM} -HipAdd-Normalised	<2.635, 2.635-2.965 or >2.965	<2.555, 2.555-2.905 or >2.905
UnRatio-ISOM-HipAbd/HipAdd	<0.956, 0.956-1.095 or >1.095	<0.92, 0.92-1.015 or >1.015
BilaRatio-PT _{ISOM} -HipAbd	No Asymmetry (<10%) or Asymmetry (≥10%)	
BilaRatio-PT _{ISOM} -HipAdd	No Asymmetry (<10%) or Asymmetry (≥10%)	

Bila: bilateral; Uni: unilateral; ISOM: isometric; PT: peak torque; Abd: abduction; Add: adduction.

Reference

1. Thorborg K, Petersen J, Magnusson SP, Hölmich P. Clinical assessment of hip strength using a hand-held dynamometer is reliable. *Scand J Med Sci Sports* 2010;20:493-501.