



This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under All Rights Reserved license:

Staddon, Philip L ORCID logoORCID: <https://orcid.org/0000-0002-7968-3179>, Reinsch, Sabine, Olsson, Pål A, Ambus, Per, Lüscher, Andreas, Jakobsen, Iver and Briones, Maria Jesus (2014) A decade of free-air CO₂ enrichment increased the carbon throughput in a grass-clover ecosystem but did not drastically change carbon allocation patterns. *Functional Ecology*, 28 (2). pp. 538-545. doi:10.1111/1365-2435.12183

Official URL: <http://dx.doi.org/10.1111/1365-2435.12183>

DOI: <http://dx.doi.org/10.1111/1365-2435.12183>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/6894>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

A decade of free-air CO₂ enrichment increased the carbon throughput in a grass-clover ecosystem but did not drastically change carbon allocation patterns

Philip L. Staddon^{†,1,2}, Sabine Reinsch^{*1}, Pål A. Olsson³, Per Ambus¹, Andreas Lüscher^{4,5} and Iver Jakobsen¹

¹Department of Chemical and Biochemical Engineering, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark; ²School of Biology, University of Nottingham, Nottingham NG7 2RD, UK; ³Biodiversity, Department of Biology, Lund University, Ecology Building, SE-223 62 Lund, Sweden; ⁴Institute of Agricultural Sciences, ETH Zurich, Universitätstrasse 2, 8092 Zurich, Switzerland; and ⁵Forage Production/Grassland Systems, Agroscope Reckenholz-Tänikon ART, Reckenholzstrasse 191, 8046 Zurich, Switzerland

Summary

1. The response of the soil carbon cycle to increasing atmospheric CO₂ concentration has far reaching consequences for the ecosystem carbon balance under future climatic conditions. We report on work carried out in the Swiss free-air CO₂ enrichment (FACE) experiment, where we used *in situ* ¹³CO₂ labelling to determine whether elevated CO₂ (+230 μL L⁻¹) concentration changes the fate of recently assimilated carbon in the soil microbial community.

2. Elevated CO₂ (eCO₂) concentration had an overall positive effect on microbial abundance ($P < 0.001$) with the gram-negative bacteria showing significantly increased quantities.

3. Gram-negative bacteria and saprotrophic fungi tended to utilize a higher amount of recently assimilated carbon under eCO₂. Arbuscular mycorrhizal fungi (AMF) utilized plant-assimilated carbon within 1 day after the ¹³CO₂ pulse and ¹³C uptake patterns in AMF suggest that carbon transfer is faster under eCO₂ concentration than under ambient CO₂ (aCO₂). Additionally, the respiration of recently assimilated carbon was significantly higher under eCO₂ than aCO₂ concentration.

4. Our data suggest that elevated atmospheric CO₂ concentration accelerated and increased the utilization of recently assimilated carbon by the microbial community without changing the microbial community composition drastically.

5. We conclude that a higher standing soil microbial biomass under eCO₂ concentration was the key cause for the higher carbon flow through the plant–soil system. Carbon utilization by microbial functional groups was only little affected by a decade of CO₂ enrichment.

Key-words: ¹³C labelling, arbuscular mycorrhizal fungi, C flux, compound-specific ¹³C analysis, elevated atmospheric CO₂, free-air CO₂ enrichment, neutral lipid fatty acids, phospholipid fatty acids, soil carbon

*Correspondence author. E-mail: sare@kt.dtu.dk

[†]Present address. ECEHH, University of Exeter Medical School, Truro, TR1 3HD, UK.

Introduction

Human-induced environmental change is a major societal concern (IPCC 2007). Predicting and mitigating the impacts of human activity on the Earth's climate is therefore receiving considerable attention. In particular, understanding how terrestrial ecosystems will respond to elevated atmospheric CO₂ (eCO₂) concentration is a priority research area in view of the existing feedbacks between the global carbon cycle and climate change (Luo *et al.* 1996; Carney *et al.* 2007; Selsted *et al.* 2012). The impact of eCO₂ concentration on carbon cycling has been investigated on small-scale mesocosms (Sowerby *et al.* 2000; Carney *et al.* 2007; Drigo *et al.* 2013), but also in large-scale field experiments (Van Kessel *et al.* 2000a; Deneff *et al.* 2007; Hill *et al.* 2007; Selsted *et al.* 2012). One of the first free-air CO₂ enrichment (FACE) experiments was the Swiss FACE that aimed to understand the impact of eCO₂ concentration in combination with nitrogen (N) fertilization and management practice on grass-clover vegetation using *Lolium perenne* L. and *Trifolium repens* L. (Hebeisen *et al.* 1997). It has been reported that after more than 8 years, photosynthesis was increased in both species (Ainsworth *et al.* 2003a,b), which is in accordance with responses in other grasses (Albert *et al.* 2011) and forbs (Ainsworth & Long 2005). Increased uptake of atmospheric CO₂ has frequently led to the hypothesis that the increased carbon flow into the ecosystem could increase soil carbon sinks, but results are ambiguous.

Extensive research in the Swiss FACE showed that increased photosynthesis did not necessarily lead to a higher carbon storage potential of the soil (Sowerby *et al.* 2000; Van Kessel *et al.* 2000a; Drissner *et al.* 2007) and this has also been reported by others (Fitter *et al.* 1996; Carney *et al.* 2007; Selsted *et al.* 2012). This result is not surprising because increased carbon availability in the ecosystem does not only affect the carbon accumulated in plant biomass (shoots and roots; Hebeisen *et al.* 1997; Suter *et al.* 2002) but also assimilated carbon released into the soil matrix, which stimulates microbial activity and maintains plant available nutrient levels in soil (Fontaine, Mariotti & Abbadie 2003; Paterson *et al.* 2007; Talbot, Allison & Treseder 2008). A stimulation of the microbial biomass results in carbon being bound in the microbial community, and at the same time, these microbes mineralize soil organic matter (SOM) and release carbon by respiratory processes (Carney *et al.* 2007).

In the Swiss FACE, the microbial community has been intensively studied (Hartwig *et al.* 2000; Lüscher *et al.* 2000; Staddon, Jakobsen & Blum 2004; Drissner *et al.* 2007). The microbial biomass was stimulated under eCO₂ concentration (Sowerby *et al.* 2000; Drissner *et al.* 2007), but increased microbial activity led to an enhanced soil CO₂ efflux counterbalancing the carbon input (Sowerby *et al.* 2000). Increased soil CO₂ efflux has also been observed for a temperate heathland where eCO₂ in combination with warming and periodic summer drought induced soil carbon loss by 21% (Selsted *et al.* 2012).

One major microbial group associated with clover is the nitrogen-fixing gram-negative rhizobia bacteria that were found to balance soil nutrient levels under eCO₂ concentration in the Swiss FACE (Lüscher *et al.* 2000; Hartwig *et al.* 2002). In the grass-clover mixture, 59% of the nitrogen was supplied from atmospheric nitrogen fixation under ambient CO₂ (aCO₂) concentration, and this amount was increased by 12% when plants were exposed to 600 ppm CO₂ (Zanetti *et al.* 1996).

Fungi were another important microbial group in the Swiss FACE: fungal enzyme production was increased under eCO₂ concentration and fungal abundance rose under low nitrogen fertilization (Drissner *et al.* 2007). Fungi have been shown to be a relevant driver of soil CO₂ efflux under eCO₂ concentrations (Carney *et al.* 2007) due to their ability to increase nutrient availability in the soil by mineralizing SOM (Carney *et al.* 2007; Drissner *et al.* 2007). Another important microbial group in terms of nutrient balance of an ecosystem is the group of mycorrhizal fungi. Mycorrhizal fungi provide nutrients, especially phosphorus, to plants in exchange for plant carbon. This is especially important in the

rhizosphere where phosphorous can be depleted and mycorrhizal hyphae can reach beyond the zone of nutrient depletion. In the Swiss FACE, arbuscular mycorrhizal fungi (AMF) were positively affected by eCO₂ conditions combined with low nitrogen fertilization (Staddon, Jakobsen & Blum 2004).

Here, we used an *in situ* ¹³C₂ pulse labelling approach (Ostle *et al.* 2000; Leake *et al.* 2006) to track and quantify the carbon flow through the ecosystem under ambient and elevated atmospheric CO₂ concentrations. We measured the ¹³C enrichment of microbial fatty acid biomarkers to determine the effect of eCO₂ on the relative activity of major soil microbial functional groups. In particular, the ¹³C label content of phospholipid fatty acids (PLFA) and neutral lipid fatty acids (NLFA) can be used to compare the relative activity of different groups of soil microorganisms, for example, gram-negative and gram-positive bacteria, saprotrophic and AMF (Treonis *et al.* 2004; Olsson & Johnson 2005). Elevated CO₂ is known to affect soil microbial communities, principally via altered carbon input quality, and clearly affects the relative activity of different soil microorganisms (Montealegre *et al.* 2002). We hypothesize that elevated CO₂ concentration changes the fate of carbon in the soil microbial community towards a higher carbon recovery in gram-negative bacteria, saprotrophic fungi and AMF.

Materials and methods

THE SWISS FACE EXPERIMENT

The research was carried out at the Swiss FACE site, Lindau, Switzerland (Hebeisen *et al.* 1997), where elevated atmospheric CO₂ concentration (+230 μL L⁻¹) was imposed for 10 years from 1993. The lowland grassland site and treatments are fully described elsewhere (Schortemeyer *et al.* 1996). There were three replicate FACE rings and three replicate control (ambient) rings yielding a total of six labelling chambers. The labelling work reported here was entirely carried out in the mixed *L. perenne* L. and *T. repens* plots under low N addition (14 g m⁻² year⁻¹).

PULSE LABELLING WITH ¹³ CO₂

The ¹³C label was delivered to the vegetation as ¹³CO₂ at 99 atom% ¹³C (Staddon *et al.* 2003) at two CO₂ concentrations, that is, ambient (aCO₂, 370 μL L⁻¹) and elevated (eCO₂, 600 μL L⁻¹). This was achieved using an inexpensive and highly flexible method developed for labelling in the field, which, importantly, allowed the delivery of the ¹³C label simultaneously at different CO₂ concentrations. Briefly, this method involved six individual labelling units consisting of 250 L capacity gas impermeable Tedlar bags, which were filled with air containing ¹³CO₂ at 370 or 600 μL L⁻¹. The 99 atom% ¹³CO₂ was mixed into bottled CO₂-free air to achieve the required concentration. The labelled air was then pumped at a flowmeter controlled rate of 1.0 L min⁻¹ to purpose built 20 cm diameter, 15 cm high, transparent acrylic labelling chambers, placed flush with the soil surface. The labelling areas were selected randomly within the low N mixed vegetation plots. Exhaust air from the labelling chambers was expelled 10 m outside the FACE and control rings. Labelling was carried out on 13 September 2002 in the afternoon and lasted for 3-5 h under sunny conditions.

We tested whether ¹³C from the ¹³CO₂ labelling pulse entered the soil directly in significant quantity by applying a ¹³CO₂ pulse to vegetation that was kept dark (labelling chambers covered in opaque plastic; Staddon *et al.* 2003). Labelling in darkened chambers was carried out outside the FACE rings. Gas samples in unlabeled area were taken at least 2 m away from the labelled area. Soil-respired CO₂ was collected in the same manner as for the actual labelling experiment (detailed below). We found that the ¹³C content of soil-respired CO₂ in unlabeled area and the labelled darkened area was similar at the relevant times after labelling.

GAS, PLANT AND SOIL SAMPLE COLLECTION

Gas sampling was carried out to investigate the ^{13}C content in soil CO_2 efflux 2 h and 1, 2 and 4 days after $^{13}\text{CO}_2$ labelling. Samples were taken within the labelled area and at 5 and 15 cm away from the outer edge of the labelled area. Soil-respired CO_2 was collected using one 10 mL Exetainer vial per sampling point (Labco Ltd., High Wycombe, UK) that was placed upside down on the soil surface (Staddon *et al.* 2003). Vials were flushed with ambient air before the incubation so that, within treatments, they all had the same starting point. Care was taken not to include shoot respiration within the trapped soil-respired CO_2 . The tubes were left in place for 4 h, after which, each tube was carefully lifted from the soil surface and capped. The soil-respired CO_2 samples were stored at ambient temperature.

Samples of above-ground biomass were taken in all labelled areas to investigate the ^{13}C content 2 h, 1 and 2 days after $^{13}\text{CO}_2$ labelling. Samples were taken within the labelled area and were obtained as close as possible to where the Exetainer vials were located.

Soil cores (2 cm diameter) were taken in all labelled areas to examine the fate of recently assimilated carbon (^{13}C) into the soil microbial community and to assess the microbial biomass 2 h and 1, 2, 3, 4, 6 and 14 days after the $^{13}\text{CO}_2$ pulse. Cores were taken to a depth of 10 cm at the same location as the gas samples. Above-ground biomass samples and soil cores were frozen at $-18\text{ }^\circ\text{C}$ until analysis.

CARBON-13 ANALYSES OF GAS AND PLANT SAMPLES

The ^{13}C content of CO_2 (soil respiration) was analysed in subsamples transferred to evacuated 2 mL crimp seal vials for immediate processing. The vial headspace was purged into the He carrier stream of a PreCon trace gas preparation-concentration unit (Thermo Electron, Bremen, Germany) interfaced with a HP 6890 gas chromatograph coupled in continuous flow mode to a Finnigan MAT Delta PLUS isotope ratio mass spectrometer (Thermo Electron). As working standard, we used commercial CO_2 calibrated against certified $^{13}\text{CO}_2$ standards (Messer Griesheim, Krefeld, Germany).

Total carbon and the $^{13}\text{C}/^{12}\text{C}$ ratio in plant material was measured by Dumas combustion on an elemental analyser (CE 1110; Thermo Electron, Milan, Italy) coupled in continuous flow mode to the Delta PLUS. Analyses were performed on dried, finely ground 2–3 mg samples weighed out into tin combustion capsules. As working standard for ^{13}C determinations, commercial CO_2 calibrated against certified ^{13}C -sucrose material (IAEA, Vienna, Austria) was used.

EXTRACTION, IDENTIFICATION AND ^{13}C ANALYSES OF FATTY ACIDS

The extracted PLFAs were selected as being specific to various groups of soil micro-organisms: general biota [16:0, 18:0 (both PLFAs synthesized by microbes and plants)], gram-negative bacteria (cy17:0, cy19:0, 18:1x7), gram-positive bacteria (i15:0, a15:0, i16:0, i17:0, a17:0, 10Me17), actinomycetes (10Me16, 10Me18) and saprotrophic fungi (18:1x9, 18:2x6,9). The NLFA 16:1x5 as specific to AMF was chosen as it has been shown to be a more sensitive indicator of AMF hyphae in soil than the PLFA 16:1x5 (Olsson 1999).

Lipids within soil (roots removed) were extracted in a one-phase mixture of citrate buffer, methanol and chloroform (0-8:2:1, v/v/v, pH 4-0). The lipids were fractionated into neutral lipids, glycolipids and phospholipids on pre-packed silica columns (100 mg sorbent mass, Varian Medical Systems, Palo Alto, CA, USA) as described by van Aarle & Olsson (2003). The fatty acid residues in neutral lipids and phospholipids were transformed into free fatty acid methyl esters and identified and quantified by gas

chromatography.

The ^{13}C enrichment in fatty acid methyl esters was determined in a 20-20 IRMS (PDZ Europa Scientific Instruments, Crewe, UK) interfaced with a Hewlett Packard 6890 gas chromatograph (Hewlett Packard, Palo Alto, CA, USA). The chromatographic conditions were as described by Olsson *et al.* (2005). The ^{13}C values were calculated based on atom% ^{13}C of the reference CO_2 gas, injected three times at the beginning and end of a chromatographic run. The reference CO_2 was standardized with the Pee Dee Belemnite (PDB) standard. The precision of the reference gas ^{13}C was 0.2‰. Integration for each peak was checked and corrected manually.

Carbon isotopic enrichment was highest in above-ground vegetation and was expressed as atom% ^{13}C excess (APE) and calculated as follows:

$$\text{APE}(\%) = {}^{13}\text{C content}_{\text{sample}} - {}^{13}\text{C content}_{\text{control}} \quad \text{Eqn 1}$$

where the $^{13}\text{C content}_{\text{sample}}$ ground vegetation after the is the measured ^{13}C content in above- $^{13}\text{CO}_2$ pulse, and $^{13}\text{C content}_{\text{control}}$ is the ^{13}C content in above-ground vegetation before the experiment.

Measured ^{13}C enrichments (^{13}C) Δ fatty acids and soil respiration are calculated as:

$$\Delta^{13}\text{C}(\text{‰}) = \delta^{13}\text{C}_{\text{sample}} - \delta^{13}\text{C}_{\text{control}} \quad \text{Eqn 2}$$

where $\delta^{13}\text{C}_{\text{sample}}$ is the $^{13}\text{C}/^{12}\text{C}$ ratio of individual PLFAs and soil respiration after the $^{13}\text{CO}_2$ pulse and $\delta^{13}\text{C}_{\text{control}}$ before the $^{13}\text{CO}_2$ pulse.

DATA ANALYSES

Statistical analyses were performed using R version 2.12.1 (www.R-project.org). All data were checked for normality and where necessary log-transformed. Analysis of variance (ANOVA) was performed to test for effects of CO_2 concentration on ^{13}C content in above-ground plant biomass. Sampling time was included as fixed variable to test for the effect of time.

Fatty acid quantities are presented as averages of sampling times and replicates (FACE rings) because fatty acid abundances are not assumed to change over the investigated time period. Standard errors were calculated using error propagation. The overall effect of eCO_2 on fatty acid abundances was tested using a MANOVA , and the effect of eCO_2 on individual fatty acids was analysed with a linear mixed-effect model (lme) using 'FACE ring' as a random variable because fatty acid quantities were consistently lower in one of the FACE rings.

The effect of eCO_2 on $\Delta^{13}\text{C}$ in microbial functional groups (averages of ^{13}C values of functional group specific fatty acids) was analysed using the 'lme' function with 'FACE ring/time' as random variable. The model was extended for a 'varIdent' structure that accounts for inhomogeneous variances if the data did not show homogeneous variances. The same model was used to test for the effect of CO_2 concentration on soil-respired $\Delta^{13}\text{CO}_2$.

Results

Plant ^{13}C content decreased with time but plant ^{13}C enrichment was not different between CO_2 treatments (Fig. 1). Microbial fatty acid abundances were significantly higher under eCO_2 compared to aCO_2 concentration (ANOVA : $P < 0.001$). This CO_2 -induced increase was lowest for actinomycetes ($9 \pm 4\%$) and highest for AMF (40%; Fig. 2). Gram-negative-specific cy17:0 and cy19:0 PLFAs were significantly more abundant in the eCO_2 treatment ($P < 0.05$). The same trend was observed for the gram-positive-specific biomarkers i15:0 ($P = 0.098$) and i16:0 ($P = 0.065$) and the actinomycetes PLFA 10Me16:0 ($P = 0.057$). The AMF-specific NLFA 16:1x5 also tended to be more abundant under eCO_2 ($P = 0.061$).

The $\Delta^{13}\text{C}$ in fatty acids was affected by functional group ($P = 0.046$) and CO_2 concentration ($P = 0$ -

025) when considering all time points. There was no significant effect of eCO₂ on $\Delta^{13}\text{C}$ in microbial functional groups. However, eCO₂ had a tendency to increase the $\Delta^{13}\text{C}$ in gram-negative bacteria and saprotrophic fungi, whereas AMF were more enriched in ¹³C when exposed to aCO₂ concentration (Fig. 3). The $\Delta^{13}\text{C}$ in gram-positive bacteria ($P = 0.40$) and actinomycetes ($P = 0.50$) were unaffected by eCO₂ (Fig. S1, Supporting information).

There was a substantial difference in $\Delta^{13}\text{C}$ of soil-respired CO₂ between aCO₂ and eCO₂ within the labelling plots (Fig. 4). The ¹³C enrichment of soil-respired CO₂ was considerably higher at eCO₂ than aCO₂. At 5 cm distance from the labelled plots, the ¹³CO₂ efflux pattern was similar at aCO₂ and eCO₂ over the 4-day sampling period (data not shown). $\Delta^{13}\text{C}$ was on average 4‰ in aCO₂ and was increased by ~2-5‰ under eCO₂ concentration. At 15 cm distance from the labelled plots, aCO₂ and eCO₂ plots showed a ~5‰ $\Delta^{13}\text{C}$ in soil CO₂ efflux. In aCO₂ plots, the measured $\Delta^{13}\text{C}$ increased to ~8‰ and approached zero thereafter. In eCO₂ plots, measured $\Delta^{13}\text{C}$ approached zero already 1 day after the labelling pulse.




Fig. 1. Above-ground ¹³C enrichment at ambient and elevated CO₂ concentration over time. ¹³C enrichment is expressed as ¹³C atom% excess. Effect of the CO₂ treatment was statistically analysed with an analysis of variance (ANOVA) and is given in the upper right corner. Enrichments are presented as means \pm SE.

Discussion

PLANT CARBON UPTAKE

Our results show the same ¹³C enrichment of above-ground biomass under aCO₂ and eCO₂ concentrations (Fig. 1). However, we can assume that the uptake of ¹³CO₂ was higher under eCO₂ conditions because we know from earlier studies at the Swiss FACE that eCO₂ concentration affects the photosynthetic capacity of plants and results in increased carbon uptake (Ainsworth *et al.* 2003a, b). Seen in this context, our results suggest that photo-assimilated carbon is not stored in above-ground biomass but transported below-ground under the assumption that plant biomass is similar or higher under eCO₂ compared to aCO₂ (Ainsworth & Long 2005; Kongstad *et al.* 2012). This is a reasonable assumption because the labelling pulse was deployed in late summer when temperate plants increase the allocation of carbon to below-ground parts to prepare for the winter season. Increased fixation of CO₂ and a more rapid transfer of photo-assimilates below-ground under eCO₂ concentration (Hebeisen *et al.* 1997; Daepf, Nösberger & Lüscher 2001) create a bigger carbon pool for plant and below-ground activity but, at the same time, increase the nutrient demand for plants and microbes.

MICROBIAL COMMUNITY RESPONSE TO ELEVATED CO₂

Elevated CO₂ significantly increased the soil microbial PLFA abundance by ~16% compared to aCO₂ concentration (Fig. 2). This finding is in accordance with previous results from the same experimental site where total PLFA abundances were increased by about 25% (Drissner *et al.* 2007). Microbial PLFA abundance and thus biomass (Frostegård & Bååth 1996) likely increased due to the higher carbon input to the soil that stimulated microbial growth and activity due to, for example, rhizodeposition.

In particular, the abundance of gram-negative bacteria was significantly increased under eCO₂ concentration. Gram-negative bacteria are fast growing and highly dependent on plant photo-assimilates (Paterson *et al.* 2007; Jin & Evans 2010; Garcia-Pausas & Paterson 2011) and the increase of root growth (Jongen *et al.* 1995; Arndal *et al.* 2013) and connected rhizodeposition under eCO₂ can therefore have a facilitating effect on gram-negative bacteria.

In soil, the free-living gram-negative bacteria *Azotobacter spp.* (Strandberg & Wilson 1968) and the symbiotic (gram-negative) rhizobia are able to fix atmospheric nitrogen, and thus, they are a key component in the nitrogen balance of the soil. However, in the present study, it was not possible to




Fig. 2. Quantities of microbial phospholipid fatty acid (nmol g⁻¹ dry soil) and the neutral lipid fatty acid 16:1x5 under ambient and elevated CO₂ concentrations. The effect of the eCO₂ treatment is given in the upper right corner. Bars represent means ± SE. *indicates significant differences on a level of $P \leq 0.05$.

determine whether these specific groups were part of the general increase in gram-negative bacteria biomass. In general, in other studies from the Swiss FACE, nodule formation was increased by 17–50% due to the symbiosis between plants and rhizobia under eCO₂ concentration (Schortemeyer *et al.* 1996; Montealegre *et al.* 2000). Nitrogen-fixing bacteria counterbalanced the expected negative effect of decreased above-ground biomass and lower nitrogen yield and maintained a similar C/N balance under aCO₂ and eCO₂ concentrations (Hartwig *et al.* 2000; Lüscher *et al.* 2000). Additionally, the increased nitrogen demand arising from increased carbon input under eCO₂ concentration was entirely supplied by nitrogen-fixing symbionts in the grass-clover mixture (Zanetti *et al.* 1996). However, the utilization of recently assimilated carbon by free-living nitrogen-fixing bacteria should be investigated in greater detail to confirm their dependence on rhizodeposits.

The AMF biomarker (NLFA 16:1x5) tended to be positively affected by eCO₂ concentration. Even

though this result was not significant, this trend is not a random effect because the extraradical mycorrhizal hyphae density was increased from 0.3 to 0.8 m g⁻¹ soil in aCO₂ to 1.0– 1.7 m g⁻¹ soil in eCO₂ by October 2002, (Staddon, Jakobsen & Blum 2004) that is, 1 month after the pulse labelling. AMF are known to act as a carbon sink (Jakobsen & Rosendahl 1990; Olsson & Johnson 2005), and due to their symbiotic connection with the host plant, AMF are the first group to receive recently plant-assimilated carbon (Olsson & Johnson 2005).

CHANGED MICROBIAL ACTIVITY

Arbuscular mycorrhizal fungi displayed distinct ¹³C enrichment in the biomarker NLFA 16:1x5 under labelling at the two CO₂ concentrations (Fig. 3c). Under aCO₂ concentration, uptake of recently assimilated CO₂ peaked at 80‰ d¹³C between 2 and 3 days after labelling and decreased thereafter. In contrast, under eCO₂ concentration, the maximum d¹³C was measured to be ~44‰ 1 day after labelling. However, the lower D¹³C in the eCO₂ treatment is a result of pool dilution because extraradial mycorrhizal hyphae density was ~50% increased (Staddon, Jakobsen & Blum 2004) in soils with eCO₂ concentration. The measured isotopic value must therefore be diluted by 50% when rhizodeposition was similar in eCO₂ and aCO₂ treatments. When considering the isotopic dilution effect, the level of NLFA would be similar.

However, carbon transport has been shown to be faster under eCO₂ than under aCO₂ concentration (Selsted *et al.* 2012). Furthermore, the symbiosis between AMF and plant roots facilitate a very fast transfer of recently assimilated carbon between plants and AMF (Moyano, Kutsch & Schulze 2007) in exchange for nutrients (Jakobsen & Rosendahl 1990; Johnson, Leake & Read 2002; Olsson & Johnson 2005). In a ¹⁴C labelling experiment, ¹⁴C-labelled carbon that was taken up by plants was re-allocated below-ground and peaked in associated AMF within 16 h after the labelling pulse (Jakobsen & Rosendahl 1990). Our second sampling took place 1 day after the labelling implying that our measuring window was too wide to detect the early peak of enrichment in the eCO₂ treatment under fast carbon transfer conditions. Under aCO₂ concentration, carbon transfer is slower than under eCO₂ because the labelling peak seems to occur between 1 and 2 days after the labelling pulse. Thus, the symbiotic lifestyle of AMF results in a rapid transfer of carbon from the plant, and this carbon transfer is faster under eCO₂ than under aCO₂ concentrations. A more detailed evaluation of the impact of AMF on the net carbon balance of the system could be achieved with more frequent samplings.

Fungi did not take up much of the recent plant-assimilated carbon under aCO₂ concentration (Fig. 3b). In contrast, fungi grown under eCO₂ concentration showed a dynamic ¹³C uptake. At the Swiss FACE experiment, it has been reported that fungal and gram-negative-specific PLFA abundance were reduced under high nitrogen fertilization compared to the low fertilization treatment (Drissner *et al.* 2007). Thus, under low nitrogen fertilization, the abundance of fungi and gram-negative bacteria is important for, for example, maintaining the nutrient balance under induced nutrient stress in the eCO₂ treatment. In addition, soil enzyme activities were increased by 17–42% under eCO₂ concentration in October 2002 (Drissner *et al.* 2007), suggesting a crucial role of the fungal group in the maintenance of available nutrients.

Gram-negative bacteria under eCO₂ concentration received recently assimilated ¹³C after a time delay of about 3 days (Fig. 3a). The observed peak enrichment at day three fits to the observation that carbon is first utilized by AMF and becomes available from rhizodeposits for bacteria and other soil organisms later (Olsson & Johnson 2005). The significant increase in gram-negative biomarker quantities coupled with a higher Δ¹³C indicates a high carbon demand in that microbial group which could be related to the effect of eCO₂ on rhizodeposition and the facilitation of gram-negative bacteria or to nitrogen-fixing

bacteria that have been shown to be abundant in this ecosystem (Lüscher *et al.* 2000; Drissner *et al.* 2007).




Fig. 3. $\Delta^{13}\text{C}$ (%) in microbial functional groups at ambient and elevated CO_2 concentrations over time: (a) gram-negative bacteria (average of cy17:0, cy19:0 and 18:1 ω 7), (b) saprophytic fungi (average of 18:1 ω 9 and 18:2 ω 9) and (c) arbuscular mycorrhizal fungi neutral lipid fatty acids (NLFA 16:1x5). Effects of the CO_2 treatment are given in the upper right corners. Data are presented as means \pm SE.




Fig. 4. Soil-respired $\Delta^{13}\text{C}$ (&) at ambient and elevated CO_2 concentration. Effect of the CO_2 treatment is given in the upper right corner. Data are represented as means \pm SE.

SOIL CO₂ EFFLUX

Soil CO₂ efflux showed a significantly higher $\Delta^{13}\text{C}$ under eCO₂ concentration (Fig. 4). To evaluate the actual amount of respired ¹³C under aCO₂ and eCO₂ concentrations, the isotopic values need to be weighted with the soil CO₂ efflux. However, soil CO₂ efflux was not assessed in this experiment, but it has been convincingly shown that eCO₂ concentration increases soil CO₂ efflux in grasslands (Luo *et al.* 1996; Selsted *et al.* 2012) and other ecosystems (Carney *et al.* 2007; Hyvönen *et al.* 2007). Therefore, the increased $\Delta^{13}\text{C}$ of soil-respired CO₂ under eCO₂ conditions is probably also associated with a higher soil CO₂ efflux and the respiration of recently assimilated ¹³C would therefore be higher under eCO₂ concentration.

Soil respiration is mainly composed of root and microbial respiration. Increased root biomass in *L. perenne* (Hebeisen *et al.* 1997; Van Kessel *et al.* 2000a; Daepf, Nösberger & Lüscher 2001) under eCO₂ concentration is a potential source of recently assimilated ¹³C to the soil CO₂ efflux. Furthermore, increased symbiosis of N-fixing bacteria with *T. repens* (Schortemeyer *et al.* 1996; Zanetti *et al.* 1996; Montealegre *et al.* 2000) is another potential source of ¹³CO₂ production. Also, a higher fungal importance under low nitrogen fertilization (Drissner *et al.* 2007) and an increased utilization of ¹³C compared at aCO₂ can add a substantial amount of ¹³C to the soil CO₂ efflux. CO₂ originating from these sources together can result in the observed higher $\Delta^{13}\text{C}$ in soil respiration.

The main contribution of AMF to the measured $\Delta^{13}\text{C}$ in soil CO₂ efflux under eCO₂ probably appeared within 16 h after the pulse labelling (Jakobsen & Rosendahl 1990) and was not captured. This assumption is in agreement with the finding that mycorrhizal fungi are strongly dependent on photosynthetic activity (Moyano, Kutsch & Schulze 2007) and a faster carbon turnover under eCO₂ concentration would therefore result in an earlier ¹³C enrichment peak. This is supported by the observed lateral ¹³C transport in 15 cm distance from the labelled plots where the ¹³C enrichment peak in soil respiration occurred 1 day after labelling. This peak was delayed in the aCO₂ compared to eCO₂ treatment and can be explained by carbon transport within the mycorrhizal network that can cover relatively large distances (Johnson, Leake & Read 2002).

SOIL CARBON STORAGE POTENTIAL

Plant-assimilated carbon can be distributed to different carbon pools, but will eventually be released back to the atmosphere. The increased carbon input to the soil under eCO₂ led to increased soil microbial biomass (Sowerby *et al.* 2000; Drissner *et al.* 2007) but the soil-respired CO₂ per unit microbial biomass stayed unchanged (Sowerby *et al.* 2000) likely due to higher rates of SOM mineralization (Drissner *et al.* 2007). After 4 years of eCO₂ fumigation in the Swiss FACE, the soil carbon pool remained unchanged under eCO₂ concentration as shown in previous studies (Van Kessel *et al.* 2000a,b). Our observations agree with those studies that reported a stimulated microbial community under eCO₂ and high impact of the fungal group in the studied ecosystem. Our results suggest that eCO₂ concentration increases the carbon throughput of the ecosystem (Figs 3 and 4) due to a higher soil carbon input and an increased soil microbial biomass. A higher carbon throughput but unchanged carbon allocation patterns under aCO₂ and eCO₂ concentrations point towards an unchanged net ecosystem carbon balance which is in agreement with previous observations (Van Kessel *et al.* 2000a,b).

Conclusions

We presented the fate of carbon into above-ground and below-ground carbon pools in mixed grass-clover vegetation in a ¹³CO₂ pulse labelling experiment after a decade of CO₂ fumigation. In this agricultural

relevant vegetation mixture, gram-negative bacteria and fungi have been shown to be important for the nutrient balance under eCO₂ especially. Our data suggest that the increase in microbial biomass is counterbalancing the increased plant nutrient demand under eCO₂ concentration. Furthermore, the increase in below-ground carbon allocation is balanced by a higher microbial activity, and thus, the release of CO₂ back to the atmosphere is increased. We conclude that, due to these balancing mechanisms, a decade of eCO₂ concentration did not drastically change carbon allocation in the studied grass-clover ecosystem.

Acknowledgements

We thank the European Commission for awarding P. L. Staddon a Marie Curie Individual Fellowship. P. A. Olsson thanks the Swedish Research Council for financial support. We thank all the staff involved with the Swiss FACE experiment at ETH Zürich for making this work possible, especially I. Popa for help with sample collection. We also thank N. Ostle at CEH Merlewood for providing the labelling chambers and A. H. Fitter, J. P. Grime and K. Thompson for comments on an earlier version of this paper.

References

- van Aarle, I.M. & Olsson, P.A. (2003) Fungal lipid accumulation and development of mycelial structures in two arbuscular mycorrhizal fungi. *Applied and Environmental Microbiology*, 69, 6762–6767.
- Ainsworth, E.A. & Long, S.P. (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist*, 165, 351–372.
- Ainsworth, E.A., Davey, P.A., Hymus, G.J., Osborne, C.P., Rogers, A., Blum, H. et al. (2003a) Is stimulation of leaf photosynthesis by elevated carbon dioxide concentration maintained in the long term? A test with *Lolium perenne* grown for 10 years at two nitrogen fertilization levels under Free Air CO₂ Enrichment (FACE). *Plant, Cell and Environment*, 26, 705–714.
- Ainsworth, E.A., Rogers, A., Blum, H., Nösberger, J. & Long, S.P. (2003b) Variation in acclimation of photosynthesis in *Trifolium repens* after eight years of exposure to Free Air CO₂ Enrichment (FACE). *Journal of Experimental Botany*, 54, 2769–2774.
- Albert, K.R., Mikkelsen, T.N., Michelsen, A., Ro-Poulsen, H. & Van der Linden, L. (2011) Interactive effects of drought, elevated CO₂ and warming on photosynthetic capacity and photosystem performance in temperate heath plants. *Journal of Plant Physiology*, 168, 1550–1561.
- Arndal, M.F., Merrild, M.P., Michelsen, A., Schmidt, I.K., Mikkelsen, T.N. & Beier, C. (2013) Net root growth and nutrient acquisition in response to predicted climate change in two contrasting heathland species. *Plant and Soil*, 369, 615–629.
- Carney, K.M., Hungate, B.A., Drake, B.G. & Megonigal, J.P. (2007) Altered soil microbial community at elevated CO₂ leads to loss of soil carbon. *Proceedings of the National Academy of Sciences United States of America*, 104, 4990–4995.
- Daepf, M., Nösberger, J. & Lüscher A. (2001) Nitrogen fertilization and developmental stage alter the response of *Lolium perenne* to elevated CO₂. *New Phytologist*, 150, 347–358.
- Denef, K., Bubenheim, H., Lenhart, K., Vermeulen, J., Van Cleemput, O. & Boeckx, P. (2007) Community shifts and carbon translocation within metabolically-active rhizosphere microorganisms in grasslands under elevated CO₂. *Biogeosciences*, 4, 769–779.
- Drigo, B., Kowalchuk, G.A., Knapp, B.A., Pijl, A.S., Boschker, H.T.S. & Van Veen, J.A. (2013) Impacts of 3 years of elevated atmospheric CO₂ on rhizosphere carbon flow and microbial community dynamics. *Global Change Biology*, 19, 621–636.
- Drissner, D., Blum, H., Tscherko, D. & Kandeler, E. (2007) Nine years of enriched CO₂ changes the function and structural diversity of soil micro-organisms in a grassland. *European Journal of Soil Science*, 58, 260–269.
- Fitter, A.H., Self, G.K., Wolfenden, J., Van Vuuren, M.M.I., Brown, T.K., Williamson, L. et al. (1996) Root production and mortality under elevated atmospheric carbon dioxide. *Plant and Soil*, 187, 299–306.

- Fontaine, S., Mariotti, A. & Abbadie, L. (2003) The priming effect of organic matter: a question of microbial competition? *Soil Biology and Biochemistry*, 35, 837–843.
- Frostegård, A. & Bååth, E. (1996) The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. *Biology and Fertility of Soils*, 22, 59–65.
- Garcia-Pausas, J. & Paterson, E. (2011) Microbial community abundance and structure are determinants of soil organic matter mineralisation in the presence of labile carbon. *Soil Biology and Biochemistry*, 43, 1705–1713.
- Hartwig, U.A., Lüscher, A., Daupp, M., Blum, H., Soussana, J.-F. & Nösberger, J. (2000) Due to symbiotic N₂ fixation, five years of elevated atmospheric pCO₂ had no effect on the N concentration of plant litter in fertile, mixed grassland. *Plant and Soil*, 224, 43–50.
- Hartwig, U.A., Lüscher, A., Nösberger, J. & Van Kessel, C. (2002) Nitrogen-15 budget in model ecosystems of white clover and perennial rye-grass exposed for four years at elevated atmospheric pCO₂. *Global Change Biology*, 8, 194–202.
- Hebeisen, T., Lüscher, A., Zanetti, S., Fischer, B.U., Hartwig, U.A., Frehner, M. et al. (1997) Growth response of *Trifolium repens* L. and *Lolium perenne* L. as monocultures and bi-species mixture to free air CO₂ enrichment and management. *Global Change Biology*, 3, 149–160.
- Hill, P.W., Marshall, C., Williams, G.G., Blum, H., Harmens, H., Jones, D.L. et al. (2007) The fate of photosynthetically-fixed carbon in *Lolium perenne* grassland as modified by elevated CO₂ and sward management. *New Phytologist*, 173, 766–777.
- Hyvönen, R., Agren, G.I., Linder, S., Persson, T., Cotrufo, M.F., Ekblad, A. et al. (2007) The likely impact of elevated [CO₂], nitrogen deposition, increased temperature and management on carbon sequestration in temperate and boreal forest ecosystems: a literature review. *The New Phytologist*, 173, 463–480.
- IPCC. (2007) *Climate change 2007: the physical science basis. Contribution of working group I. Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. Averyt, M. Tignor & H. Miller), pp. 20–35. Cambridge University Press, UK and New York, NY, USA.
- Jakobsen, I. & Rosendahl, L. (1990) Carbon flow into soil and external hyphae from roots of mycorrhizal cucumber plants. *New Phytologist*, 115, 77–83.
- Jin, V.L. & Evans, R.D. (2010) Microbial ¹³C utilization patterns via stable isotope probing of phospholipid biomarkers in Mojave Desert soils exposed to ambient and elevated atmospheric CO₂. *Global Change Biology*, 16, 2334–2344.
- Johnson, D., Leake, J.R. & Read, D.J. (2002) Transfer of recent photosynthate into mycorrhizal mycelium of an upland grassland: short-term respiratory losses and accumulation of ¹⁴C. *Soil Biology and Biochemistry*, 34, 1521–1524.
- Jongen, M., Jones, M.B., Hebeisen, T., Blum, H. & Hendrey, G. (1995) The effects of elevated CO₂ concentrations on the root growth of *Lolium perenne* and *Trifolium repens* grown in a FACE* system. *Global Change Biology*, 1, 361–371.
- Kongstad, J., Schmidt, I.K., Riis-Nielsen, T., Arndal, M.F., Mikkelsen, T.N. & Beier, C. (2012) High resilience in heathland plants to changes in temperature, drought, and CO₂ in combination: results from the CLIMAITE experiment. *Ecosystems*, 15, 269–283.
- Leake, J.R., Ostle, N.J., Rangel-Castro, J.I. & Johnson, D. (2006) Carbon fluxes from plants through soil organisms determined by field ¹³CO₂ pulse-labelling in an upland grassland. *Applied Soil Ecology*, 33, 152–175.
- Luo, Y., Jackson, R.B., Field, C.B. & Mooney, H.A. (1996) Elevated CO₂ increases belowground respiration in California grasslands. *Oecologia*, 108, 130–137.
- Lüscher, A., Hartwig, U.A., Suter, D. & Nösberger, J. (2000) Direct evidence that symbiotic N₂ fixation in fertile grassland is an important trait for a strong response of plants to elevated atmospheric CO₂. *Global Change Biology*, 6, 655–662.
- Montealegre, C.M., Van Kessel, C., Blumenthal, J.M., Hur, H.-G., Hartwig, U.A. & Sadowsky, M.J. (2000) Elevated atmospheric CO₂ alters microbial population structure in a pasture ecosystem. *Global Change Biology*, 6, 475–482.
- Montealegre, C.M., Van Kessel, C., Russelle, M.P. & Sadowsky, M.J. (2002) Changes in microbial activity and composition in a pasture ecosystem exposed to elevated atmospheric carbon dioxide. *Plant and Soil*, 243, 197–207.
- Moyano, F.E., Kutsch, W.L. & Schulze, E.-D. (2007) Response of mycorrhizal, rhizosphere and soil basal

- respiration to temperature and photo- synthesis in a barley field. *Soil Biology and Biochemistry*, 39, 843–853.
- Olsson, P.A. (1999) Signature fatty acids provide tools for determination of the distribution and interactions of mycorrhizal fungi in soil. *FEMS Microbiology Ecology*, 29, 303–310.
- Olsson, P.A. & Johnson, N.C. (2005) Tracking carbon from the atmosphere to the rhizosphere. *Ecology Letters*, 8, 1264–1270.
- Olsson, P.A., Van Aarle, I.M., Gavito, M.E., Bengtson, P. & Bengtsson, G. (2005) ^{13}C incorporation into signature fatty acids as an assay for carbon allocation in arbuscular mycorrhiza. *Applied and Environmental Microbiology*, 71, 2592–2599.
- Ostle, N., Ineson, P., Benham, D. & Sleep, D. (2000) Carbon assimilation and turnover in grassland vegetation using an in situ $^{13}\text{CO}_2$ pulse labeling system. *Rapid Communications in Mass Spectrometry*, 14, 1345–1350.
- Paterson, E., Gebbing, T., Abel, C., Sim, A. & Telfer, G. (2007) Rhizode-position shapes rhizosphere microbial community structure in organic soil. *New Phytologist*, 173, 600–610.
- Schortemeyer, M., Hartwig, U.A., Hendrey, G.R. & Sadowsky, M.J. (1996) Microbial community changes in the rhizosphere of white clover and perennial ryegrass exposed to free air carbon dioxide enrichment (FACE). *Soil Biology and Biochemistry*, 28, 1717–1724.
- Selsted, M.B., Linden, L., Ibrom, A., Michelsen, A., Larsen, K.S., Pedersen, J.K. et al. (2012) Soil respiration is stimulated by elevated CO_2 and reduced by summer drought: three years of measurements in a multifactor ecosystem manipulation experiment in a temperate heathland (CLIMAITE). *Global Change Biology*, 18, 1216–1230.
- Sowerby, A., Blum, H., Gray, T.R.G. & Ball, A.S. (2000) The decomposition of *Lolium perenne* in soils exposed to elevated CO_2 : comparisons of mass loss of litter with soil respiration and soil microbial biomass. *Soil Biology and Biochemistry*, 32, 1359–1366.
- Staddon, P.L., Jakobsen, I. & Blum, H. (2004) Nitrogen input mediates the effect of free-air CO_2 enrichment on mycorrhizal fungal abundance. *Global Change Biology*, 10, 1678–1688.
- Staddon, P.L., Ostle, N., Dawson, L.A. & Fitter, A.H. (2003) The speed of soil carbon throughput in an upland grassland is increased by liming. *Journal of Experimental Botany*, 54, 1461–1469.
- Strandberg, G. & Wilson, P. (1968) Formation of the nitrogen-fixing enzyme system in *Azotobacter vinelandii*. *Canadian Journal of Microbiology*, 14, 25–31.
- Suter, D., Frehner, M., Fischer, B.U., Nösberger, J. & Lüscher, A. (2002) Elevated CO_2 increases carbon allocation to the roots of *Lolium perenne* under free-air CO_2 enrichment but not in a controlled environment. *New Phytologist*, 154, 65–75.
- Talbot, J.M., Allison, S.D. & Treseder, K.K. (2008) Decomposers in disguise: mycorrhizal fungi as regulators of soil C dynamics in ecosystems under global change. *Functional Ecology*, 22, 955–963.
- Treonis, A.M., Ostle, N.J., Stott, A.W., Primrose, R., Grayston, S.J. & Ineson, P. (2004) Identification of groups of metabolically-active rhizosphere microorganisms by stable isotope probing of PLFAs. *Soil Biology and Biochemistry*, 36, 533–537.
- Van Kessel, C., Horwath, W.R., Hartwig, U., Harris, D. & Lüscher, A. (2000a) Net soil carbon input under ambient and elevated CO_2 concentrations: isotopic evidence after 4 years. *Global Change Biology*, 6, 435–444.
- Van Kessel, C., Nitschelm, J., Horwath, W.R., Harris, D., Walley, F., Lüscher, A. et al. (2000b) Carbon-13 input and turn-over in a pasture soil exposed to long-term elevated atmospheric CO_2 . *Global Change Biology*, 6, 123–135.
- Zanetti, S., Hartwig, U.A., Lüscher, A., Hebeisen, T., Frehner, M., Fischer, B.U. et al. (1996) Stimulation of symbiotic N_2 fixation in *Trifolium repens* L. under elevated atmospheric pCO_2 in a grassland ecosystem. *Plant Physiology*, 112, 575–583.