



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document, This is an Accepted Manuscript of an article published by Taylor & Francis in Journal of Mental Health on 13 January 2017, available online:
<http://dx.doi.org/10.1080/09638237.2016.1276535>:

Gorczyński, Paul Filip, Sitch, Matthew ORCID: 0000-0001-6521-7102 and Faulkner, Guy (2017) Examining methods, messengers and behavioural theories to disseminate physical activity information to individuals with a diagnosis of schizophrenia: a scoping review. Journal of Mental Health. pp. 1-10. doi:10.1080/09638237.2016.1276535

Official URL: <http://dx.doi.org/10.1080/09638237.2016.1276535>

DOI: <http://dx.doi.org/10.1080/09638237.2016.1276535>

EPrint URI: <http://eprints.glos.ac.uk/id/eprint/7074>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Examining methods, messengers, and behavioural theories to disseminate physical activity information to individuals with a diagnosis of schizophrenia: A scoping review

Paul Filip Gorczynski, Matthew Sitch & Guy Faulkner

Abstract

Background: Many individuals with a diagnosis of schizophrenia are not active and lack the necessary knowledge and confidence to become and stay active. To develop effective physical activity promotion interventions, it is necessary to identify credible messengers and effective methods to disseminate physical activity information to this population.

Aims: The purpose of this scoping review was to identify and examine knowledge mobilization theories, messengers, and methods used to disseminate physical activity information to individuals with a diagnosis of schizophrenia.

Method: This scoping review followed the methodological framework proposed by Arksey and O'Malley (2005).

Results: In total, 43 studies and 7 reviews identified multiple messengers and methods used to disseminate physical activity information to individuals with a diagnosis of schizophrenia, but few attempts to structure information theoretically. Findings do not point to which messengers or methods are most effective or which theories should be used to construct information interventions. Studies show that physical activity information should be provided in an individualised manner from staff who could easily connect with patients.

Conclusions: Few researchers have addressed the physical activity information needs of individuals with a diagnosis of schizophrenia. Researchers need to examine and implement effective knowledge mobilization strategies for this population.

Declaration of interest: Funding for this scoping review was provided by the Brain and Behaviour Research Foundation.

Keywords: Schizophrenia, physical activity, exercise, information, dissemination

Examining methods, messengers, and behavioural theories to disseminate physical activity information to individuals with a diagnosis of schizophrenia: A scoping review

Introduction

Physical activity can help individuals with a diagnosis of schizophrenia improve their health (Faulkner, Gorczynski, & Arbour-Nicitopoulos, 2013; Firth, Cotter, Elliot, French, & Yung, 2015; Gorczynski & Faulkner, 2010), yet people with this diagnosis are less active than those in the general population (Lindamer, et al., 2008; Stubbs et al., 2016). Low levels of physical activity and high levels of sedentary behaviour (see Stubbs, Williams, Gaughran, & Craig, 2016) are concerning given the majority of individuals with a diagnosis of schizophrenia are either overweight or obese and have high rates of type 2 diabetes mellitus (Hennekens, Hennekens, Hollar, & Casey, 2005; Vancampfort et al., 2015; Vancampfort et al., 2016). The high rate of obesity and diabetes in this population has resulted in life expectancies that are considerably lower than those in the general population (Laursen, Munk-Olsen, & Vestergaard, 2012; Walker, McGee, & Druss, 2015). Given that obesity and diabetes are independent risk factors for cardiovascular disease, strategies like increasing physical activity, are needed to decrease the incidence of morbidity and mortality in this population.

Research has shown that despite high interest in being active, individuals with a diagnosis of schizophrenia may lack the necessary knowledge and confidence to become and stay active (Johnstone, Nicol, Donaghy, Larie, 2009). Providing health and physical activity information to this population has been suggested as a possible strategy to increase physical activity participation and improve overall health (McDevitt, Snyder, Miller, & Wilbur, 2006; Faulkner, Gorczynski, & Cohn, 2009; Gorczynski, Faulkner, & Cohn, 2014). Providing relevant, skill based health information can help individuals obtain the necessary directions to change their behaviours and become active (Kreps, 2011). Although recent research has

shown that knowledge of the benefits of physical activity alone may not be sufficient to change activity behaviours in psychiatric populations (Happell, Stanton, Hoey, & Scott, 2014), providing such information may help individuals identify and overcome specific physical activity barriers and develop the skills and confidence necessary to become active. Useful and relevant information delivered in an efficient manner has also been identified as being an integral part of larger and more complex health promotion campaigns. The Ecological Model for active living has associated strong information environments with motivating individuals to become more active (Sallis, Cervero, Ascher, Henderson, Kraft, & Kerr, 2006). Despite this need for greater health and physical activity information, no research has examined the design or effectiveness of knowledge mobilization strategies to deliver physical activity information to individuals with a diagnosis of schizophrenia. In effect, no systematic efforts have been made to understand target audience needs, how messages should be framed, and which methods and messengers should be used to disseminate physical activity information in this population. This scoping review aims to fill this gap and lay the groundwork necessary to design effective knowledge mobilization campaigns to help individuals with a diagnosis of schizophrenia become and stay active. Scoping reviews aim to map out the key concepts and gaps of a research area and the main sources of evidence available. They are inclusive of different study designs, topics, and sources of literature and allow for a participatory approach to obtaining and using information (Martin Ginis et al., 2012; Lavis, Robertson, Woodside, McLeod, & Abelson, 2003; Faulkner et al., 2009; Arksey & O'Malley, 2005). The purpose of this scoping review was to identify and examine knowledge mobilization theories, messengers, and methods used to disseminate physical activity information to individuals with a diagnosis of schizophrenia. Additionally, where possible, barriers and facilitators to delivering physical activity information for each messenger and method were examined.

Methods

This scoping review followed the methodological framework proposed by Arksey and O'Malley (2005) as well as previous research by Faulkner et al. (2009). This process was carried out in five stages:

1) Identifying the research questions. The questions of this scoping review were designed to identify gaps in existing literature regarding knowledge mobilization about physical activity information for individuals with a diagnosis of schizophrenia. Specifically, these questions were designed to explore knowledge mobilization theories, messengers, and methods with respect to physical activity information dissemination. These questions included: a) What are the knowledge mobilization theories used to disseminate physical activity information for individuals with a diagnosis of schizophrenia?; b) Who are credible and preferred messengers for physical activity information for individuals with a diagnosis of schizophrenia?; and c) What are the preferred methods of information dissemination about physical activity for individuals with a diagnosis of schizophrenia? Information pertaining to barriers and facilitators to delivering physical activity information for each messenger and method were also extracted and examined. Questions were based on previous work conducted by Faulkner et al. (2009) as well as the knowledge mobilization model described by Lavis et al. (2003).

2) Identifying relevant studies. The first two authors searched for and selected relevant literature that pertained to our three research questions. Two sources of information were searched for relevant literature, including: 1) published studies; and 2) contextual literature. For published studies, no limits were imposed on study types or study dates. The following databases were searched in March 2016: PSYCHINFO, PUBMED, PSYCHARTICLES, and GOOGLE SCHOLAR. A search strategy used by Faulkner and colleagues (2009) in a

scoping review examining knowledge mobilization for individuals with spinal cord injury was used for this review. The following terms were used to search databases: schizophrenia, physical activity, exercise, information, source, needs, knowledge, mobilization, message, messenger, dissemination, help, management, and health promotion. The reference lists of included studies were also searched for additional studies. Published studies were included in the results if they were written in English, involved individuals with a diagnosis of schizophrenia, discussed aspects of information dissemination for physical activity, and contained information related to any of the three questions of the scoping review. No time limits were imposed on this search. For contextual literature, systematic reviews, meta-analyses, and book chapters were examined. These sources of literature were identified through searches of the same databases used to identify published studies, using the same search terms. A ‘review of reviews’ was conducted to identify the best current information in order to develop rigorous knowledge mobilization strategies to promote physical activity to individuals diagnosed with schizophrenia.

3) Study selection. The first two study authors examined study titles and abstracts and determined whether studies could be considered for inclusion within the scoping review. The authors met throughout the review process to ensure the identification, review, and selection process was consistent. The authors independently reviewed literature to ensure it met inclusion criteria (i.e., provided information that addressed any of the three questions of the scoping review). When disagreements arose, the two authors reached a consensus after a discussion.

4) Charting the data. A data-charting form was used to extract relevant information on each identified piece of literature. The first two authors met to compare extraction results for consistency and made modifications where necessary. Information that was extracted and charted included: author(s); year; study location; definition of physical activity or exercise;

description of knowledge mobilization theories; messengers; methods; results; and any other key findings.

5) Collating, summarizing, and presenting the results. Findings were thematically organized with respect to mobilization theories, messengers, and methods. Extracted data from our identified literature was written in a narrative manner. Given the diversity of studies retrieved, no attempts were made to assess the quality of evidence presented.

Results

A total of 50 articles (43 studies and 7 reviews) were identified from the literature search that provided an answer to at least one of the stated research questions. Full results of the research articles can be seen in Table 1. Progression of the literature search can be seen in Figure 1. below.

<<insert Figure 1. Here>>

Studies were carried out in a number of nations globally, with the majority being conducted in North America (n=20) and Europe (n=12). Sample sizes ranged from 3 (Faulkner & Sparkes, 1999) to 732 (Motlova, Dragomirecka, & Kitzlerova, 2009), although with the exception of three studies, all were under 100 participants.

Study settings were predominantly hospitals and mental health centres, utilising samples of individuals who were in-patients. Two studies (Gomes et al., 2014; Warren et al., 2011) were set in university campus facilities with out-patient populations recruited through psychiatric day-care units. A range of exercises and types of physical activity were used in studies, with the most common being walking and jogging (Attux et al., 2013; Beebe et al., 2009; Beebe et al., 2013; P. Bernard et al., 2013; Faulkner & Sparkes, 1999; Gorczynski, Faulkner, Cohn, & Remington, 2014a; Maggouritsa et al., 2014; Melamed et al., 2008; Methapatara & Srisurapanont, 2011; Sailer et al., 2015; Warren et al., 2011; Wu, Wang, Bai, Huang, & Lee, 2007). Other studies used Yoga (Behere et al., 2011; Duraiswamy, Thirthalli,

Nagendra, & Gangadhar, 2007), video games (Leutwyler, Hubbard, Vinogradov, & Dowling, 2012) and small-sided competitive games (Gomes et al., 2014).

Knowledge Mobilisation Theories

Nine studies made reference to theory when disseminating information about physical activity. Studies that made reference to theory utilised a range of theories including the Trans-theoretical model (see Prochaska, Johnson, & Lee, 2009) (Bernard et al. 2013), Cognitive and Behavioural Therapy (Attux et al., 2013), Psychoanalysis (Pesek et al., 2011), mental contrasting and implementation strategies (see Oettingen & Gollwitzer, 2010) (Sailer et al., 2006), Self-Determination Theory (see Deci & Ryan, 2011) (Gorczyński et al., 2014a, Gorczyński et al., 2014b), and Social Cognitive Theory (see Bandura, 1986) (Beebe et al., 2010; Beebe et al., 2013). The use of these theories was integrated into the design and implementation of studies and appeared to contribute toward the development of interventions that aimed to help individuals diagnosed with schizophrenia progressively improve their levels of physical activity. For example Bernard et al. (2013) utilised the Trans-theoretical model of behaviour change (Prochaska, Johnson, & Lee, 2009) to guide the content of a counselling intervention, integrating core principles of the model (e.g., *processes of change* and *decisional balance*) into the intervention. Gorczyński et al. (2014a and 2014b) utilised aspects from the Trans-theoretical model, Social Cognitive Theory, and Self-Determination Theory (Deci & Ryan, 2011) to guide their exercise counselling intervention. The thorough application of theory served to guide not only knowledge mobilisation through effective counselling practice, but also gauge behaviour change by examining movement through stages of change. Sailer et al. (2006) applied a theory based Mental Contrasting and Implementing Intentions approach to guide a goal setting intervention to increase exercise in individuals with a diagnosis of schizophrenia. Sailer et al. paid attention to the importance of

social context, creating an autonomy-focused environment for successful goal pursuit. They found support for the approach in improving attendance rates at scheduled exercise sessions

Messengers of physical activity information

Twenty-nine studies identified a variety of messengers who provided physical activity information to individuals with a diagnosis of schizophrenia. These messengers can be separated into three broad categories that include mental health care professionals, research staff, and non-specified. Mental health care professionals were the primary physical activity message providers and included care workers (Faulkner & Sparkes, 1999), clinical staff (Littrell, Hilligoss, Kirshner, Petty, & Johnson, 2003; Pelham & Campagna, 1991; Sailer et al., 2015; Warren et al., 2011), nurses (R. Bernard et al., 1990; Melamed et al., 2008; Motlova et al., 2009; Niv, Cohen, Hamilton, Reist, & Young, 2014), therapists (e.g. physiotherapists) (Duraiswamy et al., 2007; Stubbs et al., 2014a, Stubbs et al., 2014b), psychiatrists (Chen, Chen, & Huang, 2009; Pesek, Mihoci, Medved, & Solinc, 2011), and exercise psychologists (Dodd, Duffy, Stewart, Impey, & Taylor, 2011). Other studies stated that research staff were the primary physical activity messengers (Beebe et al., 2009; Beebe et al., 2010; Beebe et al., 2013; Gorczynski et al., 2014a; Gorczynski, Faulkner, Cohn, & Remington, 2014b; Leutwyler et al., 2012; Methapatara & Srisurapanont, 2011), whilst other studies did not specify who were messengers (Acil, Dogan, & Dogan, 2008; Maggouritsa et al., 2014; McKibbin et al., 2006; Wu et al., 2007). Although many messengers were identified, the effectiveness of messengers was not assessed. A number of studies recommended the use of multiple messengers (Marzolini, Jensen, & Melville, 2009; Tetlie & Polit, 2009). For example Tetlie and Polit (2009) utilised psychiatric nurses to provide personalised messages regarding exercise participation while exercise instructors provided the exercise program. Only one study mentioned the use of patients' relatives in the provision of physical activity messages to enhance intervention effectiveness, citing that the use of

relatives was of particular importance where patients live with their families (Attux et al., 2013).

Methods of Physical Activity Information Dissemination

A total of 13 studies provided a wide range of methods to disseminate information regarding physical activity to individuals with a diagnosis of schizophrenia. A number of studies utilised structured educational sessions to inform study participants (Attux et al., 2013; Littrell et al., 2003; McKibbin et al., 2006; Motlova et al., 2009; Niv et al., 2014; Pesek et al., 2011). These sessions were often multidimensional in message delivery. For example, Littrell et al. (2003) utilised a combination of verbal and written information, reading aloud, discussion of topics, written exercises, quizzes and educational games to convey messages regarding physical activity. The use of classroom-based education sessions were typically part of wider interventions designed to improve health behaviours in general (e.g., diet, prescription drug use) and specific measures of physical activity were not taken. Other studies opted to use an approach which promoted independence amongst participants by providing physical activity information in a leaflet (Methapatara & Srisurapanont, 2011) or encouraging participants to keep an exercise diary (Chen et al., 2009; Gorczynski et al., 2014b; Kwon et al., 2006). Gorczynski, Faulkner, Cohn, and Remington (2013) aimed to increase stair use within a mental health centre with the use of stair riser banners that prompted patients and staff to “Take the stairs, stay healthy”. Whilst results over a 6-week period indicated no significant change in stair use amongst patients, the study’s use of stair risers to disseminate physical activity information had an effect on increasing staff stair usage, potentially creating a more activity conscious hospital environment. Similarly, Gorczynski et al. (2014a) examined the feasibility of accelerometer use with individuals with a diagnosis of schizophrenia, and presented that with some compliance strategies an objective measure of physical activity might be used as a motivator to be active, a finding that echoed

Beebe and Faust-Harris' (2012) work. How messages regarding physical activity are disseminated to patients appeared to be an oversight for many studies in that exercise programs were implemented without mention of the methods used to educate participants of what activities should be done, how they should be performed, or why they should be done.

Barriers and Facilitators to Message Dissemination

A number of studies (n=6) provided insight into some of the barriers and facilitators to physical activity message dissemination, (Crone, 2007; Gorczynski, Faulkner, & Cohn, 2013; Johnstone, Nicol, Donaghy, & Lawrie, 2009; Leutwyler, Hubbard, Jeste, & Vinogradov, 2013; Rastad, Martin, & Asenlof, 2014; Stubbs et al., 2014a; Stubbs et al., 2014b). Stubbs et al. (2014a and 2014b) examined the perceptions of physiotherapists' working in mental health on the assessment, benefits and delivery of physical activity in individuals with a diagnosis of schizophrenia. A key finding was that the inclusion of physiotherapists in physical activity programs was felt to be a facilitator to physical activity participation. Physiotherapists perceived that they had the necessary theoretical knowledge and clinical skills required to lead and oversee physical activity programs with individuals with a diagnosis of schizophrenia. Leutwyler et al. (2013) similarly posited how psychiatric staff felt that they played a key role in modelling and motivating physical activity for patients. Furthermore, it was reported how staff perceived patients were reluctant to join groups and preferred one-to-one support for physical activity. The personal support of health care staff was identified as a facilitator to message dissemination, with the impetus placed on messengers whom patients knew for a long period of time, staff whom patients trusted and felt connected with (Rastad et al., 2014). Similarly, Crone (2007) outlined that individuals with a diagnosis of schizophrenia attending a walking intervention thought that their participation was facilitated by the interpersonal skills of the intervention staff. Patients' trust and respect of messengers is a recurrent theme emerging from studies as an important factor

influencing physical activity message dissemination. For example, Johnstone et al., (2009) highlighted social anxiety as a significant barrier to physical activity participation with many individuals living with a diagnosis of schizophrenia finding it difficult to leave their home environment. Effective messengers were those who individuals trusted and respected including family and friends, long-term health care providers, and fellow patients. Other barriers included a lack of choice in physical activities, misconceptions about physical activity such as only intensive exercise was meaningful or you have to lose weight to be active, and negative expectations of physical activity such as fear of not succeeding (Rastad et al. 2014; Gorczynski et al., 2013). These barriers pose important considerations for physical activity intervention design as researchers and health providers need to carefully select information messengers who are trusted and familiar to patients and what information they are conveying.

Review of Reviews

Many review papers exist detailing the literature that has sought to examine physical activity and exercise interventions for individuals with a diagnosis of schizophrenia. From the range of review papers available, seven provide important points with regard to literature pertaining to messengers and the provision of messages in the promotion of physical activity (Faulkner, Cohn, & Remington, 2010; Faulkner & Gorczynski, 2014; Faulkner, Gorczynski, & Arbour-Nicitopolous, 2013; Gross, Vancampfort, Stubbs, Gorczynski, & Soundy, 2016; Holley, Crone, Tyson, & Lovell, 2011; Soundy et al., 2014; Vancampfort et al., 2011).

Almost unanimous agreement was found in review studies regarding concern over the lack of theoretical underpinning in research studies' design and implementation which has led to an absence of understanding of the possible mechanisms responsible for intervention success. Gross et al.'s (2015) review of the value of social support in the promotion of physical activity offers useful insight into the importance of considering support in the operation of

physical activity interventions. Findings from 23 studies indicated that although informational support was the most abundant form of social support in studies, esteem support was the most valued by staff and patients. Gross et al. iterate the importance of social support for physical activity intervention outcome with emphasis placed on patient's sensation of trust and value generated through supportive exchanges with intervention organisers, particularly during early stages of intervention implementation. The importance of social support provision was echoed by other review studies including Faulkner et al. (2010) and Soundy et al. (2014) where support was considered an essential feature of message dissemination and barrier confrontation. For example, Soundy et al. (2014) outlines how research has identified that social anxiety is a prominent barrier to physical activity participation and effective support from other patients and care providers was seen to be a vital aspect influencing intervention outcome.

Discussion

This scoping review examined knowledge mobilization theories, messengers, and methods used to disseminate physical activity information to individuals with a diagnosis of schizophrenia. In total, 50 research articles, including studies and reviews, pointed to multiple messengers and methods used to disseminate physical activity information to people individuals with a diagnosis of schizophrenia, but few attempts to structure information using a theoretical approach. Most common messengers and methods of physical activity included clinical and research staff and educational approaches, respectively. Findings from the examined studies and reviews revealed that physical activity information should be provided in an individualised manner from staff who could easily connect with patients in a trusting manner. A number of researchers also recognized that physical activity is a social experience and emphasised that strategies should be put in place to help individuals with a diagnosis of schizophrenia address any fears or anxieties they may have with being social or being active. The incorporation of family and friends and other caregivers within physical activity programs may be a way to help address these anxieties and fears. Overall physical activity information should be provided in a manner that affords individuals with a diagnosis of schizophrenia the ability to be autonomous, feel connected to others in a supportive manner, and informed about the physical activity choices they make so as to improve their physical, mental, and social health.

In line with previous reviews examining physical activity interventions in individuals with diagnosis schizophrenia (Faulkner, Cohn, & Remington, 2010; Faulkner & Gorczynski, 2014; Faulkner, Gorczynski, & Arbour-Nicitopolous, 2013; Gross, Vancampfort, Stubbs, Gorczynski, & Soundy, 2016; Holley, Crone, Tyson, & Lovell, 2011; Soundy et al., 2014; Vancampfort et al., 2011), there was a lack of theory utilised in the studies found for this scoping review. The few studies that used theory in their design and application were able to

illustrate how theories may be used to inform and enhance aspects crucial to message design and transmission. Overall, theory based physical activity interventions are more effective than those designed not using any theory (Kahn et al., 2002). The use of theory can specifically help improve information dissemination by guiding message construction and the selection of helpful messengers and methods. Additionally, the use of theory in the construction of interventions can address specific known barriers individuals with a diagnosis of schizophrenia face to receiving information about being active. Two important areas that researchers may wish to focus on are ways of providing individuals with a diagnosis of schizophrenia information that aids with motivational support and impaired cognition (Vancampfort et al., 2013). Previous research has shown significant positive correlations between autonomous regulation, both extrinsic regulation (i.e., identified regulation) and intrinsic regulation, and levels of physical activity (Vancampfort et al., 2013). Providing physical activity information that is easily attainable, states the benefits of regular activity, and easy to understand that ultimately strengthens an individual's freedom of choice and ability to connect with others may prove beneficial to helping individuals with a diagnosis of schizophrenia find and participate in enjoyable physical activities both initially and for a prolonged period of time (Teixeira, Carraça, Markland, Silva, & Ryan, 2012). Such provisions to constructing and distributing health information to ensure all individuals, regardless of ability or impairment, can improve their overall wellbeing should be ensured by healthcare providers and healthcare systems, as has recently been done by the National Health Service (NHS) in the United Kingdom through the Accessible Information Standard (see NHS, 2016). Addressing this lack of theory in future research is essential and necessary to improve the overall quality of physical activity interventions for individuals with a diagnosis of schizophrenia (Faulkner & Gorczynski, 2014). Furthermore, the use of theory

can improve our understanding of the mechanisms underpinning physical activity message mobilisation within this population.

With regard to messengers and methods of physical activity information dissemination, there is a general lack of consideration within interventions as to how information should be transmitted to individuals with a diagnosis of schizophrenia. Additionally, the effectiveness of messengers and methods to provide physical activity information to individuals with a diagnosis of schizophrenia has not been addressed. Studies that were considerate of messengers (e.g. Marzolini et al., 2007; Tetlie & Polit, 2007) identified that using multiple messengers and family members may prove beneficial to overall information dissemination and physical activity participation. In line with previous research that has investigated the use of family interventions in the treatment of psychosis (Harvey & O'Hanlon, 2013), the involvement of family, friends, and caregivers to promote physical activity seems like a natural fit. Overall, family members want to be involved in the care and treatment of their relatives living with psychosis (Parker et al., 2010). Involving family, friends, and caregivers to learn about physical activity opportunities for their relatives may afford individuals with a diagnosis of schizophrenia the support that is necessary to learn about multiple physical activity options in their respective communities and then engage in physical activity through a supported manner. Physical activity may also prove to be a benefit for family, friends, and caregivers who may be experiencing caregiver fatigue and potential mental health problems of their own (Awad & Voruganti, 2008). Findings from a recent study show that carers are ready to be involved in research that will enhance the overall treatment of their loved ones living with serious mental health issues (Ashcroft, Wykes, Taylor, Crowther, & Szmukler, 2016). Further research into the involvement of physical activity programming into family psycho-education is needed and warranted.

The lack of theoretical consideration in the design of interventions and specifically to guide message mobilisation in studies is also of concern. Owing to the range and diversity of study designs and settings it is not possible to assess the contribution of theoretical consideration on study efficacy. However, the contribution of theory appears to enhance the robustness of study design and implementation by informing and justifying researchers' decision making. This enhancement can also be seen in research by Bradshaw et al. (2010) who did not make reference to a specific knowledge mobilisation theory per se but rather utilised the Medical Research Council (MRC) framework for the development and evaluation of randomised control trials to guide their study (Craig, Dieppe, Macintyre, Michie, Nazareth, & Petticrew, 2008). The MRC framework incorporates a range of theoretical underpinnings designed to guide practice and its use within Bradshaw and colleagues' study led to a clearly progressive, effective intervention.

The findings of this review offer several implications for healthcare professionals as to how best deliver physical activity information to individuals with a diagnosis of schizophrenia. Findings from the practice of physiotherapy shows that physiotherapists are comfortable with prescribing and helping individuals with a diagnosis of schizophrenia become physically active (Stubbs et al., 2014a; Stubbs et al., 2014b; Vancampfort, Rosenbaum, Schuch, Ward, Probst, & Stubbs, 2016). Unfortunately, similar work has not occurred with other healthcare professions. Research is needed to ensure other healthcare professions feel confident in their knowledge of physical activity guidelines, physical activity opportunities available to their patients, and methods in which they can effectively provide physical activity information to their patients. Such research would help ensure that appropriate training is created and offered to help healthcare professionals gain the necessary confidence to prescribe physical activity to their patients. Healthcare professionals need to also take steps to help individuals with a diagnosis of schizophrenia feel autonomous and

supported in their ability to become physically active. Providing information about various physical activity options is essential as is ensuring individuals with a diagnosis of schizophrenia have the necessary support to continue to be active. Incorporating family, friends, and other caregivers is one step to ensuring individuals with a diagnosis of schizophrenia can become and stay active. Healthcare professionals also need to better understand which methods of information dissemination are best for their patients. Further research is needed to identify effective strategies to deliver physical activity information to patients and at what stage of the recovery process.

A number of limitations with the current scoping review must be mentioned. First, it is possible that all relevant studies may not have been identified despite our systematic approach based on a similar scoping review to locating necessary literature. Second, scoping reviews offer a narrative or descriptive account of the literature in a particular field of study, and do not offer a synthesis of which intervention is best (Arksey & O'Malley, 2005). Findings from this current scoping review cannot point to which messengers or methods are most effective with respect to physical activity information dissemination to individuals with a diagnosis of schizophrenia or which theories should be used to construct information interventions. Rather, this scoping review offers suggestions for further targeted inquiry. The health benefits of physical activity are well known for both clinical and non-clinical populations. In the context of schizophrenia, the greatest challenge is not a lack of evidence; it is absent, inconsistent, or failed implementation of the evidence. This scoping review provides a foundation for future efforts in addressing effective knowledge translation.

References

- Acil, A., Dogan, S., & Dogan, O. (2008). The effects of physical exercises to mental state and quality of life in patients with schizophrenia. *Journal of Psychiatric and Mental Health Nursing*, 15(10), 808-815.
- Arksey, H., & O'Malley, L. (2005). Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology*, 8(1), 19-32.
- Ashcroft, J., Wykes, T., Taylor, J., Crowther, A., & Szmukler, G. (2016). Impact on the individual: What do patients and carers gain, lose, and expect from being involved in research? *Journal of Mental Health*, 21(1), 28-35.
- Attux, C., Martini, L. C., Elkis, H., Tamai, S., Freirias, A., Camargo, M. d., Mateus, M. D., Mari Jde, J., Reis, A. F., & Bressan, R. A. (2013). A 6-month randomized controlled trial to test the efficacy of a lifestyle intervention for weight gain management in schizophrenia. *BMC Psychiatry*, 13(1), 60.
- Awad, A. G., & Voruganti, L. N. (2008). The burden of schizophrenia on caregivers: A review. *Pharmacoeconomics*, 26(2), 149-162.
- Bandura A. (1986). *Social foundations of thought and action*. Upper Saddle River, NJ: Prentice Hal.
- Beebe, L. H., Burk, R., McIntyre, K., Smith, K., Velligan, D., Resnick, B., Tavakoli, A., Tennison, C., & Dessieux, O. (2009). Motivating Persons with Schizophrenia Spectrum Disorders to Exercise: Rationale and Design. *Clinical Schizophrenia and Related Psychoses*, 3(2), 111-116.
- Beebe, L. H., Smith, K., Burk, R., Dessieux, O., Velligan, D., Tavakoli, A., & Tennison, C. (2010). Effect of a Motivational Group Intervention on Exercise Self-Efficacy and Outcome Expectations for Exercise in Schizophrenia Spectrum Disorders. *Journal of the American Psychiatric Nurses Association*, 16(2), 105-113.

- Beebe, L. H., Smith, K. D., Roman, M. W., Burk, R. C., McIntyre, K., Dessieux, O. L., Tavakoli, A., & Tennison, C. (2013). A pilot study describing physical activity in persons with schizophrenia spectrum disorders (SSDS) after an exercise program. *Issues in Mental Health Nursing*, 34(4), 214-219.
- Behere, R., Arasappa, R., Jagannathan, A., Varambally, S., Venkatasubramanian, G., Thirhalli, J., Subbakrishna, D. K., Nagendra, H. R., & Gangadhar, B. (2011). Effect of yoga therapy on facial emotion recognition deficits, symptoms and functioning in patients with schizophrenia. *Acta Psychiatrica Scandinavica*, 123(2), 147-153.
- Bernard, P., Esseul, E., Raymond, E., Dandonneau, L., Xambo, J., Carayol, M., & Ninot, G. (2013). Counseling and exercise intervention for smoking reduction in patients with schizophrenia: a feasibility study. *Archives of Psychiatric Nursing*, 27(1), 23-31.
- Bernard, R., Johnsen, E. C., Killworth, P. D., McCarty, C., Shelley, G. A., & Robinson, S. (1990). Comparing four different methods for measuring personal social networks. *Social Networks*, 12(3), 179-215.
- Chen, C. K., Chen, Y. C., & Huang, Y. S. (2009). Effects of a 10-week weight control program on obese patients with schizophrenia or schizoaffective disorder: a 12-month follow up. *Psychiatry and Clinical Neuroscience*, 63(1), 17-22.
- Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew. (2008). Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ*, 337, 979-983.
- Crone, D. (2007). Walking back to health: a qualitative investigation into service users' experiences of a walking project. *Issues in Mental Health Nursing*, 28(2), 167-183.
- Deci, E. L., & Ryan, R. M. (2011) Self-determination theory. In P. Van Lange, A. Kruglanski & T. Higgins (Eds.), *Handbook of Theories of Social Psychology* (Vol. 1, pp. 416-433). Thousand Oaks, CA: SAGE Publications.

- Dodd, K. J., Duffy, S., Stewart, J. A., Impey, J., & Taylor, N. (2011). A small group aerobic exercise programme that reduces body weight is feasible in adults with severe chronic schizophrenia: a pilot study. *Disability and Rehabilitation*, 33(13-14), 1222-1229.
- Duraiswamy, G., Thirthalli, J., Nagendra, H. R., & Gangadhar, B. N. (2007). Yoga therapy as an add-on treatment in the management of patients with schizophrenia--a randomized controlled trial. *Acta Psychiatrica Scandinavica*, 116(3), 226-232.
- Faulkner, G., Cohn, T., & Remington, G. (2010). Interventions to reduce weight gain in schizophrenia. *Cochrane Database of Systematic Reviews*, 1, CD005148.
- Faulkner, G., & Gorczynski, P. (2014). Schizophrenia. In A. Clow & S. Edmunds (Eds.), *Physical Activity and Mental Health* (pp. 215-235). Champaign, IL: Human Kinetics.
- Faulkner, G., Gorczynski, P., & Arbour-Nicitopolous, K. (2013). Exercise as an adjunct treatment for schizophrenia. In P. Ekkekakis (Ed.), *Routledge Handbook of Physical Activity and Mental Health* (pp. 541-555). New York, NY: Routledge.
- Faulkner, G., Gorczynski, P., Arbour, K., Letts, L., Wolfe, D., & Martin Ginis, K. (2009). Messengers and methods of disseminating health information among individuals with spinal cord injury: A scoping review. In T.C. Berkovsky (Ed.), *Handbook of Spinal Cord Injuries: Types, Treatments and Prognosis* (pp. 349-374). Hauppauge, NY: Nova Science Publishers, Inc.
- Faulkner, G., Gorczynski, P., & Cohn T. (2009). Psychiatric illness and obesity: Recognizing the 'Obesogenic' nature of an inpatient psychiatric setting. *Psychiatric Services*, 60(4), 538-541.
- Faulkner, G., & Sparkes, A. (1999). Exercise as therapy for schizophrenia: An ethnographic study. *Journal of Sport and Exercise Psychology*, 21, 52-69.

- Firth, J., Cotter, J., Elliot, R., French, R., & Yung, A. R. (2015). A systematic review and meta-analysis of exercise interventions in schizophrenia patients. *Psychological Medicine*, 45, 1343-1361.
- Gomes, E., Bastos, T., Probst, M., Ribeiro, J. C., Silva, G., & Corredeira, R. (2014). Effects of a group physical activity program on physical fitness and quality of life in individuals with schizophrenia. *Mental Health and Physical Activity*, 7(3), 155-162.
- Gorczyński, P., & Faulkner, G. (2010). Exercise therapy for schizophrenia. *Cochrane Database of Systematic Reviews*, 5, CD004412.
- Gorczyński, P., Faulkner, G., & Cohn, T. (2013). Dissecting the Obesogenic Environment of a Psychiatric Setting: Client Perspectives. *Canadian Journal of Community Mental Health*, 32(3), 51-68.
- Gorczyński, P., Faulkner, G., Cohn, T., & Remington, G. (2013). Effects of point-of-choice prompts on stair usage in a psychiatric setting. *Psychiatric Services*, 64(5), 498.
- Gorczyński, P., Faulkner, G., Cohn, T., & Remington, G. (2014a). Examining strategies to improve accelerometer compliance for individuals living with schizophrenia. *Psychiatric Rehabilitation Journal*, 37(4), 333-335.
- Gorczyński, P., Faulkner, G., Cohn, T., & Remington, G. (2014b). Examining the efficacy and feasibility of exercise counseling in individuals with schizophrenia: A single-case experimental study. *Mental Health and Physical Activity*, 7(3), 191-197.
- Gross, J., Vancampfort, D., Stubbs, B., Gorczyński, P., & Soundy, A. (2016). A narrative synthesis investigating the use and value of social support to promote physical activity among individuals with schizophrenia. *Disability and Rehabilitation*, 38(2), 123-150.
- Happell, B., Stanton, R., Hoey, W., & Scott, D. (2014). Knowing is not doing: The relationship between health behaviour knowledge and actual health behaviours in

people with serious mental illness. *Mental Health and Physical Activity*, 7(3), 198-204.

Harvey, C., & O'Hanlon, B. (2013). Family psycho-education for people with schizophrenia and other psychotic disorders and their families. *Australian and New Zealand Journal of Psychiatry*, 47(6), 516-520.

Hennekens, C.H., Hennekens, A.R., Hollar, D., & Casey, D.E. (2005). Schizophrenia and increased risks of cardiovascular disease. *American Heart Journal*, 150(6), 1115-1121.

Holley, J., Crone, D., Tyson, P., & Lovell, G. (2011). The effects of physical activity on psychological well-being for those with schizophrenia: A systematic review. *British Journal of Clinical Psychology*, 50(1), 84-105.

Johnstone, R., Nicol, K., Donaghy, M., & Lawrie, S. (2009). Barriers to uptake of physical activity in community-based patients with schizophrenia. *Journal of Mental Health*, 18(6), 523-532.

Kahn, E.B., Ramsey, L. T., Brownson, R. C., Heath, G. W., Howze, E. H., Powell, K. E., Stone, E.J., Rajab, M.W., & Corso, P. (2002). The effectiveness of interventions to increase physical activity: A systematic review. *American Journal of Preventative Medicine*, 22(4suppl.), 73-107.

Kreps, G.L. (2001) The evolution and advancement of health communication inquiry. In: W. B. Gudykunst (Ed.), *Communication Yearbook 24* (pp. 232-254). Thousand Oaks: Sage.

Kwon, J. S., Choi, J.-S., Bahk, W.-M., Yoon Kim, C., Hyung Kim, C., Chul Shin, Y., Park, B. J., & Geun Oh, C. (2006). Weight management program for treatment-emergent weight gain in olanzapine-treated patients with schizophrenia or schizoaffective disorder: a 12-week randomized controlled clinical trial. *Journal of Clinical Psychiatry*, 67(4), 547-553.

- Laursen, T.M., Munk-Olsen, T., & Vestergaard, M. (2012). Life expectancy and cardiovascular mortality in persons with schizophrenia. *Current Opinion in Psychiatry*, 25(2), 83-88.
- Lavis, J.N., Robertson, D., Woodside, J.M., McLeod, C.B., & Abelson J. (2003). How can research organizations more effectively transfer research knowledge to decision makers? *The Milbank Quarterly*, 81, 221-248.
- Leutwyler, H., Hubbard, E. M., Jeste, D. V., & Vinogradov, S. (2013). "We're not just sitting on the periphery": a staff perspective of physical activity in older adults with schizophrenia. *Gerontologist*, 53(3), 474-483.
- Leutwyler, H., Hubbard, E. M., Vinogradov, S., & Dowling, G. (2012). Videogames to promote physical activity in older adults with schizophrenia. *Games for Health: Research, Development, and Clinical Applications*, 1(5), 381-383.
- Lindamer, L.A., McKibbin, C., Norman, G.J., Jordan, L., Harrison, K., Abeyesinhe, S., & Patrick, K. (2008). Assessment of physical activity in middle-aged and older adults with schizophrenia. *Schizophrenia Research*, 104(1-3), 294-301.
- Littrell, K. H., Hilligoss, N. M., Kirshner, C. D., Petty, R. G., & Johnson, C. G. (2003). The effects of an educational intervention on antipsychotic-induced weight gain. *Journal of Nursing Scholarship*, 35(3), 237.
- Magouritsa, G., Kokaridas, D., Theodorakis, I., Patsiaouras, A., Mouzas, O., Dimitrakopoulos, S., & Diggelidis, N. (2014). The effect of a physical activity programme on improving mood profile of patients with schizophrenia. *International Journal of Sport and Exercise Psychology*, 12(3), 273-284.
- Martin Ginis, K.A., Latimer-Cheung, A., Corkum, S., Ginis, S., Anathasopoulos, P., Arbour-Nicitopoulos, K., & Gainforth, H. (2012). A case study of a community-university multidisciplinary partnership approach to increasing physical activity participation

among people with spinal cord injury. *Translational Behavioral Medicine*, 2(4), 516-522.

Marzolini, S., Jensen, B., & Melville, P. (2009). Feasibility and effects of a group-based resistance and aerobic exercise program for individuals with severe schizophrenia: A multidisciplinary approach. *Mental Health and Physical Activity*, 2(1), 29-36.

McDevitt, J., Snyder, M., Miller, A., & Wilbur, J. (2006). Perceptions of barriers and benefits to physical activity among outpatients in psychiatric rehabilitation. *Journal of Nursing Scholarship*, 38(1), 50-55.

McKibbin, C. L., Patterson, T. L., Norman, G., Patrick, K., Jin, H., Roesch, S., Mudalliar, S., Barrio, C., O'Hanlon, K., Griver, K., Sirkin, A., & Jeste, D. V. (2006). A lifestyle intervention for older schizophrenia patients with diabetes mellitus: a randomized controlled trial. *Schizophrenia Research*, 86, 36-44.

Melamed, Y., Stein-Reisner, O., Gelkopf, M., Levi, G., Sivan, T., Ilievici, G., Rosenberg, R., Weizman, A., & Bleich, A. (2008). Multi-modal weight control intervention for people with persistent mental disorders. *Psychiatric Rehabilitation Journal*, 31(3), 194-200.

Methapatara, W., & Srisurapanont, M. (2011). Pedometer walking plus motivational interviewing program for Thai schizophrenic patients with obesity or overweight: a 12-week, randomized, controlled trial. *Psychiatry and Clinical Neuroscience*, 65(4), 374-380.

Motlova, B. L., Dragomirecka, E., & Kitzlerova, E. (2009). Weight control programme for schizophrenia: bridge between psychiatrists and primary care physicians. *European Psychiatry*, 24(7), 490-491.

National Health Service (2016). NHS England launches accessible information standard.

Retrieved August 10, 2016, from <https://www.england.nhs.uk/2015/07/access-info-standard/>

Niv, N., Cohen, A. N., Hamilton, A., Reist, C., & Young, A. S. (2014). Effectiveness of a psychosocial weight management program for individuals with schizophrenia. *The Journal of Behavioural Health Services & Research*, 41(3), 370-380.

Oettingen, G., & Gollwitzer, P.M. (2010). Strategies of setting and implementing goals: Mental contrasting and implementation intentions. In J. E. Maddux & J. P. Tangney (Eds.), *Social Psychological Foundations of Clinical Psychology* (pp. 114-135). New York, NY: Guilford Press.

Parker, G., Arksey, H., & Harden, M. (2010). Meta-review of international evidence on interventions to support carers. York, UK: Social Policy Research Unit.

Pelham, T. W., & Campagna, P. D. (1991). Benefits of exercise in psychiatric rehabilitation of persons with schizophrenia. *Canadian Journal of Rehabilitation*, 4(3), 159-168.

Pesek, M. B., Mihoci, J., Medved, K., & Solinc, N. P. (2011). Long-term groups of patients with psychosis: Physical Activity and Medical Treatment. *Psychiatria Danubina*, 23(1), 149-154.

Prochaska, J. J., Johnson, S., & Lee, P. (2009). The transtheoretical model of behaviour change. In S. A. Shumaker, J. Ockene & K. Riekert (Eds.), *Handbook of Health Behaviour Change* (3rd ed., pp. 59-84). New York, NY: Springer Publishing Company, LLC.

Rastad, C., Martin, C., & Asenlof, P. (2014). Barriers, benefits, and strategies for physical activity in patients with schizophrenia. *Physical Therapy*, 94(10), 1467-1479.

Sailer, P., Wieber, F., Pröpster, K., Stoewer, S., Nischk, D., Volk, F., & Odenwald, M. (2015). A brief intervention to improve exercising in patients with schizophrenia: a

- controlled pilot study with mental contrasting and implementation intentions (MCII). *BMC Psychiatry*, 15(1).
- Sallis, J.F., Cervero, R.B., Ascher, W., Henderson, K.A., Kraft, M.K., & Kerr, J. (2006). An ecological approach to creating active living communities. *Annual Review of Public Health*, 27, 297-322.
- Soundy, A., Freeman, P., Stubbs, B., Probst, M., Coffee, P., & Vancampfort, D. (2014). The transcending benefits of physical activity for individuals with schizophrenia: a systematic review and meta-ethnography. *Psychiatry Research*, 220(1-2), 11-19.
- Stubbs, B., Firth, J., Berry, A., Schuch, F. B., Rosenbaum, S., Gaughran, F., ... Vancampfort, D. (2016). How much physical activity do people with schizophrenia engage in?: A systematic review, comparative meta-analysis and meta-regression. *Schizophrenia Research*. doi:10.1016/j.schres.2016.05.017
- Stubbs, B., Soundy, A., Probst, M., De Hert, M., De Herdt, A., Parker, A., & Vancampfort, D. Stubbs, B., Soundy, A., Probst, M., De Hert, M., De Herdt, A., & Vancampfort, D. (2014a). Understanding the role of physiotherapists in schizophrenia: an international perspective from members of the International Organisation of Physical Therapists in Mental Health (IOPTMH). *Journal of Mental Health*, 23(3), 125-129.
- Stubbs, B., Soundy, A., Probst, M., De Hert, M., De Herdt, A., Parker, A., & Vancampfort, D. (2014b). The assessment, benefits and delivery of physical activity in people with schizophrenia: a survey of members of the International Organization of Physical Therapists in Mental Health. *Physiotherapy Research International*, 19(4), 248-256.
- Stubbs, B., Williams, J., Gaughran, F., & Craig, T. (2016). How sedentary are people with psychosis? A systematic review and meta-analysis. *Schizophrenia Research*, 171(1-3), 103-109.

- Tetlie, T., & Polit, M. C. H. C. (2009). Using exercise to treat patients with severe mental illness: how and why? *Journal of Psychosocial Nursing & Mental Health Services*, 47(2), 32-40.
- Teixeira, P. J., Carraça, E. V., Markland, D. Silva, M. N., & Ryan, R. M. (2012). Exercise, physical activity, and self-determination theory: A systematic review. *International Journal of Behavioral Nutrition and Physical Activity*, 9(78).
- Vancampfort, D., Correll, C. U., Galling, B., Probst, M., De Hert, M., Ward, P. B., Rosenbaum, S., Gaughran, F., Lally, J., & Stubbs, B. (2016). Diabetes mellitus in people with schizophrenia, bipolar disorder and major depressive disorder: a systematic review and large scale meta-analysis. *World Psychiatry*, 15(2), 166-74.
- Vancampfort, D., De Hert, M., Vansteenkiste, M., De Herdt, A., Scheewe, T. W., Soundy, A., Stubbs, B., & Probst, M. (2013). The importance of self-determined motivation towards physical activity in patients with schizophrenia. *Psychiatry Research*, 210(3), 812-818.
- Vancampfort, D., Rosenbaum, S., Schuch, F. B., Ward, P. B., Probst, M., & Stubbs, B. (2016). Prevalence and predictors of treatment dropout from physical activity interventions in schizophrenia: a meta-analysis. *General Hospital Psychiatry*, 39, 15-23.
- Vancampfort, D., Stubbs, B., Mitchell, A. J., De Hert, M., Wampers, M., Ward, P. B., Rosenbaum, S., & Correll, C. U. (2015). Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: a systematic review and meta-analysis. *World Psychiatry*, 14(3), 339-47.
- Vancampfort, D., Sweers, K., Probst, M., Mitchell, A. J., Knapen, J., & De Hert, M. (2011). Quality assessment of physical activity recommendations within clinical practice

guidelines for the prevention and treatment of cardio-metabolic risk factors in people with schizophrenia. *Community Mental Health Journal*, 47(6), 703-710.

Walker, E.R., McGee, R. E., & Druss, B. G. (2015). Mortality in mental disorders and global disease burden implications: a systematic review and meta-analysis. *JAMA Psychiatry*, 72(4), 334-41.

Warren, K. R., Ball, M. P., Feldman, S., Liu, F., McMahon, R. P., & Kelly, D. L. (2011). Exercise program adherence using a 5-kilometer (5K) event as an achievable goal in people with schizophrenia. *Biological Research for Nursing*, 13(4), 383-390.

Wu, M.-K., Wang, C.-K., Bai, Y.-M., Huang, C.-Y., & Lee, S.-D. (2007). Outcomes of obese, clozapine-treated inpatients with schizophrenia placed on a six-month diet and physical activity program. *Psychiatric Services*, 58(4), 544-550.

Figure 1. Flow diagram




Table 1. Research articles included in the scoping review

| Authors | Year | Location | Participants | Setting | Study Design | PA | Knowledge mobilisation Theory | Methods of Info. Dissemination | Messengers |
|------------------------------|-------|-------------|---------------|-----------------------------|-----------------------------------|--|---|---|---|
| Acil et al. | 2008 | Turkey | n=20 | in and out patient hospital | RCT | not specified | not specified | not specified | not specified |
| Attux et al. | 2013 | Brazil | n=160 | out-patients | RCT | walking, moderate and vigorous PA | not specified | education, DVD | mental health professions, DVD, patient relatives |
| Beebe & Faust Harris | 2012 | USA | n=24 | community health centre | feasibility | pedometer use | not specified | verbal instruction | research staff |
| Beebe et al. | 2009 | USA | n=10 | mental health center | intervention | walking | not specified | exercise program | not specified |
| Beebe et al. | 2010 | USA | n=97 | mental health center | RCT | walking | social cognitive theory | verbal, visual aids, demonstration. Forum for discussion | research staff |
| Beebe et al. | 2013 | USA | n=22 | mental health center | follow-up from | walking | social cognitive theory | verbal, visual aids, demonstration. | research staff |
| Behere et al. | 2011 | India | n=56 | not specified | intervention | yoga, exercise | not specified | not specified | not specified |
| Belcher | 1988 | USA | n=1 | nursing home | case study | walking | not specified | walking as a deterrent | staff member |
| Bernard et al. | 2014 | France | n=12 | inpatients | intervention | walking | transtheoretical model | counselling | psychiatric nurse, exercise specialist |
| Bradshaw et al. | 2010 | UK | n=45 | in and out patients | intervention | Self-report PA | MRC framework | discuss problems and solutions | occupational therapists, social worker |
| Chamove | 1986 | UK | n=40 | hospital and hostel | intervention | keep fit, gardening, swimming | not specified | not specified | nurses |
| Chen et al. | 2009 | Taiwan | n=33 | hospital | intervention | aerobic activity | not specified | PA diaries, exercise program | psychiatrist, exercise specialist |
| Crone | 2007 | UK | n=4 | NA | Examine barriers and facilitators | walking | NA | NA | psychiatric nurse, NA |
| Dodd et al. | 2011 | Australia | n=8 | residential unit | intervention | circuit training and walking | not specified | exercise program | exercise psychologist, staff members |
| Duraiswamy et al. | 2007 | India | n=41 | inpatients | intervention | yoga, exercise | not specified | exercise program | trained therapist |
| Faulkner & Sparkes | 1999 | UK | n=3 | hostel | ethnography | walking/swimming | not specified | support | care workers |
| Gomes et al. | 2014 | Portugal | n=19 | sports faculty | intervention | small sided games; basketball, soccer, volleyball, walking handball, jogging | not specified | support, positive feedback | PE teacher, research staff |
| Gorczyński et al. | 2014a | Canada | n=4 | mental health clinic | feasibility | accellerometer use | not specified | accellerometer | research staff |
| Gorczyński et al. | 2014b | Canada | n=4 | mental health clinic | intervention | mixed, self set goals | Self-efficacy theory transtheoretical model Self-determination theory | exercise counseling | research staff |
| Gorczyński et al. | 2013 | Canada | n=25 | in patient | Examine barriers and facilitators | NA | NA | NA | NA |
| Gorczyński, Faulkner, et al. | 2013 | Canada | not specified | mental health unit | intervention | stair use | not specified | point of choice prompts | point of choice prompt |
| Johnstone et al. | 2009 | UK | n=27 | community dwelling patients | examine barriers and facilitators | NA | NA | NA | NA |
| Kwon et al. | 2006 | South Korea | n=48 | clinical centers | RCT | not specified | cognitive and behavioural therapy | exercise diaries, PA discussion | exercise coordinator |
| Leutwyler et al. | 2012a | USA | n=15 | not specified | feasibility | active video games | not specified | video games | research staff |
| Leutwyler et al. | 2012b | USA | n=23 | NA | Examine barrier and facilitators | NA | NA | NA | NA |

Physical activity information

| | | | | | | | | |
|-----------------------------|-------------------------------|------------|---------------------------|--------------------------------------|---|---|---|---|
| Littrell et al. | 2003 USA | n=75 | not specified | intervention | not specified | not specified | verbal and written information. reading aloud, quizzes discussion of topics, and educational games | clinician |
| Maggouritsa et al. | 2014 Greece | n=30 | inpatient hospital | intervention | walking, aerobic exercises, balance and coordination | not specified | exercise program | not specified |
| Marzolini et al. | 2009 Canada | n=13 | community centre | intervention | walking, weight training | not specified | exercise program | Cardiac rehab. specialist nurse, social worker |
| McKibbin et al. | 2012 USA | n=57 | board and care facilities | RCT | self-report PA | not specified | teach and query training methods, mnemonic aids, print materials in large font and limit text | not specified |
| Melamed et al | 2008 USA | n=59 | inpatient hospital | intervention | walking | not specified | exercise program | nurses |
| Methapatara & Srisurapanont | 2011 Thailand | n=64 | inpatient hospital | RCT | walking | motivational interveiwing | leaflets, pedometer | research staff |
| Srisurapanont | | | | | | | | |
| Motlova et al. | 2009 Czech Republic | n=732 | mental health units | intervention | not specified | not specified | education | psychiatric nurses |
| Niv et al. | 2014 USA | n=109 | not specified | intervention | not specified | not specified | handouts, knowledge quizzes, education principles suitable for schizophrenia | nurse care coordinator |
| Pelham & Campagna | 1991 Canada | n=3 | mental health clinic | single subject | bicycle ergometer | not specified | not specified | clinic staff |
| Pelham et al. | 1993 Canada | n=11,10,15 | not specified | 3 comparisons | bicycle ergometer | not specified | exercise program | not specified |
| Pesek et al. | 2011 USA | n=32 | out-patient clinic | intervention | attitudes towards PA | psychoanalytic group approach | psycho education, cognitive techniques, non-structural conversation and clarifications | psychiatrist, psychiatric nurse |
| Rasad et al. | 2014 Sweden | n=20 | out-patient clinic | examine barriers and facilitators | NA | NA | NA | health care professionals |
| Sailer et al. | 2015 Germany & Switzerland | n=36 | inpatients hospital | intervention | jogging | mental contrasting & implementation intentions (MCII) | MCII strategies identify barriers and solutions and goal setting | research and nursing staff |
| Stubbs et al. | 2014 multinational | n=151 | NA | Examine barriers and facilitators | NA | NA | NA | physical therapists |
| Tetlie | 2009 Norway | n=15 | hospital | intervention | gym, swimming, hiking | not specified | exercise program | nurses and exercise instructors |
| Warren et al. | 2010 USA | n=17 | campus research facility | feasibility | walking/jogging | not specified | exercise program | clinical staff, research staff |
| Wu et al | 2015 Taiwan | n=20 | psychiatric day care unit | intervention | HIIT | not specified | exercise program | specialist in internal medicine |
| Wu et al | 2007 Taiwan | n=53 | hospital | RCT | walking/stairs | not specified | exercise program | not specified |

