

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document, Manuscript has been published online as Ahead of Print: Accepted author manuscript version reprinted, by permission, from International Journal of Sports Physiology and Performance, 2020, https://doi.org/10.1123/ijspp.2019-0624. © Human Kinetics, Inc. and is licensed under All Rights Reserved license:

Salter, Jamie, De Ste Croix, Mark B ORCID logoORCID: https://orcid.org/0000-0001-9911-4355, Hughes, Jonathan ORCID logoORCID: https://orcid.org/0000-0002-9905-8055, Weston, Matthew and Towlson, Christopher (2021) Monitoring practices of training load and biological maturity in UK soccer academies. International Journal of Sports Physiology and Performance, 16 (3). pp. 395-406. doi:10.1123/ijspp.2019-0624

Official URL: https://doi.org/10.1123/ijspp.2019-0624 DOI: http://dx.doi.org/10.1123/ijspp.2019-0624 EPrint URI: https://eprints.glos.ac.uk/id/eprint/8306

## Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

# Monitoring practices of training load and biological maturity in UK soccer academies

Jamie Salter<sup>1</sup>, Mark B.A. De Ste Croix<sup>2</sup>, Jonathan D. Hughes<sup>2</sup>, Matthew Weston<sup>3</sup> and Christopher Towlson<sup>4</sup> <sup>1</sup>School of Sport, York St John University, York, UK <sup>2</sup>School of Sport and Exercise, University of Gloucestershire, Gloucester, UK <sup>3</sup>School of Health and Life Sciences, Teesside University, Middlesbrough, UK <sup>4</sup>School of Life Sciences, Sport, Health and Exercise Science. The University of Hull, Hull, UK

Submission Type: Original Investigation

Running Head: Monitoring training load and maturity

Corresponding Author: Jamie Salter School of Sport York St John University Lord Mayors Walk York, YO31 7EX, United Kingdom j.salter@yorksj.ac.uk @jay\_salter ORCiD: 0000-0002-7375-147

Abstract Word Count: 246 Main Text Word Count: 3464 Tables: 4 Figures: 0

### Abstract

## Purpose

Overuse injury risk increases during periods of accelerated growth which can subsequently impact development in academy soccer, suggesting a need to quantify training exposure. Non-prescriptive development scheme legislation could lead to inconsistent approaches to monitoring maturity and training load. Therefore, this study aims to communicate current practices of UK soccer academies towards biological maturity and training load.

#### Methods

Fourty-nine respondents completed an online survey representing support staff from male Premier League academies (n = 38) and female Regional Talent Clubs (n = 11). The survey included 16 questions covering maturity and training load monitoring. Questions were multiple-choice or unipolar scaled (agreement 0-100) with a magnitude-based decision approach used for interpretation.

#### Results

Injury prevention was deemed *highest* importance for maturity (83.0 ± 5.3, mean ±SD) and training load monitoring (80.0 ± 2.8). There were *large* differences in methods adopted for maturity estimation and *moderate* differences for training load monitoring between academies. Predictions of maturity were deemed *comparatively low* in importance for bio-banded (biological classification) training (61.0 ± 3.3) and *low* for bio-banded competition (56.0 ± 1.8) across academies. Few respondents reported maturity (42%) and training load (16%) to parent/guardians, and only 9% of medical staff were routinely provided this data.

# **Conclusions**

Although consistencies between academies exist, disparities in monitoring approaches are likely reflective of environment-specific resource and logistical constraints. Designating consistent and qualified responsibility to staff will help promote fidelity, feedback and transparency to advise stakeholders of maturity-load relationships. Practitioners should consider biological categorisation to manage load prescription to promote maturity appropriate dose-responses and help reduce non-contact injury risk.

Keywords: maturation, training load, monitoring, injury, adolescence, soccer

# Introduction

For academy soccer players, the pubertal growth period is a particularly sensitive time and should be managed with caution<sup>1,2</sup>. This period coincides with progressive, age specific increases in prescribed training exposure (hours), irrespective of individual biological maturation based on the development scheme legislation (policy)<sup>3,4</sup>. Elite Player Performance Pathway (EPPP)<sup>3</sup> and FA Women's Talent Pathway for Regional Talent Clubs (RTC)<sup>4</sup> policy provides recommendations for multifaceted components of player development, including minimum weekly training time, staff requirements, monitoring training load and biological maturity. The systematic increases in training exposure across both genders predominantly reflect development stage informed increases in weekly training load (20-50% depending on academy category) with adolescent players<sup>5</sup>. Most injuries within adolescent soccer are non-contact and soft tissue in nature<sup>6,7</sup> suggesting that these injuries may be attributable to

inadequate training load prescription or growth-related physical and anthropometrical changes<sup>8,9</sup>. Significant time loss through injury, or illness may have major implications for (de)selection and long-term development<sup>10</sup>.

Most (58-69%) injuries within professional soccer academies occur during training rather than match-play. Injuries peak following periods of relatively increased (relative risk of 3.5 following pre-season) or reduced training exposure (mid-season break)<sup>6,11,12</sup>. These findings are consistent with adult populations, where large (>10%) and sudden fluctuations in training load can amplify injury risk<sup>15</sup>. This highlights the importance of quantifying training load to mitigate injury risk<sup>14</sup>, particularly during periods of accelerated biological development<sup>1</sup>. Consequently, to enhance long-term development and improve the sensitivity of (de)selection criteria, fluctuations in physical and functional attributes of players owing to maturity, and the associated response to training exposure, should be monitored and communicated to key stakeholders (e.g. coaches, medical staff and parents/guardians)<sup>15</sup>.

EPPP and RTC policies aim to outline minimum standards for each category to facilitate adequate talent development environments for players. Adherence to these standards are assessed and used to classify each academy (e.g., category 1/tier 1) in return for financial investment and associated prestige helping with recruitment and retention. Yet, the extent of EPPP guidelines is somewhat non-prescriptive and open to interpretation (e.g. '188.2. anthropometric assessments' and '188.7. monitoring of physical exertion [Category 1 academies only]<sup>3</sup>', with no minimum expected monitoring standards or guidelines provided in RTC legislation<sup>4</sup>. Although this ambiguity facilitates context and environment specific approaches which are warranted<sup>16</sup>, it may subconsciously reduce consistency and generate opportunity for '*mixed-practice*' rather than '*best-practice*'.

Various methods to predict maturity status and timing exist with each having logistical, systematic or resource-based confines<sup>17</sup>. Similar limitations exist for training load monitoring which influences the methods adopted by academies<sup>16</sup>. As a result, debate remains around approaches to monitoring training load and which combination of internal (e.g. heart rate, rating of perceived exertion [RPE]) or externally derived metrics (e.g. total distance covered, activity profiles) offer most value for academy practitioners<sup>16</sup>.

Previous surveys investigating training load monitoring have been conducted within professional populations<sup>18,19</sup> and identified varied approaches to collating and disseminating data to stakeholders, with resource and communication-based limitations apparent. Despite strong evidence outlining its relevance within academy settings, no such attempt to investigate current practices of maturity and training load monitoring within male or female academy soccer currently exists. Assessing the current extent of, and manner in which both male and female academies monitor these factors, would provide a platform to develop practice and subsequently optimise development. Therefore, given likely disparities in situational, logistical and environmental factors that govern both male and female academy practices, the aim of the current study was to establish and compare current perceptions and perceived barriers of practitioners to maturity and training load monitoring within UK soccer academies.

# Methods

#### Design

A cross-sectional survey design was used to ascertain perceptions of staff from male (EPPP) and female (RTC) academies during the first trimester (August to December) of the 2017/18 soccer season. Following ethical approval from the University ethics committee and in

accordance with the Declaration of Helsinki, voluntary informed consent was included prior to survey completion. No personal details of the respondent or club were requested to maintain respondent anonymity. Two eligibility questions 1) *Have you already completed the survey?* (Yes or No); 2) *Are you currently working with academy players within an EPPP or RTC setting?* (EPPP, RTC or No) followed the consent page to prevent duplicate responses and ensure construct validity respectively. Each respondent was required to state which professional league their club competed in, the academy category (e.g. Cat/RTC), job role, employment status accompanied by which age category (Foundation [<9 to <12 years], Youth Development [<13 to <16 years], Professional Development [<18 to <23 years]) they primarily worked with.

## **Subjects**

118 respondents started the survey, however, there were 23 incomplete responses and 46 respondents failed eligibility criteria (question 2) and were excluded from analysis. In total, 49 respondents completed the survey (Cat1: n = 15 [31%]; Cat2: n = 13 [27%]; Cat3: n = 10 [20%]; RTC: n = 11 [22%]). Most respondents worked in the Youth Development Phase (YDP; 57%) or Professional Development Phase (PDP; 39%); with 4% working with the Foundation Phase (FP). Most responses were from sport science support staff (sport scientists, strength and conditioning coaches, athletic development or physical development coaches; 77%) with medical (physiotherapists, sports therapists, rehabilitation specialist or doctor; 15%) and technical coaching staff (lead or age group coach; 8%) providing the remainder of the responses. Most of the respondents were employed either full-time (57%) or part-time (23%), with a smaller number of responses coming from sessional staff (hourly paid; 14%) and internship students (6%). Most respondents worked for Championship (43%) or Premier

League (29%) clubs, but some responses were from League One (14%), League 2 (6%) and clubs within the National League or below (8%).

# Methodology

Content validity<sup>20</sup> of the initial survey was reviewed via communications between the research team and practitioners (n = 5) and academics (n = 4) with experience of academy soccer and survey-based studies. This process removed five questions, combined six questions into three and had language amendments for clarity. The final survey consisted of 16 questions that included 2 unipolar (0 = not *important*; 100 = highly *important*) and 6 multiple choice questions each, covering two concepts: 1) *monitoring of biological maturity* and 2) *training load monitoring*. Response analysis to establish internal consistency of each concept using Cronbach's alpha<sup>21</sup> yielded alphas rated as 'good', which ranged from 0.78 [95% confidence interval 0.72 to 0.86] (*monitoring of biological maturity*) to 0.83 [0.72 to 0.86] (*training load monitoring*). The survey was then published using an online survey tool (surveymonkey.com, California, Palo Alto, USA), with completion time of ~10 minutes. A web-link invite to participate was distributed to coaches, sport science support staff and medical practitioners within EPPP and RTC clubs via personal networks and social media.

# Statistical Analysis

Responses from the multiple-choice questions were converted into a proportion of the total number of respondents from each academy category. Independent-group proportion differences for multiple choice questions were calculated with the following scale used to classify magnitudes of difference 10%, 30%, 50%, 70% and 90% as *small, moderate, large, very large* and *extremely large* respectively<sup>22</sup>. Given the small sample size and the large number of inferences, we elected to use moderate as our threshold for meaningful differences.

Numerical data from unipolar-scaled questions were rank ordered and presented as mean ±SD to qualitatively illustrate perceived importance. To facilitate distribution-based interpretations and overcome the limitations of few verbal anchors on the unipolar scale, four perception levels were devised based on percentage thresholds of the overall mean; lowest (<25%), comparatively low (25% to 50%), comparatively high (50% to 75%) and highest (>75%<sup>23</sup>). Inferential analysis (ANOVA) was conducted using JASP computer software (v0.11.1, Amsterdam, Netherlands) to establish independent group mean differences in perceived importance and 99% compatibility limits (CL) to reduce inferential error rates, which were subsequently translated into probabilistic terms using a customised Magnitude-Based Decisions (MBD) spreadsheet<sup>24</sup>. A clear standardised difference for non-clinical substantiveness of 10% was adopted, as this is considered the smallest important effect threshold for between-group differences<sup>22</sup>. Only those effects that were above the smallest important effect were reported and these were then interpreted against the following Bayesian scale: 0.5% most unlikely or almost certainly not; 0.5-5% very unlikely; 5-25% unlikely or possibly not; 25-75% possibly; 75-95% likely or probably; 95-99% very likely; and 99.5% most *likely*<sup>24</sup> to express uncertainty. For both approaches to analysis, all comparisons were made against EPPP Cat1 academies. In light of the EPPP infrastructure being more mature than RTC, and these Cat1 academies fulfilling significant requirements to be awarded this status, they should be regarded as the benchmark of best practice within UK academy football.

### Results

Table 1 Perceived importance (mean  $\pm$  SD) of biological maturity estimations between clubs sorted by percentiles (sample mean  $\pm$  SD), with chances that the true magnitude of difference is important. Effects below the smallest important threshold are not reported. All comparisons made against Category 1 academies (Cat1).

|                                       | 0.1            | G 12          | G 12          | DTC          | 11          |  |
|---------------------------------------|----------------|---------------|---------------|--------------|-------------|--|
|                                       | Catl           | Cat2          | Cat3          | RIC          | Mean        | Between-group differences and probability of                         |
|                                       | (n - 15)       | (n - 12)      | (n - 10)      | (n - 11)     | (n - 40)    | Important differences  |
|                                       | (n = 15)       | (n = 13)      | (n = 10)      | (n = 11)     | (n = 49)    | Mean difference ±99% CL  |
| Perceived level of importance         | e of the estin | nations of bi | iological ma  | uturity for  |             |  |
| <sup>H</sup> injury prevention        | 79 ± 13        | 84 ± 19       | 79 ± 11       | 91 ± 10      | $83 \pm 14$ | <i>Possibly</i> , RTC 11%; ±11%                                      |
| <sup>CH</sup> overall player          | $74 \pm 15$    | $87 \pm 14$   | $80 \pm 12$   | $80 \pm 12$  | $80 \pm 14$ | <i>Possibly</i> , Cat3 6%; ±15%                                      |
| development                           |                |               |               |              |             |  |
| <sup>CH</sup> load management         | $79 \pm 10$    | $79 \pm 20$   | $75 \pm 12$   | $80 \pm 21$  | $78 \pm 16$ |  |
| CHcoach feedback                      | $75 \pm 11$    | $80 \pm 12$   | $72 \pm 9$    | $76 \pm 10$  | $76 \pm 11$ |  |
| <sup>CH</sup> player feedback         | $58\pm18$      | $73 \pm 19$   | $72 \pm 14$   | $81 \pm 14$  | $71 \pm 19$ | Likely, Cat2 15%; ±17%; Cat3 14%; ±18%; Very Likely,                 |
|                                       |                |               |               |              |             | 23%; ±19%  |
| <sup>CL</sup> player retention        | $72 \pm 13$    | $78 \pm 22$   | $64 \pm 22$   | $59 \pm 19$  | $68 \pm 19$ | <i>Possibly</i> , Cat3 -8%: ±21%; RTC -13%; ±20%                     |
| <sup>CL</sup> reports to parents      | 64 ± 13        | 75 ± 22       | $56 \pm 22$   | 75 ± 19      | 68 ± 17     | <i>Possibly</i> , Cat2 11%; ±16%; Cat3 -8%; ±17%; RTC 11%; ±16%      |
| <sup>CL</sup> player recruitment      | $71 \pm 16$    | $71 \pm 22$   | $67 \pm 17$   | $58 \pm 24$  | $67 \pm 20$ | <i>Possibly</i> , RTC -14%; ±21%                                     |
| <sup>CL</sup> bio-banded training     | $59\pm27$      | $64 \pm 23$   | $57 \pm 21$   | $63 \pm 22$  | $61 \pm 23$ |  |
| <sup>L</sup> club legislation         | $54 \pm 17$    | $60 \pm 24$   | $51 \pm 26$   | $64 \pm 15$  | $58 \pm 21$ |  |
| <sup>L</sup> bio-banded competition   | $53 \pm 28$    | 57 ± 32       | 55 ± 23       | 57 ± 21      | $56 \pm 26$ |  |
| <sup>L</sup> EPPP/RTC legislation     | $59\pm15$      | $50 \pm 28$   | $39 \pm 25$   | $52 \pm 26$  | $50 \pm 23$ | <i>Likely</i> , Cat3 -20%; ±23%                                      |
| What are the primary barrier          | s to implem    | enting estim  | ations of bio | ological mat | urity?      | ·  |
| <sup>CH</sup> time constraints        | $57 \pm 23$    | $65 \pm 33$   | $73 \pm 28$   | $66 \pm 26$  | $65 \pm 27$ | <i>Possibly</i> , Cat3 16%; ±29%                                     |
| <sup>CH</sup> staffing numbers        | $47 \pm 27$    | $42 \pm 35$   | $76 \pm 33$   | $47 \pm 32$  | $53 \pm 33$ | <i>Likely</i> , Cat3 29%; ±34%                                       |
| <sup>CH</sup> resource limitations    | $30 \pm 19$    | $31 \pm 26$   | $59 \pm 29$   | $45 \pm 33$  | $41 \pm 28$ | <i>Possibly</i> , RTC 15%; ±28%; <i>Very Likely</i> , Cat3 29%; ±29% |
| <sup>CH</sup> staffing competency     | $41 \pm 26$    | $37 \pm 28$   | $32 \pm 26$   | $53 \pm 32$  | $41 \pm 28$ | <i>Possibly</i> , RTC 12%; ±29%                                      |
| <sup>CL</sup> coach support           | $37\pm26$      | $38 \pm 35$   | $42 \pm 27$   | $31 \pm 23$  | $37 \pm 28$ |  |
| <sup>CL</sup> financial budget        | $25 \pm 24$    | $30 \pm 31$   | $53 \pm 37$   | $35 \pm 27$  | $36 \pm 31$ | Possibly, Cat2 5%; ±30%; RTC 10%; ±32%; Likely, Cat3                 |
| limitations                           |                |               |               |              |             | 28%; ±33%  |
| <sup>CL</sup> management support      | $36 \pm 28$    | $36 \pm 32$   | $35 \pm 26$   | $26 \pm 21$  | $33 \pm 27$ | <i>Possibly</i> , RTC - 10%; ±29%                                    |
| <sup>CL</sup> Parent/guardian support | $17 \pm 16$    | $26 \pm 32$   | $27 \pm 22$   | $29 \pm 30$  | $25 \pm 25$ | Possibly, Cat3 10%; ±28% RTC 12%; ±27%                               |

Perceived importance: 0 = not important, 100 = highly important; Perception level: Llowest; CL comparatively low; CH comparatively high; H highest

Probability of important differences: <0.5%, most unlikely; 0.5-5%, very unlikely; 5-25%, unlikely; 25-50%, possibly; 75-95%, likely; 95-99.5%, very likely; >99.5% most

likely (Hopkins, 2019)

Cat1, Category 1 academy; Cat2, Category 2 academy, Cat3, Category 3 academy; RTC, Regional Talent Club.

#### **Biological Maturity**

Injury prevention was identified as *highest* importance for estimation of maturity across academy groups, with overall athletic development, load management, coach and player feedback considered *comparatively high* (Table 1). Legislative expectations from clubs and governing bodies as well as bio-banded competition were considered *lowest* importance. Cat1 academies placed more importance on EPPP legislation than Cat3 academies and a *likely* to *very likely* lower importance on player feedback than all other academies. Time constraints,

staff numbers, resource limitations and staff competency were all perceived to be *comparatively higher* barriers to implementing maturity predictions (Table 1). Staff numbers and resource limitations are *likely* to *very likely* bigger barriers in lower ranked academies than Cat1. Coach support, financial budget limitations, management and parental/guardian support were all perceived as *comparatively low* barriers, with differences between Cat1, Cat3 and RTC academies *possible* to *likely*.

Table 2 Number of responses (percentages) and qualitative differences magnitude for questions relating to biological maturation estimations. All comparisons made against Category 1 academies (Cat1) with only magnitudes of Small or greater reported.

| Question and Responses   | Cat1             | Cat2             | Cat3             | RTC              | Proportion Difference        |  |  |  |
|--|------------------|------------------|------------------|------------------|------------------------------|--|--|--|
| _  | ( <i>n</i> = 15) | ( <i>n</i> = 13) | ( <i>n</i> = 10) | ( <i>n</i> = 11) | Magnitude                    |  |  |  |
| Which approach is primarily adapted for estimating biological maturity?        |                  |                  |                  |                  |                              |  |  |  |
| Predication of adult height  | 9 (60)           | 1 (8)            | 6 (60)           | 5 (46(           | Small: RTC; Large: Cat2      |  |  |  |
| Maturity offset  | 5 (33)           | 12 (92)          | 3 (30)           | 3 (27)           | Large: Cat2                  |  |  |  |
| Skeletal maturity  | 0 (0)            | 0 (0)            | 0 (0)            | 2 (18)           | Small: RTC                   |  |  |  |
| Other  | 1 (7)            | 0 (0)            | 1 (10)           | 1 (9)            |                              |  |  |  |
| Who is primarily responsible for collecting biological maturation data?        |                  |                  |                  |                  |                              |  |  |  |
| Medical staff  | 1 (7)            | 2 (15)           | 0 (0)            | 3 (28)           | Small: RTC                   |  |  |  |
| Sport Science support staff  | 14 (93)          | 11 (85)          | 8 (80)           | 8 (72)           | Small: Cat3; RTC             |  |  |  |
| Other  | 0 (0)            | 0 (0)            | 2 (20)           | 0 (0)            | Small: Cat3                  |  |  |  |
| *Who is biological maturity data reported to?                                  |                  |                  |                  |                  |                              |  |  |  |
| Academy manager  | 10 (67)          | 8 (62)           | 7 (70)           | 6 (55)           |                              |  |  |  |
| Lead age group coach   | 12 (80)          | 12 (92)          | 8 (80)           | 9 (82)           | Small: Cat2                  |  |  |  |
| Age group coaches  | 14 (93)          | 10 (77)          | 7 (70)           | 9 (82)           | Small: Cat2, Cat 3, RTC      |  |  |  |
| Medical staff  | 15 (100)         | 11 (85)          | 9 (90)           | 9 (82)           | Small: Cat2, Cat3, RTC       |  |  |  |
| Sport Science support staff  | 14 (93)          | 12 (92)          | 9 (90)           | 9 (82)           | Small: RTC                   |  |  |  |
| Intern/student   | 2 (13)           | 6 (46)           | 2 (20)           | 2 (18)           | Large: Cat2                  |  |  |  |
| Player   | 7 (47)           | 5 (39)           | 5 (50)           | 7 (64)           | Small: RTC                   |  |  |  |
| Parent/guardian  | 1 (7)            | 5 (39)           | 4 (40)           | 9 (82)           | Moderate: Cat2, Cat3; Very   |  |  |  |
|  |                  |                  |                  |                  | large: RTC                   |  |  |  |
| What is the primary method of feedback on biological maturation estimations?   |                  |                  |                  |                  |                              |  |  |  |
| Infographic  | 1 (7)            | 0 (0)            | 0 (0)            | 0 (0)            |                              |  |  |  |
| Verbal communication   | 1 (7)            | 2 (15)           | 1 (10)           | 8 (73)           | Large: RTC                   |  |  |  |
| Visual presentation  | 9 (60)           | 8 (62)           | 6 (60)           | 2 (18)           | Moderate: RTC                |  |  |  |
| Written report   | 4 (27)           | 3 (23)           | 3 (30)           | 1 (9)            | Small: RTC                   |  |  |  |
| *When using biological maturity to group players, what activities is this for? |                  |                  |                  |                  |                              |  |  |  |
| Pitch-based sessions   | 8 (25)           | 8 (29)           | 4 (25)           | 2 (25)           | Small: Cat3; Moderate: RTC   |  |  |  |
| Gym-based sessions   | 7 (22)           | 8 (29)           | 4 (25)           | 4 (50)           | Small: Cat2, RTC             |  |  |  |
| Recovery sessions  | 0 (0)            | 0 (0)            | 0 (0)            | 1 (12.5)         |                              |  |  |  |
| Competitive fixtures   | 5 (16)           | 2 (7)            | 1 (6)            | 0 (0)            | Small: Cat2, Cat3; Moderate: |  |  |  |
|  |                  |                  |                  |                  | RTC                          |  |  |  |
| Ad-hoc fixtures  | 7 (22)           | 6 (21)           | 3 (19)           | 1 (12.5)         | Small: Cat3; Moderate: RTC   |  |  |  |
| Specific fixtures  | 5 (16)           | 4 (14)           | 4 (25)           | 0 (0)            |                              |  |  |  |

\*Question permitted multiple responses

Scale of magnitudes: <10%, trivial; 10-30%, small; 30-50%, moderate; 50-70%, large, 70-90%, very large; >90%, huge22

Cat1, Category 1 academy; Cat2, Category 2 academy, Cat3, Category 3 academy; RTC, Regional Talent Club.

There were *large* differences between the methods of maturity estimation utilised by Cat1 and Cat2 academies (Table 2). Cat1, 3 and RTC academies preferred the prediction adult height whist Cat2 had a clear preference for maturity offset (i.e. time from peak height velocity). Sport Science support staff were primarily responsible for collection of maturity data consistently across all academies. There were no small to large differences in the methods used by academies communicate maturity feedback and *moderate* to *very large* differences suggesting that fewer Cat1 academies report this data to parents/guardians. There were small to moderate differences that suggests that academy status is linked to the activities influenced by maturity status monitoring (i.e. pitch-based training, competitive fixtures etc).

|   | Cat1             | Cat2             | Cat3             | RTC              | Mean        | Between-group differences and <i>probability</i> of important differences |
|---|------------------|------------------|------------------|------------------|-------------|---|
|   | ( <i>n</i> = 15) | ( <i>n</i> = 13) | ( <i>n</i> = 10) | ( <i>n</i> = 11) | (n = 49)    | Mean difference ±99% CL   |
| Perceived level of importance for monit                                 | toring train     | ing load for     |                  |                  |             |   |
| <sup>H</sup> injury prevention  | 80 ± 17          | 80 ± 24          | $77 \pm 16$      | $84\pm19$        | $80 \pm 19$ |   |
| <sup>CH</sup> coach feedback  | $80 \pm 10$      | $72 \pm 26$      | $74 \pm 7$       | $66 \pm 21$      | $73 \pm 19$ | <i>Possibly</i> , RTC -14%; ±19%  |
| <sup>CH</sup> prescription of training                                  | $72 \pm 18$      | $70 \pm 17$      | $61 \pm 23$      | $80\pm9$         | $71 \pm 19$ | <i>Possibly</i> , Cat3 -14%; ±20%   |
| <sup>CH</sup> individualisation of training                             | $71\pm18$        | $65 \pm 21$      | $71 \pm 10$      | $77 \pm 13$      | $71 \pm 17$ |   |
| <sup>CH</sup> overall player development                                | $75\pm18$        | $65 \pm 25$      | $73 \pm 12$      | $68 \pm 20$      | $70 \pm 20$ | <i>Possibly</i> , Cat2 -10%; ±20%   |
| <sup>CH</sup> systematic progression                                    | $66 \pm 22$      | $68 \pm 15$      | $68 \pm 15$      | $63 \pm 21$      | $66 \pm 21$ |   |
| <sup>CL</sup> player feedback   | $62 \pm 21$      | $52\pm26$        | $69 \pm 10$      | $72 \pm 7$       | $64 \pm 20$ | <i>Possibly</i> , Cat2 -10%; ±19%   |
| CLEPPP/RTC legislation  | $57 \pm 22$      | $44 \pm 26$      | $53 \pm 13$      | $47\pm28$        | $50 \pm 24$ | <i>Likely</i> , Cat2 -13%; ±24%   |
| <sup>CL</sup> player retention  | $45 \pm 26$      | $44 \pm 25$      | $57 \pm 24$      | $48 \pm 25$      | $49 \pm 25$ | <i>Possibly</i> , Cat3 12%; ±28%  |
| <sup>CL</sup> Parent/guardian feedback                                  | $32 \pm 18$      | $47 \pm 31$      | $51 \pm 15$      | $56\pm21$        | $47 \pm 24$ | <i>Likely</i> , Cat2 15%; ±23%; Cat3 19%; ±25%;                           |
|   |                  |                  |                  |                  |             | RTC 24%; ±24%   |
| <sup>CL</sup> club legislation  | $48\pm19$        | $39 \pm 21$      | $50 \pm 13$      | $45\pm27$        | $46 \pm 21$ |   |
| <sup>CL</sup> player recruitment  | $45 \pm 26$      | $27 \pm 23$      | $44 \pm 25$      | $40\pm28$        | $39\pm26$   | <i>Possibly</i> , Cat2 -18%; ±26%   |
| What are the primary barriers to implementing training load monitoring? |                  |                  |                  |                  |             |   |
| <sup>CH</sup> resource limitations                                      | $54 \pm 34$      | 64 ± 29          | $84 \pm 24$      | $80 \pm 9$       | $71 \pm 32$ | Possibly, Cat2 10%; ±31%; Likely, Cat3                                    |
|   |                  |                  |                  |                  |             | 30%; ±34%   |
| <sup>CH</sup> staffing numbers  | $59\pm28$        | $69\pm28$        | $80\pm26$        | $63\pm29$        | $67\pm28$   | Possibly, Cat2 10%; ±28%; Likely, Cat3                                    |
|   |                  |                  |                  |                  |             | 21%; ±31%   |
| <sup>CH</sup> financial budget limitations                              | $57 \pm 31$      | $72\pm29$        | $82\pm18$        | $50\pm31$        | $65\pm30$   | Possibly, Cat2 15%; ±29%; Likely, Cat3                                    |
|   |                  |                  |                  |                  |             | 25%; ±31%   |
| <sup>CL</sup> limited opportunity for intervention                      | $48 \pm 26$      | $69 \pm 33$      | $63\pm28$        | $53\pm28$        | $58\pm29$   | <i>Possibly</i> , Cat3 15% ±32%; <i>Likely</i> , Cat2 2%;                 |
|   |                  |                  |                  |                  |             | 1 ±29%  |
| <sup>CL</sup> staffing competency                                       | $38 \pm 28$      | $43 \pm 27$      | $44 \pm 24$      | 55 ± 32          | $45 \pm 28$ | <i>Likely</i> , RTC 17%; ±30%   |
| <sup>CL</sup> coach support   | $31 \pm 20$      | $51 \pm 38$      | $37 \pm 24$      | $42 \pm 26$      | $40 \pm 28$ | <i>Possibly</i> , Cat3 6%; ±30%; RTC 11%;                                 |
|   |                  |                  |                  |                  |             | ±30%; <i>Likely</i> , 20%; ±28%   |
| <sup>CL</sup> management support  | $43 \pm 28$      | $39 \pm 38$      | $34 \pm 25$      | $30 \pm 22$      | $36 \pm 29$ | <i>Possibly</i> . Cat3 9%: ±32%: RTC 13%: ±32%                            |

Table 3 Perceived importance (mean  $\pm$  SD) of training load monitoring between clubs sorted by percentiles (sample mean  $\pm$  SD), with chances that the true magnitude of difference is important. Effects below the smallest important threshold are not reported. All comparisons made against Category 1 academies (Cat1).

Perceived importance: 0 = not important, 100 = highly important; Perception level: <sup>L</sup> lowest; <sup>CL</sup> comparatively low; <sup>CH</sup> comparatively high; <sup>H</sup> highest

Probability of important differences: <0.5%, most unlikely; 0.5-5%, very unlikely; 5-25%, unlikely; 25-50%, possibly; 75-95%, likely; 95-99.5%, very likely; >99.5%, most likely (Hopkins, 2019)

Cat1, Category 1 academy; Cat2, Category 2 academy; Cat3, Category 3 academy, RTC, Regional Talent Club

# **Training Load**

Monitoring training load is deemed *highest* importance for injury prevention (Table 3). Player recruitment, retention, parent/guardian and player feedback and legislative purposes were considered *comparatively low* importance. Responses suggest Cat 1 academies *likely* share load monitoring information with parent/guardians less often than other academies.

Resource limitations, staffing numbers, financial budget limitations and limited intervention opportunity were all considered *comparatively high* barriers to training load monitoring (Table

3). Cat3 academies *likely* find these barriers more prominent than Cat1. Management and coach support, staff competency and limited opportunity for intervention were *comparatively low* barriers to training load monitoring. A *possible to likely* differences in coach support may infer greater coach buy-in within Cat1 academies than others. Additionally, it is *likely* that RTC academies perceived staff competency as a greater barrier than Cat1 academies.

*Moderate* differences suggest that Cat1 academies utilise RPE and coach perception less than other academies in preference for external training load measures (Table 4). *Small to moderate* differences suggest that Cat1 academies favour customised spreadsheets to the Performance Management Application (PMA), conversely it is worth noting that the PMA is not available for RTC academies which likely influenced between-group comparisons. Training load data was mostly collated by Sport Science support staff with *moderate* differences between Cat1 and RTC academies. *Moderate* differences suggest Cat1 academies report training load data to age group coaches more frequently than other academies, but less to lead age group coaches than Cat2 academies.

| Question and Responses   | Cat1     | Cat2     | Cat3     | RTC      | Proportion Difference Magnitudes |  |  |
|--|----------|----------|----------|----------|----------------------------------|--|--|
|  | (n = 15) | (n = 13) | (n = 10) | (n = 11) |                                  |  |  |
| What is the primary approach to training load monitoring?      |          |          |          |          |                                  |  |  |
| GPS devices  | 7 (47)   | 4 (31)   | 0 (0)    | 0 (0)    | Small: Cat2; Moderate: Cat3, RTC |  |  |
| Rating of Perceived Exertion                                   | 6 (40)   | 3 (23)   | 7 (70)   | 8 (73)   | Small: Cat2; Moderate: Cat3; RTC |  |  |
| Physiological (TRIMP)  | 1 (7)    | 0 (0)    | 0 (0)    | 0 (0)    |                                  |  |  |
| Coach perceptions  | 1 (7)    | 4 (31)   | 2 (20)   | 1 (9)    | Small: Cat2, RTC                 |  |  |
| Support staff perceptions                                      | 0 (0)    | 0 (0)    | 1 (10)   | 0 (0)    | Small: Cat3                      |  |  |
| Wellness data  | 0 (0)    | 0 (0)    | 0 (0)    | 2 (18)   | Small: RTC                       |  |  |
| Verbal discussion  | 0 (0)    | 2 (15)   | 0 (0)    | 0 (0)    | Small: Cat2                      |  |  |
| How is your training load data compiled?                       |          |          |          |          |                                  |  |  |
| Player Management Application                                  | 4 (27)   | 4 (31)   | 5 (50)   | 0 (0)    | Small: Cat2, RTC                 |  |  |
| Customised spreadsheet   | 9 (60)   | 8 (62)   | 3 (30)   | 9 (82)   | Small:RTC; Moderate: Cat3        |  |  |
| Monitoring application   | 1 (7)    | 0 (0)    | 0 (0)    | 1 (9)    |                                  |  |  |
| Other  | 1 (7)    | 1 (8)    | 2 (20)   | 1 (9)    | Small: Cat3                      |  |  |
| Who is primarily responsible for collating training load data? |          |          |          |          |                                  |  |  |
| Academy manager  | 0 (0)    | 0 (0)    | 1 (10)   | 0 (0)    | Small: Cat3                      |  |  |
| Lead age group coach   | 0 (0)    | 1 (7)    | 1 (10)   | 1 (9)    | Small: Cat3                      |  |  |
| Age group coaches  | 0 (0)    | 1 (7)    | 0 (0)    | 1 (9)    |                                  |  |  |
| Medical staff  | 0 (0)    | 1 (7)    | 1 (10)   | 2 (18)   | Small: Cat3, RTC                 |  |  |
| Sport Sciences support staff                                   | 14 (93)  | 9 (69)   | 7 (70)   | 6 (55)   | Small: Cat2, Cat3; Moderate: RTC |  |  |
| Intern/student   | 1 (7)    | 1 (7)    | 0 (0)    | 1 (9)    |                                  |  |  |
| Players  | 0 (0)    | 0 (0)    | 0 (0)    | 0 (0)    |                                  |  |  |
| Who is training load data reported to?                         |          |          |          |          |                                  |  |  |
| Academy manager  | 0 (0)    | 0 (0)    | 2 (20)   | 3 (27)   | Small: Cat3, RTC                 |  |  |
| Lead age group coach   | 4 (27)   | 8 (62)   | 2 (20)   | 0 (0)    | Small: RTC; Moderate: Cat2       |  |  |
| Age group coach  | 8 (53)   | 1 (8)    | 2 (20)   | 4 (36)   | Small: RTC; Moderate: Cat2, Cat3 |  |  |
| Medical staff  | 0 (0)    | 0 (0)    | 0 (0)    | 1 (9)    |                                  |  |  |
| Sport Science support staff                                    | 1 (7)    | 2 (15)   | 1 (10)   | 0 (0)    |                                  |  |  |
| Player   | 1 (7)    | 1 (8)    | 0 (0)    | 1 (9)    |                                  |  |  |
| Other  | 1 (7)    | 1 (8)    | 3 (30)   | 2 (18)   | Small: Cat3, RTC                 |  |  |
| How frequently are training load reports compiled?             |          |          |          |          |                                  |  |  |
| Daily  | 9 (60)   | 6 (46)   | 2 (20)   | 2 (18)   | Small: Cat2; Moderate: Cat3, RTC |  |  |
| Weekly   | 5 (33)   | 2 (15)   | 2 (20)   | 5 (46)   | Small: Cat2, Cat3, RTC           |  |  |
| Monthly  | 0 (0)    | 1 (8)    | 1 (10)   | 1 (9)    | Small: Cat3                      |  |  |
| Quarterly  | 0 (0)    | 0 (0)    | 0 (0)    | 2 (18)   | Small: RTC                       |  |  |
| Bi-annually  | 0 (0)    | 0 (0)    | 1 (10)   | 0 (0)    |                                  |  |  |
| Annually   | 1 (7)    | 0 (0)    | 1 (10)   | 0 (0)    |                                  |  |  |
| Other  | 0 (0)    | 4 (31)   | 3 (30)   | 1 (9)    | Moderate: Cat2                   |  |  |

Table 4 Number of responses (percentages) and qualitative differences magnitude for questions relating to training load monitoring. All comparisons made against Category 1 academies (Cat1) with only magnitudes of Small or greater reported.

*Question permitted multiple responses* 

Scale of magnitudes: <10%, trivial; 10-30%, small; 30-50%, moderate; 50-70%, large; 70-90%, very large; >90%, huge<sup>22</sup>

Cat1, Category 1 academy; Cat2, Category 2 academy; Cat3, Category 3 academy, RTC, Regional Talent Club

## Discussion

This study represents the first attempt to establish perceptions of monitoring of maturity and

training load in UK soccer academies. Given inherent differences between the two constructs,

findings are discussed individually.

## **Biological Maturity**

Practitioners agreed that injury prevention was of *highest* importance for predicting maturity characteristics. Responses indicate that practitioners recognise associations between maturity characteristics and amplified injury risk, and that monitoring maturity positively influences long-term outcomes<sup>1</sup>. Yet, there is disparity concerning protocols employed to predict maturity between academies, with indicators of timing (offset) and status (percentage of predicted adult height) prominent. *'Other'* responses may include a maturity ratio, growth velocity curves or skeletally derived methods (e.g. body dimensions)<sup>25</sup>. Both dominant protocols are advocated by the legislative bodies, however Cat1, Cat3 and RTC academies demonstrated a greater reliance on the prediction of adult height, with C2 favouring maturity offset (Table 2). Their prevalence is likely attributable to the 'non-invasive' and logistically simple algorithm-based protocols, yet evidence has previously outlined limitations in somatic assessment of maturity in comparison with more invasive skeletal protocols<sup>17</sup>. Consequently, it is imperative that practitioners are cognisant of the relevant methodological limitations and accurate (de)selection evaluations.

Despite being pivotal for categorisation, practitioners unanimously perceived maturity prediction of *comparatively low* importance for biologically classified training and *lowest* for competitions. This is perhaps surprising given the recent rise of bio-banded male soccer tournaments supported by the EPPP, in which players are categorised by their current biological maturity<sup>26</sup>. The relative immaturity of the Women's FA Talent Pathway could explain the *comparatively low* importance placed on this by RTC clubs. Bio-banding is largely considered "an alternative method of categorising players, according to maturity status rather

than their chronological age category, with the assumption that this will alleviate (de)selection bias associated with earlier and/or later maturing players."<sup>27</sup>

Bio-banding is a relatively new concept that has until recently traditionally adopted a talent development and selection focus, and therefore the relevance of bio-banding for managing load and injury was possibly overlooked within survey responses. It is reasonable to think that biological constraints within training and match-play would reduce physical variation and help coaches adequately stimulate players to reduce the typically increased injury incidence around biological growth spurts<sup>2,26</sup>. Evidence suggests trends in injury type throughout maturation, with late maturers having more osteochondral disorders and earlier maturers having more tendinopathies<sup>11</sup>. These non-traumatic injuries are largely preventable, which supports that biologically appropriate training prescription may help reduce the incidence of certain injuries through more effective manipulation of intensity. Therefore, practitioners are encouraged to consider the wider benefits of biological categorisation to optimise training load to facilitate biologically relevant content<sup>1</sup>.

Time constraints, resource limitations, staff number and competency were considered as *comparatively high* barriers particularly in lower ranked academies, which could negatively impact validity of maturity predictions, <sup>28</sup>. Even when maturity assessments are stringently controlled, prediction equations can vary 0.1 to 0.2 years between weekly measures<sup>29</sup>. Therefore, anthropometric data collection requires precise measurements to reduce systematic error, which may be compromised in the absence of adequately trained or experienced staff, equipment or time. Whether these data are sport science led as prevalent in the survey, or medical staff led, consistency is paramount to reduce systematic error and thus safeguard data fidelity (i.e. inter-rater reliability)<sup>25</sup>. Importantly, the quality of internal communication between support, medical and technical staff within soccer clubs has been linked with injury

rates and match availability<sup>15</sup>. Therefore, academies that designate responsibility of maturity monitoring to specifically trained staff will likely enhance transfer to positively influence athletic performance and associated caveats (i.e. reduction of injury risk).

There were *moderate* to *very large* differences between the low number of Cat1 respondents reporting maturity data to players and parent/guardians. This is surprising considering Cat1 academies perceive resources as comparatively lower barriers than Cat3 and RTC and therefore likely have better mechanisms to communicate this information effectively. Being transparent with maturity data and informing parent/guardians of the associated transient physical and functional turbulence related to growth, disadvantages (i.e. stress or anxiety) may be alleviated and may even lead to an autonomy supportive bio-psychosocial environment, reducing the likelihood of drop-out or injury<sup>30</sup>. In contrast, failure to involve stakeholders or providing a clear rationale for decision-making has been termed as '*autonomy-thwarting*' behaviour and linked to failed career progression and behavioural disengagement within soccer<sup>31</sup>.

# Training Load monitoring

Injury prevention perceived to be of *highest* importance for monitoring training load within academies. This is likely influenced by recent associations between training exposure and injury in both adult and adolescent populations<sup>32,33</sup>. Despite being of *highest* importance for injury prevention, remarkably almost no medical staff were routinely provided training load data (Table 4). This may suggest a reactive approach to injury management, opposed to a proactive approach whereby medical staff are actively involved in load management decisions. By routinely sharing training load data with medical staff (e.g. multidisciplinary team meetings), a more unified approach could better inform the process and help reduce injury

incidence<sup>15</sup>. This suggests a communication breakdown in lower ranked academies, negating the purpose of monitoring training load and possibly the impact on reducing injury burden<sup>15</sup>.

In addition, responses suggest coach and player feedback, overall development, systematic progression and individualisation and prescription of future training activities were considered of *comparatively high* importance. Although Cat1 academies reported training load to coaches 80% of the time, other academies reported this data to coaches less. On a positive note, this implies that active engagement in training load monitoring is accepted across academies, but the communication, interpretation and application of this appears to be negating impact, likely attributable to the resources available. Although these findings outline reduced impact of monitoring strategies, they correspond with similar conclusions from professional soccer<sup>18,19</sup>. These studies identified coach buy-in and discipline as prominent barriers to the effective impact of training load monitoring, implying that this problem is not an academy-isolated problem. In resolution, academies are encouraged to employ a routine load monitoring strategy enabling consistent collation and interpretation of data in line with context specific and resource appropriate objectives that fit their structure<sup>16</sup>. This should be combined with an education programme to involve all stakeholders and subsequently establish palatable dissemination strategies to enhance its application<sup>16</sup>, potentially supported by a local academic institution.

Cat1 academies utilise external training loads more than other academies, which is unsurprising based on the resource investment associated with this. This potentially explains why other academies (Cat3) perceive staff numbers, financial budgets and resource limitations, as *comparatively high* barriers to training load monitoring. Although microelectromechanical systems (MEMS) may provide a wealth of data, it does not automatically result in better monitoring outcomes as some ambiguity exists around the precision of devices and metrics to

monitor<sup>33</sup>. Research suggests combining internal and external loads offer best practice and better dose-response outcomes<sup>16</sup> to appropriately quantify the magnitude of internal response in light of the external stimulus<sup>32</sup>. This is crucial during periods of accelerated growth, considering likely fluctuations of the dose-response within adolescent soccer.

In the absence of resources to facilitate MEMS, RPE has been shown to be a suitable and valid surrogate gauge of relative psychophysical training intensity<sup>34</sup>. The application of RPE derived training load values are accessible and cost-effective, which may explain the dominant use of this within academies that reported financial and resource barriers (Cat2, Cat3 & RTC). RPE correlates well with physiological and some MEMS derived metrics, and they can be collated retrospectively with suitable validity in adolescent populations, although an approach utilising multiple markers of training load is preferable if resources permit<sup>14,34</sup>.

# Limitations

Although 49 responses are comparative to other soccer surveys ( $n = 19-41^{18,29,35}$ ), it is below that of others ( $n = 182-242^{19}$ ). It is acknowledged responses from the study represent a portion of the population and the opportunity for multiple responses from academies could lead to clustering<sup>19</sup>. The smaller sample size is somewhat negated as responses were from highperformance environments from a finite pool of UK-based academies. From anecdotal estimations, this study includes responses from approximately 38% of registered academies, from which a statistically conservative approach to inference was adopted to minimise false positive risk with power and precision results indicated by the 99% compatibility intervals for smallest important effects only. It is also acknowledged that engagement in this survey is more likely from those academies actively engaged in load and maturity monitoring, which may have influenced findings. Finally, it is noted differences between the more established EPPP and developing FA Women's Talent Pathway academies exist, and that legislations for these pathways may influence differences in responses. However, this survey provides the first comparison between the professional practices of male and female adolescent academies and was therefore considered a novel facet to the study.

## **Practical Applications**

Designating consistent responsibility for data collation to suitably qualified staff may enhance maturity and training load data dependability, engagement and help establish palatable dissemination strategies. Through this more effective feedback loop, academies will promote transparency of data and better inform stakeholders of maturity-load relationships leading to enhanced impact at group and individual levels. This interdisciplinary approach will require a more proactive, and targeted style of monitoring, to facilitate early intervention around accelerated growth periods. Finally, practitioners should consider using biological categorisation to help manage load prescription and maturity appropriate dose-response to help reduce non-contact injury risk.

## Conclusion

Survey responses suggest that routine monitoring of biological maturity and training load is commonplace within adolescent soccer and that clubs adopt monitoring practices to primarily prevent injury. But, resource and environmental constraints create natural diversity around the methodologies and success of the monitoring process which may nullify impact. Without positively impacting player development or reducing injury risk, the monitoring process is futile. Therefore, practitioners are encouraged to identify a context-specific monitoring system that can be reliably and consistently applied and communicated to players, coaches and parent/guardians efficiently.

# Acknowledgements

There are no acknowledgements beyond the author team for this study. The authors report no conflict of interest.

## References

- 1. Johnson, A., Doherty, P.J. and Freemont, A. (2009). Investigation of growth, development, and factors associated with injury in elite schoolboy footballers: prospective. *BMJ*, 338, b490.
- van der Sluis, A., Elferink-Gemser, M.T., Brink, M.S. and Visscher, C. (2014). Importance of Peak height Velocity Timing in Terms of Injuries in Talented Soccer Players. *Int J Sports Med*, 36, 327-332.
- 3. Elite Player Performance Plan. (2011). *Premier League*. English Premier League. Available online: Elite Player Performance Plan
- 4. FA Girls' England Talent Pathway 2016-17. (2016). The Football Association. Available online: FA England Women's Talent Pathway
- Wrigley, R., Drust, B., Stratton, G., Scott, M. and Gregson, W. (2012). Quantification of the typical weekly in-season training load in elite junior soccer players. *J Sports Sci*, 30, 1573-1580.
- Read, P. J., Oliver, J.L., De Ste Croix, M.B.A., Myer, G.D. and Lloyd, R.S. (2017). An audit of injuries in six English professional soccer academies. *J Sports Sci*, 10, 1-7.
- Tears, C., Chesterton, P. and Wijnbergen, M. (2018). The elite player performance plan: the impact of a new national youth development strategy on injury characteristics in a premier league football academy. *J Sports Sci*, e-publication ahead of print. Available from https://www.ncbi.nlm.nih.gov/pubmed/29478360 [Accessed on 29th October 2018].

- 8. Bowen, L., Gross, A.S., Gimpel, M. Li, F. (2016). Accumulated workloads and the acute:chronic workload ratio relate to injury risk in elite youth football players. *Br J Sports Med*, **51**, 452-459.
- 9. Kemper, G.L.J., van der Sluis, A., Brink, M.S. Visscher, C., Frencken, W.G.P. and Elferink-Gemser, M.T. (2015). Anthropometric Injury Risk Factors in Elite-standard Youth Soccer. *Int J Sports Med*, **36**, 1112-1117.
- Myer, G., Jayanthi, N., Difiori, J.P., Faigenbaum, A.D., Kiefer, A.W., Logerstedt, D. and Micheli, L.J. (2015). Sport specialisation Part 1: Does early sports specialisation increase negative outcomes and reduce the opportunity for success in young athletes? *Orthop Surgery*, 7, 437-442.
- 11. Le Gall, F., Carling, C. and Reilly, T. (2007). Biological maturity and injury in elite youth football. *Scandinavian J Med Sci Sports*, **17**, 564-572.
- Le Gall, F., Carling, C., Reilly, T., Vandewalle, H., Church, J. and Rochcongar, P. (2006). Incidence of injuries in Elite French Youth Soccer: A 10-season study. *Am J Sports Med*, **34**, 928-938.
- 13. Gabbett, T.J. (2015). The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med*, **50**, 273-280.
- 14. Fanchini, M., Ferraresi, I., Petruolo, A., Azzalan, A., Ghielmetti, R., Schena, F. and Impellizzeri, F.M. (2017). Is a retrospective RPE appropriate in soccer? Response shift and recall bias. *Sci Med Football*. Available from <u>https://doi.org/10.1080/02640414.2016.1231411</u> [Accessed 29th March 2018].
- 15. Ekstrand, J., Lundqvist, D., Davison, M., D'Hooghe, M. and Pensgaard, A.M. (2018). Communication quality between the medical team and the head coach/manager is associated with injury burden and player availability in elite football clubs. *Br J Sports Med*, e-publication ahead of print. Available from https://bjsm.bmj.com/content/bjsports/early/2018/08/21/bjsports-2018-099411.full.pdf [Accessed 18th January 2019].
- Gabbett, T.J., Nassis, G.P., Oetter, E., Pretorius, J., Johnston, N., Medina, D., Rodas., Myslinski, T., Howells, D., Beard, A. and Ryan, A. (2017). The athlete monitoring cycle. *Br J Sports Med*, **51**, 1451-1452.

- Malina, R.M., Coelho E Silva, M.J., Figueiredo, A.J., Carling, C. and Beunen, G.P. (2012). Interrelationships among invasive and non-invasive indicators of biological maturation in adolescent male soccer players. *J Sports Sci*, **30**, 1705-1717.
- 18. Akenhead, R. and Nassis, G.P. (2016). Training Load and Player Monitoring in High-Level Football: Current Practice and Perceptions. *Int J Sports Phys Perf*, **11**, 587-593.
- 19. Weston, M. (2018). Training load monitoring in elite English soccer: a comparison of practices and perceptions between coaches and practitioners. *Sci Med Football*, **2**, 216-224.
- 20. Stoszkowski, J. and Collins, D. (2016). Sources, topics and use of knowledge by coaches. *J Sports Sci*, 34:9, 794-802.
- 21. Tavakol, M. and Dennick, R. (2011). Making sense of Cronbach's alpha. *Int Journal of Medical Ed*, 2, 53-55.
- Hopkins, W.G. (2010). Linear models and effect magnitudes for research, clinical and practical applications. *Sportscience*, 14, 49-57. <a href="https://www.sportsci.org/2010/wghlinmod.htm">https://www.sportsci.org/2010/wghlinmod.htm</a>
- McCall, A., Carling, C., Nedelec, M., Davison, Le Gall, F., Berthoin, S. and Dupont, G. (2014). Risk factors, testing and preventative strategies for non-contact injuries in professional football: current perceptions and practices of 44 teams from various premier leagues. *B J Sports Med*, 48, 1352-1357.
- 24. Hopkins, W. G., (2019). A spreadsheet for Bayesian Posterior Compatibility Intervals and Magnitude-Based Decisions. *Sportscience*, **23**, 5–7. sportsci.org/2019/bayes.htm
- 25. Malina, R.M., Bouchard, C. and Bar-Or, O. (2004). Growth, Maturation and Physical Activity. Leeds: Human Kinetics.
- 26. Cumming, S., Lloyd, R.S., Oliver, J.L., Eisenmann, J.C., Malina, R.M. (2017). Biobanding in Sport: Applications to Competition, Talent Identification, and Strength and Conditioning of Youth Athletes. *Strength Cond J*, **39**, 34-47.
- 27. Reeves, M.J., Enright, K.J., Dowling, J. and Roberts, S.J. (2018). Stakeholders' understanding and perceptions of bio-banding in junior-elite football training [Pre Print]. To be published in *Soccer Society*. Available from <u>https://www.tandfonline.com/doi/abs/10.1080/14660970.2018.1432384</u> [Accessed 18th March 2018].

- 28. Buchheit, M., and Mendez-Villanueva, A. (2013). Reliability and stability of anthropometric and performance measures in highly-trained young soccer players: effect of age and maturation. *J Sports Sci*, **31**, 1332-1343.
- 29. Towlson, C., Cobley, S., Midgley A. and Lovell, R. (2017). Relative age, maturation and physical biases on position allocation in elite-youth soccer. *Int J Sports Med*, **38**, 201-209.
- Quested, E., Ntoumais, N., Viladrich, C., Haug, E., Ommundsen, Y. and Van hoye, A. (2013). Intentions to drop-out of youth soccer: A test of the basic need's theory among european youth from five countries. *Int J Sport Ex Psy*, **11**, 395-407.
- 31. Gledhill, A., Harwood, C. and Forsdyke, D. (2017). Psychosocial factors associated with talent development in football: A systematic review. *Psych Sport Ex*, **31**, 93-112.
- Jaspers, A., Kuyvenhoven, J.P., Staes, F., Frenken, W.P., Helsen, W.F. and Brink, M.S. (2018). Examination of the external and internal load indicators' association with overuse injuries in professional soccer players. *J Sci Med Sport*, **21**, 579-585.
- 33. Malone, J., Lovell, R. Varley, M.C. and Coutts, A.J. (2017). Unpacking the black box: Applications and Considerations for using GPS devices in Sport. *Int J Sports Phys Perf*, **12**, 18-26.
- 34. Impellizzeri, F.M., Rampinini, E., Coutts, A.J., Sassi, A. and Marcora, S.M. (2004). Use of RPE-based training load in soccer. *Med Sci Sport Ex*, 36, 1042-1047. DOI: 10.1249/01.MSS.0000128199.23901.2F
- 35. Read, P.J., Jimenez, P., Oliver, J.L., and Lloyd, R.S. (2017a). Injury prevention in male youth soccer: Current practices and perceptions of practitioners working at elite English academies. *J Sports Sci*, **36**, 1423-1431.