



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document, This is a pre-copyedited, author-produced version of an article accepted for publication in Military Medicine following peer review. The version of record Panagiotis Pasis, MSc, Brian Hanley, PhD, Konstantinos Havenetidis, PhD, Athanassios Bissas, PhD, Cypriot and Greek Army Military Boot Cushioning: Ground Reaction Forces and Subjective Responses, Military Medicine, Volume 178, Issue 4, April 2013, Pages e493–e497, is available online at: <https://doi.org/10.7205/MILMED-D-12-00432>. and is licensed under All Rights Reserved license:

Pasis, Panagiotis, Hanley, Brian, Havenetidis, Konstantinos and Bissas, Athanassios ORCID: 0000-0002-7858-9623 (2013) Cypriot and Greek army military boot cushioning: ground reaction forces and subjective responses. Military Medicine, 178 (4e). pp. 493-497. doi:10.7205/MILMED-D-12-00432

Official URL: <https://academic.oup.com/milmed/article/178/4/e493/4222857>

DOI: <http://dx.doi.org/10.7205/MILMED-D-12-00432>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/8394>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Cypriot and Greek Army Military Boot Cushioning: Ground Reaction Forces and Subjective Responses

Panagiotis Pasis, MSc*; Brian Hanley, PhD*; Konstantinos Havenetidis, PhD†; Athanassios Bissas, PhD*

*Carnegie Faculty, Carnegie Research Institute, Leeds Metropolitan University, United Kingdom LS6 3QS.

†Human Performance-Rehabilitation Laboratory, Faculty of Physical and Cultural Education, Hellenic Army Academy, Leoforos Vari-Koropiou Avenue, 16673 Vari, Attiki, Greece.

ABSTRACT

Lower limb injuries are a continual and serious issue for military personnel. Such injuries have been associated with the requirement to train in military boots (MBs) and might be offset with commercial insoles. In this study, ground reaction forces were measured in seven male participants wearing running shoes (RS), MBs commonly used by Cypriot and Greek Army personnel, and the MBs with two types of shock-absorbing insole. The participants performed 4-min trials at walking pace ($5 \text{ km}\cdot\text{h}^{-1}$) and running pace ($10 \text{ km}\cdot\text{h}^{-1}$) at a 5% gradient on a treadmill under all four shod conditions. The treadmill incorporated two force plates under its belt, which provided measurements of key kinetic variables. During walking, RS showed significantly lower values for impact peak force ($p < 0.01$), maximum force ($p < 0.05$), and push-off rate ($p < 0.05$) compared with other conditions, although no significant differences were found during running. Although the RS were rated significantly more comfortable than any other condition, neither insole made the MBs more comfortable to wear. With little evidence to support wholesale adoption of insoles in MBs, their use by military personnel can only be recommended on a case-by-case basis.

INTRODUCTION

Overuse injuries of the lower extremities associated with military training are a serious and continual problem, resulting in loss of manpower and training time.¹⁻⁴ The majority of musculoskeletal injuries associated with military training occur at or below the knee.⁵ For example, Havenetidis et al⁴ found that the most common injuries in Hellenic Army Academy recruits were to the ankle and foot. It has been suggested that the typical military boot (MB) worn during training may be a factor in these injuries, due partly to the inadequate cushioning they provide against shock transmission through the tissues of the lower limb.⁶ This is because the main role of MBs is to protect the foot from direct trauma (because of rough terrain, for example)¹ and protect the ankle from inversion injury⁷ rather than providing shock absorbance. However, previous research has suggested that impact forces were decreased in MBs when using an additional insole⁸ and that by using such insoles the incidence of injuries can similarly be decreased.² This is interesting given that athletic footwear and shock-absorbing insoles are often used by the civilian population to try to protect against injury⁹ by reducing the magnitude and rate of loading experienced during walking and running.¹⁰ However, other research has found that there was no benefit gained from using additional insoles in MBs,^{1,11} particularly when running¹² and so their value to Army personnel is still unclear.

Although some studies have taken an epidemiological approach in assessing the role of insoles in reduction of injury risk,^{11,13,14} others have directly measured those factors associated with lower limb injury. For example, in comparing a standard British MB with and without a commercial insole, Dixon¹⁵ used a force plate to measure ground reaction forces (GRFs) from the right foot only during running trials along a 15-m runway. She found that peak impact force and peak rate of loading were both significantly reduced when using the insole. Similar experimental setups were adopted by Dixon et al⁸ and O'Leary et al,¹⁶ but a limitation of measuring kinetic variables in walking or running is the difficulty of obtaining multiple footstrikes. This is because normal gait patterns, and consequently GRF curves, can be affected by participants targeting the force plates rather than walking or running naturally at an appropriate, realistic pace. This drawback can be avoided with the use of an instrumented treadmill with in-dwelling force plates located under the treadmill belt. Such treadmills also have the advantage that running or walking speed can be controlled and multiple steps can be measured during a single trial. These treadmills are not widely available and therefore offer a novel approach to analyze the effects of MBs on GRFs.

Although all Cypriot and Greek men are normally required to attend the Army forces for a period of between 1 and 2 years, few published data exist related to the shock properties of the MB that is used by Cypriot and Greek Army personnel. Army personnel might decide to use commercially available insoles as a means of protecting themselves from injury or pain. However, despite the enormity of the problem as reflected by the high incidence of lower limb injuries during basic training,⁴ no studies have examined the possible beneficial effects of improving shock absorption in MBs in Cypriot and Greek Army personnel. Therefore, the principal aim of this study was to investigate the GRFs generated during walking and running on an instrumented treadmill while wearing RS, MBs commonly used by Cypriot and Greek personnel, and MB with two different shock-absorbing commercial insoles in an attempt to understand the possible internal loading mechanics. The present study aimed to investigate the importance of comfort perception under these footwear conditions and how this information related to GRF data. Because of the employment of the instrumented treadmill, the findings of the study would provide valuable information not only to Cypriot and Greek Army personnel but also to other users of MBs.

METHODS

Participants

Seven healthy young adult male volunteers (24 ± 3 years; 1.73 ± 0.06 m; 79.2 ± 9.4 kg) took part in the study. The participants wore light clothing and were barefoot during the measurement of their anthropometric characteristics. All participants were normally heel strikers, free from injury on the day of testing, and experienced in treadmill running. The study was approved by the University Ethics Committee, and written informed consent was obtained from all participants before participation in the study.

Description of the Boot and Insoles

The standard MB is a rigid boot composed of an upper made of leather and a rubber sole and its mass (individually) is 0.90 kg. This particular MB is used by Cypriot and Greek Army personnel (infantry). Two commercial pairs of insoles from different manufacturers were used in the present study. The specifications of the insoles are presented below:

Insole A An ethyl vinyl acetate (EVA) insole, featuring a forefoot to aid flexibility and increased EVA heel thickness for comfort. It also has a sculpted heel area for support, an antislip texture on the underside for grip, and a toweling top surface for comfort.

Insole B This insole was developed using Sorbothane technology and consists of 100% polyurethane foam for cushioning and covered in breathable polyester fabric to wick away moisture.

Procedure

The research took place at the University Campus (Biomechanics Laboratory), where participants performed a test comprising both walking and running on the h/p/Cosmos Gaitway treadmill (Gaitway, Traunstein, Germany). This treadmill has two in-dwelling force plates with an eightchannel charge amplifier (Kistler, Winterthur, Switzerland), which can measure GRFs during locomotion. Its force range was set to 6,000 N for testing. All participants were given time to familiarize themselves with the treadmill during a separate visit. This was achieved by allowing the participants to walk and run on the treadmill at any desired speed for any period of time.

Each participant performed a warm-up at a steady pace. Afterwards, they performed stretching exercises of their preference. To imitate the kinds of training undertaken using the MBs,³ the participants then started either walking at 5 km·h⁻¹ (1.39 m·s⁻¹) or running at 10 km·h⁻¹ (2.78 m·s⁻¹) on the treadmill, at a 5% gradient, for 4 minutes. Each test was performed under the following conditions:

- Wearing running shoes (RS)
- Wearing the MB without insole (MB)
- Wearing the MB with insole A (MBA)
- Wearing the MB with insole B (MBB)

The order of testing was randomized to imitate the undefined nature of training and between each condition the participants had a rest of 4 minutes. It was decided to conduct each running and walking trial over 4 minutes to minimize any possible influence of fatigue over the course of testing and to minimize any discomfort felt in any particular shod condition. The participants wore their own RS, although the boots and both sets of insoles were newly acquired. Data were collected at 1,000 Hz during the last 30 seconds of walking and running in all four different conditions. This resulted in analysis of between 50 and 60 steps during walking and between 70 and 90 steps during running per participant during each condition. During data collection, the researcher ensured that each participant was striking the treadmill correctly; this was achieved by monitoring the participant's position on the treadmill and by checking that a full complement of force traces were recorded immediately after recording. The treadmill collected data from both left and right footstrikes. The kinetic variables that were collected and investigated included the impact peak force (IPF), the push-off rate (POR), maximum force (MF), and loading rate (LR). IPF was defined as the as highest recorded force recorded during the first 70 ms of contact with the treadmill and represented the passive peak. In conjunction with this, MF was defined as the highest force recorded during the contact phase. LR was defined as the slope of the force curve throughout the loading phase of the running cycle and is taken from the point of 10% of the IPF to the 90% point. POR was defined as the slope of the force curve during unloading, taken from the 90% of push-off peak to the point of 10%. To facilitate comparisons between participants,

GRF peak magnitudes and LRs were divided by the participants' weights and expressed in bodyweights (BW) and BW per second ($BW \cdot s^{-1}$), respectively.

Upon completion of each experimental condition, the participants answered a questionnaire (as described by House et al¹⁷), which asked them to evaluate the comfort of the RS, MBs, and MBs with insoles, by marking a position on a line that ranged from very comfortable (+10) to very uncomfortable (-10).

Statistical Analysis

Statistical analysis of the GRF variables was undertaken using PASW Statistics 18 (IBM SPSS, Chicago, Illinois, 2009). Means and standard deviations were computed for all variables. Analysis of variance and subsequent post hoc analysis (Tukey) were used to determine possible differences between footwear conditions with an α level of 5%. A Friedman test was used to analyze the questionnaire data, and Spearman's rank correlation test was used to examine possible relationships between subjective (questionnaire) and objective (GRF) data.

RESULTS

Analytical data for all variables during walking are presented in Table I. During walking, IPF was lower when participants wore their own RS than when wearing the MBs either with or without the insoles. Furthermore, MF was lower in RS than in the MBA and MBB conditions, but it was not lower in RS than in the MB condition. This was despite there being a larger absolute difference between the RS and MB than between RS and MBA or MBB. This was because of the larger range (and therefore larger standard deviation) found in the MB condition for this variable.

Table 1 GRF data for each condition during walking

Variable	Footwear condition			
	RS	MB	MBA	MBB
IBF (BW)	0.11 ± 0.08 ^a	0.43 ± 0.04 ^a	0.41 ± 0.03 ^a	0.42 ± 0.06 ^a
MF (BW)	1.15 ± 0.04 ^b	1.25 ± 0.12	1.19 ± 0.05 ^b	1.21 ± 0.04 ^b
LR ($BW \cdot s^{-1}$)	7.60 ± 1.19	8.03 ± 0.87	7.86 ± 0.74	8.13 ± 0.89
POR ($BW \cdot s^{-1}$)	10.46 ± 0.50 ^c	10.47 ± 0.86	9.91 ± 0.63 ^c	10.06 ± 0.84

IPF and MF are expressed in BW, and LR and POR are expressed in BW per second ($BW \cdot s^{-1}$). ^aSignificant difference ($p < 0.001$) between RS and MB, MBA, MBB. ^bSignificant difference ($p < 0.01$) between RS and MBA, MBB. ^cSignificant difference ($p < 0.05$) between RS and MBA.

Analytical data for conditions RS, MB, MBA, and MBB during running are shown in Table II; no significant differences were found for any of these variables.

Table 2 GRF data for each condition during running

Variable	Footwear condition			
	RS	MB	MBA	MBB
IBF (BW)	121 ± 0.10	1.22 ± 0.19	1.10 ± 0.10	1.12 ± 0.19
MF (BW)	2.33 ± 0.14	2.40 ± 0.17	2.36 ± 0.13	2.42 ± 0.08
LR ($BW \cdot s^{-1}$)	23.91 ± 4.49	22.48 ± 4.80	21.67 ± 3.35	21.90 ± 2.38
POR ($BW \cdot s^{-1}$)	18.21 ± 2.49	18.43 ± 2.81	18.33 ± 2.36	18.41 ± 2.14

IPF and MF are expressed in BW, and LR and POR are expressed in BW per second ($BW \cdot s^{-1}$)

The subjective comfort/discomfort data for each condition are presented in Figure 1.




Figure 1 Average comfort results for RS and MBs without and with insole A (MBA) and B (MBB). * Indicates a significant difference between RS and MBA ($p < 0.05$) and MBB ($p < 0.05$) conditions, and † indicates a significant difference between MB and RS ($p < 0.001$) conditions.

All participants rated the RS as the most comfortable, and they were significantly different from other conditions (Friedman = 17.4, $p < 0.001$). Mean ranks for RS, MB, MBA, and MBB were 3.9, 1.0, 2.7, and 2.4, respectively. The range of comfort–discomfort scores ($-10 =$ very uncomfortable; $0 =$ neutral; $10 =$ very comfortable) for RS, MB, MBA, and MBB were from 5 to 10, from -10 to 4, from -7 to 8 and from -8 to 7, respectively. The number of participants who rated the RS, MB, MBA, and MBB conditions in the comfortable range (greater than 0) was 7 (100%), 1 (14%), 4 (57%), and 3 (43%), respectively. Alternatively, the number of participants who rated the RS, MB, MBA, and MBB conditions in the uncomfortable range (less than zero) was 0 (0%), 9 (86%), 3 (43%), and 4 (57%), respectively. No significant correlations were found between comfort/discomfort data and GRF data. Spearman rank test correlation values are presented in Table III.

Table 3 Correlation analysis of key GRF variables with comfort scores during both running and walking

GRFs	Comfort/discomfort response			
	RS	MB	MBA	MBB
Walking				
IPF (BW)	-0.06	0.29	0.52	-0.11
MF (BW)	-0.69	-0.46	-0.28	0.25
LR ($BW \cdot s^{-1}$)	-0.13	-0.25	-0.34	0.09
POR ($BW \cdot s^{-1}$)	-0.13	-0.11	0.14	0.11
Running				
IPF (BW)	0	-0.54	-0.02	0.04
MF (BW)	0.24	-0.61	-0.57	-0.22
LR ($BW \cdot s^{-1}$)	-0.35	-0.36	-0.29	0.25
POR ($BW \cdot s^{-1}$)	0.33	-0.50	-0.65	-0.07

DISCUSSION

The aim of this study was to investigate the GRFs generated during walking and running while wearing RS, MBs, and MBs with two different shock-absorbing commercial insoles (MBA and MBB) in an attempt to understand the possible internal loading mechanics. The present data showed that during walking across all conditions, RS presented a lower GRF profile compared with MBs either with or without the shock-absorbing insoles. In particular, IPF was approximately four times greater in the three MB conditions compared with RS. However, despite the absolute values for MF, LR, and POR being higher for MBs than for RS, there were no significant differences between them. The values found for MF in both insole conditions (MBA and MBB) were found to be greater than in RS. The absence of a similar significant difference in the MB condition (despite higher absolute values) might have been due to the larger standard deviation found in the MB condition. Larger standard deviations were also found in the MB condition during the running trials for all GRF variables. These larger deviations in both forms of gait suggest that there is a wider range of individual adaptations to wearing MBs, which means that for some individuals (i.e., those encountering the greatest decrease in GRFs) the insoles might be more worthwhile and have an important benefit.

During the running trials, the study's overall results showed that no significant differences existed across all variables while wearing the running shoe compared with the three boot conditions. This might have been because of the participants adopting different running styles to accommodate the different footwear conditions so that GRFs were minimized. In particular, research suggests that individuals adapt their running style to different shoe-surface interactions¹⁸ so that changes in gait kinematics (e.g., footstrike pattern) occur to reduce impact variables, such as peak impact force, or any pain or discomfort experienced; it is possible that this may have occurred in the MB conditions. Whatever the reason, there was no advantage to either wearing RS or commercial insoles when running in terms of reducing GRF magnitudes. These results differ from those of some previous research, which did find reduced impact forces with some insoles in MBs.^{8,15} The present study measured impact forces using an instrumented treadmill rather than the more commonly used runway methodology as it eliminated the risk of participants targeting the force plate or varying their speed. It is possible that the differences in findings were a result of these different methodologies, and further research using instrumented treadmills is advised.

With regard to ratings of comfort, the findings of the present study suggested that commercially available insoles did not play a significant role in the perception of footwear comfort in the MBs. In particular, the results showed that even with an insole the MBs did not achieve the comfort perception of the RS. This is probably because of the fact that the MB provided extra weight to the foot, is much more rigid, and its general design and construction have other priorities than comfort.^{1,7} On an individual basis, all participants gave higher ratings to the two insole conditions than without the insoles, similar to earlier studies,^{17,19} but there was no significant difference overall and both insole conditions were still rated as uncomfortable by roughly half the participants. This suggests that the shock absorbance properties of the insoles were not sufficient to make the boots comfortable enough for walking and running across all individuals (as the RS were), and this is another aspect of fitting insoles, which needs to be assessed on a case-by-case basis. The low comfort scores

found for the insole conditions might be because of the other properties of the MB, which make it uncomfortable (e.g., its rigidity) and therefore cannot be overcome with an insole alone. The perception of comfort scores showed that all participants preferred walking and running while wearing a running shoe and this suggests that changes in MB design, such as softer leather and wider shoe lasts, could be beneficial.²⁰

The results of the present study indicating that insoles did not have a significant role in comfort perception when wearing the MB contrasts with the findings of other studies,^{19,21} where boots with the best combination of shock-absorbing properties and stability were rated the most comfortable by the participants. However, even in these studies,^{19,21} there was no association between the sensitivity of cushioning and the GRFs, which was supported by the present study's data where comfort ranking was also not related with any of the force variables measured. This would seem to suggest that a boot which feels comfortable does not necessarily have reduced GRF magnitudes, and therefore any risk of injury to the lower limb from impact forces needs to be assessed separately. Although the lack of significance could be partly because of the limited number of participants, we would nonetheless suggest that the perception of comfort itself is not sufficient to provide evidence for the suitability of the insoles and for their promotion among recruits. Furthermore, since the insoles in the present study were rated under a very short-term trial, the clinical value of the present results is not certain; in particular, insoles which do make MBs comfortable on initial usage should be assessed over the course of many months of being worn to measure any depreciation in quality. The participants in the present study were required to run and walk for relatively short periods of time for each condition (4 minutes); this was to avoid fatigue or pain, which could have been exacerbated by performing in unusual shod conditions. However, it is possible that longer bouts of running and walking will provide useful insights into cushioning and comfort variables in MBs, and such future research is recommended.

The loss of manpower because of lower limb injury is a serious issue for professional armies. Basic infantry training has been changed in some armies to reduce the incidence of lower limb injury, for example with reduced marching.²⁰ Nonetheless, army personnel are still required to march often considerable distances on foot while carrying heavy loads. With regard to GRF variables, program modifications may be needed so that long-term hiking with pack and equipment are performed in MBs, which resemble RS in terms of absorbance and comfort properties, rather than by just adding an insole to the MB. However, care must always be taken with new boot design as more comfortable boots are not necessarily better for injury prevention.²² The use of an instrumented treadmill for future studies on MBs is recommended because of the large number of footstrikes quickly available for analysis and the ability to record footstrikes from both right and left feet.

CONCLUSIONS

The present study illustrated that the MBs used by the Cypriot and Greek Army personnel with the use of the two specific insoles did not significantly influence IPFs during walking and running. Participants found the MBs uncomfortable, and this was not significantly offset when using the insoles. Although commanders insist that recruits wear MBs in preparation for and during war and therefore there is a need to train in them beforehand, it should be noted that an injured recruit cannot fight as well as a healthy recruit. Therefore, because of

interindividual differences in GRF patterns, it is worthwhile assessing each recruit on an individual basis for the appropriateness of inserting insoles into MBs.

REFERENCES

1. Davidson PL, Wilson SJ, Chalmers DJ, Wilson BD, McBride D: Examination of interventions to prevent common lower-limb injuries to the New Zealand Defense Force. *Mil Med* 2009; 174(11): 1196–202.
2. Franklyn-Miller A, Wilson C, Bilzon J, McCrory P: Foot orthoses in the prevention of injury in initial military training. *Am J Sports Med* 2010; 39(1): 30–7.
3. Havenetidis K, Paxinos T: Risk factors for musculoskeletal injuries among Greek Army officer cadets undergoing basic combat training. *Mil Med* 2011; 176(10): 1111–6.
4. Havenetidis K, Kardaris D, Paxinos T: Profiles of musculoskeletal injuries among Greek Army officer cadets during basic combat training. *Mil Med* 2011; 176(3): 297–303.
5. Kaufman KR, Brodine S, Shaffer R: Military training-related injuries: surveillance, research, and prevention. *Am J Prev Med* 2000; 18(3S): 54–63.
6. Schwellnus MP, Jordan G, Noakes TD: Prevention of common overuse injuries by the use of shock absorbing insoles. *Am J Sports Med* 1990; 18(6): 636–41.
7. Snyder RA, DeAngelis JP, Koester MC, Spindler KP, Dunn WR: Does shoe insole modification prevent stress fractures? A systematic review. *Hospital Special Surg J* 2009; 5(2): 92–8.
8. Dixon SJ, Waterworth C, Smith C, House CM: Biomechanical analysis of running in military boots with new and degraded insoles. *Med Sci Sports Exerc* 2003; 35(3): 472–9.
9. Razeghi M, Batt ME: Biomechanical analysis of the effect of orthotic shoe inserts. *Sports Med* 2000; 29(6): 425–38.
10. House CM, Waterworth C, Allsopp AJ, Dixon SJ: The influence of simulated wear upon the ability of shock absorbing insoles to reduce peak pressures during running when wearing military boots. *Gait Posture* 2002; 16(3): 297–303.
11. Withnall R, Eastaugh J, Freemantle N: Do shock absorbing insoles in recruits undertaking high levels of physical activity reduce lower limb injury? A randomized controlled trial. *J R Soc Med* 2006; 99(1): 32–7.
12. Ekenman I, Milgrom C, Finestone A, Begin M, Olin C, Arndt T, Burr D: The role of biomechanical shoe orthoses in tibial stress fracture prevention. *Am J Sports Med* 2002; 30(6): 866–70.
13. Larsen K, Weidich F, Leboeuf-Yde C: Can custom-made biomechanical shoe orthoses prevent problems in the back and lower extremities? A randomized, controlled intervention trial of 146 military conscripts. *J Manipulative Physiol Ther* 2002; 25(5): 236–31.
14. Milgrom C, Finestone A, Lubovsky O, Zin D, Lahad A: A controlled randomized study of the effect of training with orthoses on the incidence of weight bearing induced back pain among infantry recruits. *Spine* 2005; 30(3): 272–5.

15. Dixon S: Influence of a commercially available orthotic device on rearfoot eversion and vertical ground reaction force when running in military footwear. *Mil Med* 2007; 172(4): 446 – 50.
16. O’Leary K, Anderson Vorpahl K, Heiderscheit B: Effect of cushioned insoles on impact forces during running. *J Am Podiatr Med Assoc* 2008; 98(1): 36– 41.
17. House CM, Dixon SJ, Allsopp AJ: User trial and insulation test to determine whether shock-absorbing insoles are suitable for use by military recruits during training. *Mil Med* 2004; 169(9): 741– 6.
18. Nigg BM, Yeadon MR: Biomechanical aspects of playing surfaces. *J Sports Sci* 1987; 5(2): 117– 45.
19. Mündermann A, Stefanyshyn DJ, Nigg BM: Relationship between footwear comfort of shoe inserts and anthropometric and sensory factors. *Med Sci Sports Exerc* 2001; 33(11): 1939– 45.
20. Finestone A, Milgrom C: How stress fracture incidence was lowered in the Israeli Army: a 25-yr struggle. *Med Sci Sports Exerc* 2008; 40(11 Suppl): S623–9.
21. Williams KM, Brodine SK, Shaffer RA, Hagy J, Kaufman K: Biomechanical properties of infantry combat boot development. Naval Health Research Center Report No. 97-26, 1997; pp 1–22. The Defense Technical Information Center. Available at <http://www.dtic.mil/dtic/tr/fulltext/u2/a331427.pdf>; accessed August 30, 2012.
22. Arndt A., Westblad P, Ekenman I, Lundberg A: A comparison of external plantar loading and in vivo local metatarsal deformation wearing two different military boots. *Gait Posture* 2003; 18(2): 20 – 6.