UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document, This is the peer reviewed version of the following article: Hegyi, A, Péter, A, Finni, T, Cronin, NJ. Region-dependent hamstrings activity in Nordic hamstring exercise and stiff-leg deadlift defined with high-density electromyography. Scand J Med Sci Sports. 2018; 28: 992–1000. <https://doi.org/10.1111/sms.13016>, which has been published in final form at <https://onlinelibrary.wiley.com/doi/full/10.1111/sms.13016>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving. and is licensed under All Rights Reserved license:

**Hegyi, András, Péter, Annamária, Finni, Taija and Cronin, Neil
ORCID: 0000-0002-5332-1188 (2018) Region-dependent
hamstrings activity in Nordic hamstring exercise and stiff-leg
deadlift defined with high-density electromyography.
Scandinavian Journal of Medicine and Science in Sports, 28
(3). pp. 992-1000. doi:10.1111/sms.13016**

Official URL: <http://dx.doi.org/10.1111/sms.13016>

DOI: <http://dx.doi.org/10.1111/sms.13016>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/8881>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Region-dependent hamstrings activity in Nordic hamstring exercise and stiff-leg deadlift defined with high-density electromyography

András Hegyi | Annamária Péter | Taija Finni | Neil Cronin

Neuromuscular Research Center, Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

Correspondence:

András Hegyi, Neuromuscular Research Center, Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland. Email: andras.a.hegyi@jyu.fi

ORCID:

A. Hegyi <http://orcid.org/0000-0002-3663-0288>

T. Finni <http://orcid.org/0000-0002-7697-2813>

A. Péter <http://orcid.org/0000-0002-5197-6869>

N. J. Cronin <http://orcid.org/0000-0002-5332-1188>

Abstract

Recent studies suggest region-specific metabolic activity in hamstring muscles during injury prevention exercises, but the neural representation of this phenomenon is unknown. The aim of this study was to examine whether regional differences are evident in the activity of biceps femoris long head (BFlh) and semitendinosus (ST) muscles during two common injury prevention exercises. Twelve male participants without a history of hamstring injury performed the Nordic hamstring exercise (NHE) and stiff-leg deadlift (SDL) while BFlh and ST activities were recorded with high-density electromyography (HD-EMG). Normalized activity was calculated from the distal, middle, and proximal regions in the eccentric phase of each exercise. In NHE, ST overall activity was substantially higher than in BFlh ($d = 1.06 \pm 0.45$), compared to trivial differences between muscles in SDL ($d = 0.19 \pm 0.34$). Regional differences were found in NHE for both muscles, with different proximal-distal patterns: The distal region showed the lowest activity level in ST (regional differences, d range = 0.55-1.41) but the highest activity level in BFlh (regional differences, d range = 0.38-1.25). In SDL, regional differences were smaller in both muscles (d range = 0.29-0.67 and 0.16-0.63 in ST and BFlh, respectively) than in NHE. The use of HD-EMG in hamstrings revealed heterogeneous hamstrings activity during typical injury prevention exercises. High-density EMG might be useful in future

studies to provide a comprehensive overview of hamstring muscle activity in other exercises and high-injury risk tasks.

Keywords:

bi-articular hamstrings, electrical activity, muscle function

1 | INTRODUCTION

Hamstring injuries are the most common in sports involving high-speed running.¹⁻³ Optimizing hamstring exercise selection for prevention of and rehabilitation from hamstring injury is of major interest as low eccentric strength is thought to be one of the most significant⁴⁻⁶ but modifiable⁷⁻¹⁰ risk factors for hamstring injury. In recent decades, injury incidence has not decreased,¹¹ highlighting the need for studies examining the possible mechanisms of injury. Bi-articular components of the hamstrings (biceps femoris long head [BFlh], semitendinosus [ST], and semimembranosus) contribute to both hip extension and knee flexion.¹² Thus, complex within-muscle hamstring coordination may be required. Recent studies using muscle functional magnetic resonance imaging (mfMRI) suggest that hamstring exercises may cause heterogeneous elevation of transverse (T2) relaxation time within different hamstring muscles, implying inhomogeneous metabolic activity of different muscle regions.^{13,14} However, there is a lack of information about the neural representation of this phenomenon. Additionally, mfMRI cannot be used in real-time. Therefore, other methods of defining muscle activity are needed to comprehensively understand hamstring muscle function. Electromyography (EMG) has been widely used to define real-time ST and BFlh activity in a variety of hamstring exercises,¹⁵⁻¹⁸ but has relied on a single pair of EMG electrodes placed over each muscle. Assuming that the aforementioned mfMRI changes are reflected in the EMG activity, conventional EMG configurations may not be sufficient to describe overall muscle activity accurately.

High-density surface electromyography (HD-EMG) has been used recently to provide a more comprehensive overview of human muscle activity.¹⁹ This method has revealed regional differences (eg, proximal vs distal) in the activity of different muscles during stimulation and voluntary movements, for example, walking.²⁰⁻²³ The method has not yet been used in hamstrings, but would likely provide further insights into hamstring activity during strengthening exercises. Recent mfMRI studies suggest that hip-dominant exercises may activate proximal BFlh preferentially, while in knee-dominant exercises distal BFlh is relatively more activated,^{14,24} which may be of interest from an injury prevention perspective. Thus in this study, to examine whether heterogeneous activity of hamstring regions exists, two exercises which are frequently used in injury prevention—the knee-

dominant Nordic hamstring exercise (NHE) and the hip-dominant stiff-leg deadlift (SDL)—were investigated with HD-EMG. We aimed to examine the eccentric-only NHE and the eccentric phase of SDL to avoid contraction mode bias. We hypothesized that differences would be observed between normalized EMG activity of the proximal, middle, and distal regions of BFlh and ST during NHE and SDL.

2 | MATERIALS AND METHODS

2.1 | Participants

Twelve recreationally active young males (age 24.3 ± 3.7 years, body mass 74.2 ± 8.3 kg, height 179.3 ± 8.8 cm) with weightlifting experience and without any cardiovascular or musculo-skeletal disorders volunteered for this study. Participants had no known history of hamstring injury or any lower extremity/lower back injuries in the past 3 years. After informing the participants about the study details, they gave written consent before data collection. Testing procedures were approved by the ethics committee of the University of Jyväskylä and performed in accordance with the Declaration of Helsinki.

2.2 | Study design

In the familiarization session (10-14 days before the testing), 1 repetition maximum (1RM) was defined for SDL.²⁵ Participants also practiced the NHE and maximal voluntary isometric contractions (MVICs) with visual force-time curve feedback.

In the main testing session, after preparation and warm-up, participants lay prone in a custom-made dynamometer (UniDrive, University of Jyväskylä)²⁶ with hip joint and trunk fixed to the dynamometer bench in neutral position. For EMG signal normalization, participants performed maximal hip extension and knee flexion MVICs after a specific warm-up including 10 submaximal contractions with increasing intensity (from ~30% to ~90%). Activity along the BFlh and ST muscles was recorded during MVICs using HDEMGE arrays (Figure 1). During knee flexion contractions, the knee joint was fixed at ~20° of flexion while the lever arm of the dynamometer was strapped 2 cm above the lateral malleolus. During hip extension, participant positioning was identical but the lever arm of the dynamometer was fixed 1 cm above the knee fold. Participants were asked to perform hip extension with a slightly flexed knee to match the position in the knee flexion tasks. Three maximal contractions were performed and maintained for 2 seconds for both knee flexion and hip extension (2 minutes rest in-between). Knee flexion or hip extension alone was assumed to be insufficient to evoke maximal muscle activity in all EMG channels. Thus, participants also performed hip extension MVICs superimposed on knee flexion MVICs (3 reps, 2 minutes rest in-between). In

this task, the cuff of the dynamometer lever arm was fixed 2 cm above the lateral malleolus, and the thigh was strapped to the bench above the knee fold. Participants started by increasing knee flexion then adding hip extension, reaching maximal effort in both tasks in ~2 seconds and maintaining it for another 2 seconds. Thereafter, NHE and SDL exercises were performed in random order.

Figure 1 Fifteen-channel high-density electromyography arrays were attached over the biceps femoris long head and semitendinosus to measure activity along the muscles. Electrode grids were filled with conductive gel for proper electrode-skin contact.

2.3 | Exercise description

2.3.1 | Nordic hamstring exercise

The knee-dominant NHE (5 repetitions, 2-minutes rest in-between) was performed on a custom-made device with force transducers attached above the ankles.²⁷ Participants started from a kneeling position, with arms crossed in front of the body. They then lowered the body forward as far as possible at a constant speed of $18^\circ/\text{s}$ controlled with a metronome. The hips and torso were in neutral position throughout the range of motion. NHE was performed as a bodyweight-only exercise, and participants were not able to resist until full knee extension. Force from each leg and EMG activity along the ST and BFLh were recorded during the exercise (Video S1).

2.3.2 | Stiff-leg deadlift

In the hip-dominant SDL, the starting position was upright. Throughout the range of motion, the knees were straight but not locked, and the back was in neutral position with closed scapulae. Participants lowered the bar close to the body toward the floor until the plates touched the floor or for as long as proper technique could be maintained. In the upward movement, the hips were extended to the starting position. The downward and upward movements were each performed in 2 seconds. Five repetitions were performed at 80% 1RM, with 2-minutes rest between repetitions. Joint kinematics and EMG activity along the muscles were recorded during the exercise (Video S2).

2.4 | Data collection

2.4.1 | Electromyography

High-density electromyography preparation was performed while participants lay prone with neutral hip and knee joint angles. The right leg was measured for every participant. To determine proper HD-EMG array positioning, BFLh and ST muscle borders were determined with B-mode 2D ultrasonography (Aloka α 10, Tokyo, Japan) and the skin over the borders was marked with a pen. After skin preparation, 15-channel semidisposable EMG arrays (OT Bioelettronica, Torino, Italy) were attached along the midline between the borders for each muscle using adhesive foam and connected to the amplifier of the EMG system.

The cavities of the electrode arrays were filled with 20 μ L conducting gel for proper electrode-skin contact (Figure 1). Electrode arrays were further secured with adhesive tape to minimize movement artifact. For BFLh, channel 8/9 from the distal end of the array was aligned with the midpoint on the line between the ischial tuberosity and knee joint fold. For ST, the array was attached below the tendinous inscription²⁸ of the muscle defined with ultrasonography. A reference electrode was placed over the left wrist. Signal quality was checked during submaximal knee flexion contractions. EMG data were sampled at 2048 Hz, amplified ($\times 1000$), and digitized (EMG-USB 12-bit A/D converter; OT Bioelettronica). During each task, 15 differential signals were recorded from each muscle using BioLab software (v3.1; OT Bioelettronica).

2.4.2 | Kinematics

Before performing the exercises, reflective markers (14 mm diameter) were secured over the anterior and posterior superior iliac spine, lateral thigh, lateral epicondyle of the femur, lateral shank, lateral malleolus, calcaneus, and second metatarsal head of each side, to determine hip and knee joint angular displacements. 3D marker displacements were recorded using an 8-camera motion analysis system sampling at 250 Hz in Nexus software (Vicon Motion Systems Inc., Oxford, UK).

2.4.3 | Force

Maximal voluntary isometric contraction and NHE force data (strain gauge at the ankle, see Video S1) were collected at 1000 Hz and digitized using an A/D converter (Cambridge Electronic Design, Cambridge, UK), and recorded in Spike2 software (Cambridge Electronic Design). Force and EMG data were synchronized by sending a pulse from Spike2 to the BioLab EMG software. Spike2 software was also used to send a digital pulse to the Nexus software to synchronize EMG and kinematic data.

2.5 | Data analysis

As NHE is an eccentric-only exercise, muscle activity levels were determined in the eccentric phase of each task to make exercises comparable. For NHE, force measured from the right leg was used to determine the active lengthening phase: from the start of force increment to the instant of peak force. For SDL, the eccentric phase was defined based on muscle-tendon length change: hip and knee joint angular displacements were calculated in Nexus software based on the Plug-in Gait Model after smoothing marker trajectories with an 8-Hz low-pass Butterworth filter. Joint angular data were then imported to Matlab (MathWorks Inc, Natick, MA, USA), where ST and BFlh muscle-tendon lengths were calculated using modeling equations.²⁹

Electromyography data were band-pass filtered using a 10-500 Hz fourth-order zero-phase Butterworth filter in Matlab. For MVICs, root-mean-square (RMS) EMG was calculated from a 1-second stable plateau for each EMG channel. In NHE and SDL, RMS activity in the eccentric phase (defined as above) was calculated. RMS values across repetitions were averaged for each exercise and expressed as a percentage of the highest RMS activity of the corresponding channel during any of the MVIC tasks (%MVIC). Channels 1-5, 6-10, and 11-15 were then averaged to represent activity in the distal, middle, and proximal regions, respectively. This approach minimized the effects of muscle shift under the skin on regional EMG activity. Overall activity was defined as the average normalized RMS activity of all 15 channels for each muscle.

2.6 | Statistical analysis

The magnitude of the differences (Cohen's $d \pm 90\%$ confidence limits) between the overall activity levels of ST and BFlh muscles, and regional activity within each muscle in NHE and SDL were calculated using a custom spreadsheet.³⁰ Differences were classified as trivial (<0.2), small (≥ 0.2), moderate (≥ 0.5), or large (≥ 0.8). Nonetheless, differences with 90% confidence intervals overlapping both the positive (≥ 0.2) and negative (≤ -0.2) smallest worthwhile standardized effects were deemed to be unclear effects.³¹

3 | RESULTS

Peak NHE force was 285 ± 48 N; load for SDL was 86.9 ± 25.8 kg (mean $\pm$ SD). Figures 2 and 3 represent group average normalized activity for each EMG channel during NHE and SDL, respectively. Regional EMG activities for each individual are illustrated in Figure 4.

Figure 2 Normalized electromyographic (EMG) activity along the biceps femoris long head (upper panel) and semitendinosus (bottom panel) during Nordic hamstring exercise. The figure represents group average ($N = 12$) for each channel of the EMG array along the corresponding muscle. EMG channels were time-normalized and low-pass filtered for visualization. MVIC, maximal voluntary isometric contraction.

During NHE, activity within ST was highest in the middle region ($80.48 \pm 13.78\%$ MVIC), and differences between regions were substantial (d range = 0.55- 1.41). BFLh activity was highest in the distal region ($72.08 \pm 10.66\%$ MVIC) and lowest in the proximal region ($57.74 \pm 15.95\%$ MVIC), with small to large differences between regions (d range = 0.38-1.25). In SDL, ST activity was highest in the middle region ($40.70 \pm 9.44\%$ MVIC), and differences between regions were small to moderate (d range = 0.29-0.67). In BFLh, the proximal region displayed the lowest activity level ($32.23 \pm 8.55\%$ MVIC) and the difference between the middle and distal regions was trivial ($d = 0.16 \pm 0.27$). All within-muscle comparisons are shown in Table 1.

With respect to overall activity, ST presented substantially higher activity than BFlh in NHE ($72.31 \pm 7.33\% \text{MVIC}$ vs $63.97 \pm 10.46\% \text{MVIC}$, $d = 1.06 \pm 0.45$), but in SDL, the difference between muscles was negligible ($37.46 \pm 6.74\% \text{MVIC}$ in ST vs $36.07 \pm 8.54\% \text{MVIC}$ in BFlh, $d = 0.19 \pm 0.34$).

Figure 3 Normalized electromyographic (EMG) activity along the biceps femoris long head (upper panel) and semitendinosus (bottom panel) muscles during stiff-leg deadlift. The figure represents group average ($N = 12$) for each channel of the EMG array along the corresponding muscle. EMG channels were time-normalized and low-pass filtered for visualization. MVIC, maximal voluntary isometric contraction.

4 | DISCUSSION

This study has shown using HD-EMG that intramuscular distribution of normalized EMG activity is non-uniform along the BFlh and ST muscles; the middle-to-proximal region of ST and the distal region of BFlh showed the highest within-muscle activity, irrespective of exercise. Regarding intermuscular differences, higher activity was observed in ST compared to BFlh in NHE, but no differences between muscles were observed in SDL.

Figure 4 Regional electromyographic (EMG) activity as a percentage of the activity during maximal voluntary isometric contraction (%MVIC) in the biceps femoris long head (BFIh, panel A) and semitendinosus (ST, panel B) for each participant during Nordic hamstring exercise (NHE) and stiff-leg deadlift (SDL). Filled black markers connected with dashed lines represent group average.

Table 1 Differences (Cohen's $d \pm 90\%$ confidence limits) between the activity level of muscle regions for each muscle and exercise.

	Semitendinosus		Biceps femoris long head	
Region	Middle	Proximal	Middle	Proximal
Nordic hamstring exercise				
Distal	1.41 ± 0.79^L	0.87 ± 0.60^L	-0.87 ± 0.38^L	-1.25 ± 0.71^L
Middle	-	-0.55 ± 0.52^M	-	-0.38 ± 0.53^S
Stiff-leg deadlift				
Distal	0.67 ± 0.63^M	0.29 ± 0.44^S	-0.16 ± 0.27^T	-0.63 ± 0.35^M
Middle	-	-0.38 ± 0.39^S	-	-0.48 ± 0.29^S

T = trivial difference, S = small difference, M = moderate difference, L = large difference between regions. Positive and negative values refer to higher activity in the relatively more proximal and distal regions, respectively.

4.1 | Region-specific muscle activity

In BFlh, the region effect was larger in NHE than in SDL, with large and moderate effects, respectively. In both tasks, the distal region was activated the most. This is consistent with preferentially distal BFlh muscle use based on mfMRI studies in NHE³² and in the mechanically similar eccentric knee extension task.¹³ However, rather homogeneous muscle metabolic activity was previously found in a modified SDL,¹⁴ contrary to our study wherein differences between regions were present, although smaller compared to NHE.

In ST, large differences in activity between regions were detected during NHE, whereby the middle and proximal regions were more active than the distal region. Similarly, in a previous study, middle and proximal regions of ST were more active than the distal region in eccentric knee extension.¹³ Another previous study found that the middle region showed the highest activity in a modified SDL,¹⁴ in accordance with the current study.

Abundant innervation of BFlh and ST likely contributes to the region-specific activity of these muscles. One-to-three primary nerves innervating different BFlh regions divide into two or more primary branches^{28,33} that may contribute to region-dependent activity. ST muscle is divided into upper and lower parts by a tendinous inscription, which is an attachment zone for most of the fascicles proximally and distally.²⁸ Both parts are innervated by a separate primary motor nerve implying separate function of regions. In this study, activity was recorded from below the tendinous inscription due its relatively proximal location. ST showed heterogeneous distribution of muscle activity within this part. Small territories of motor units may be responsible for independent activity of hamstring muscle regions, which is yet to be examined.

In other muscles, for example, in biceps brachii, region-specific activity has been linked to non-uniform hypertrophy after a training intervention.³⁴ Accordingly, ST is the thickest in the upper mid-region, which is the region where the highest activity was found in our study. On the contrary, BFlh showed the highest activity in the distal region where muscle thickness is lower compared to the middle and proximal regions.³⁵ However, BFlh architecture differs between regions, with shorter fascicles and larger pennation angle in the distal compared to the proximal region.³⁵ Thus, the distal region seems to be more suited to force production than the proximal region, which is more suited to larger excursions. On the contrary, Kellis et al³⁶ found higher pennation angle in the proximal compared to the distal region in cadavers, which might be due to disagreement in BFlh pennation angles between cadaver and in vivo measurements.³⁵ Intervention studies are needed to examine associations between region-specific activity and region-specific architectural and morphological changes along the hamstrings.

4.2 | Overall muscle activity

As hamstring injuries are most common in BFlh, muscle-specific exercise selection based on the activity of individual hamstring muscles is of significant interest. Architecturally, longer fascicle length and lower pennation angle of ST compared to BFlh^{28,37} suggests different functions of these muscles.

During NHE in the current study, ST showed higher relative activity compared to BFlh, which apparently associates with increased hypertrophy in ST relative to BFlh in response to NHE training.³⁸ On the contrary, in eccentric knee extension³⁹ and NHE,¹⁸ ST and BFlh were equally activated based on bipolar EMG. However, mfMRI studies during eccentric knee extension¹³ and NHE¹⁶ found higher activity in ST compared to BFlh, similar to the current study. The latter mfMRI study analyzed 3 slices at 40%, 50%, and 60% muscle length. During NHE in the current study, between-muscle differences were not as high as those found with mfMRI, but it is likely that this mfMRI study ignored the region of highest activity within BFlh, which is distal to 40% muscle length according to the current study.

In SDL, overall activity did not differ between muscles in this study. In a previous study, the mechanically similar Romanian deadlift was classified as an ST-dominant exercise based on bipolar EMG. On the contrary, another study⁴⁰ also using conventional bipolar EMG showed higher activity in BFlh than in ST during SDL. However, the same study compared these results to mfMRI data and found that differences in T2 changes between muscles were not observed when 5 slices along the muscles were analyzed, which is in accordance with the current study using HD-EMG. Although methods were not directly compared, on the basis of the above results, it seems that HD-EMG can provide a more comprehensive estimate of overall muscle activity than conventional bipolar EMG configurations.

As noted, previous studies are not in agreement concerning the relative activity of ST and BFlh in hamstring exercises. To estimate whole-muscle activity with mfMRI, 3-5 slices have been analyzed along hamstring muscles.^{15,38,40} However, this approach is not directly comparable to EMG studies wherein data are usually collected from a small muscle region. Large differences in spatial resolution between these methods may be one reason for discrepancies between studies, although mfMRI and EMG also assess different physiological mechanisms; metabolic and neural activity, respectively. As a limitation, EMG is prone to cross-talk. In this study, EMG activity measured from the ST and BFlh could have been contaminated by the activity of the semimembranosus and the short head of the biceps femoris, respectively. This possibility was presumably minimized by several factors: careful electrode array location using ultrasonography; applying 10 mm inter-electrode distance⁴¹ and using electrodes with a relatively shallow pick-up area; examining male athletes with relatively thin

subcutaneous tissue over the hamstrings. We also tried to improve EMG normalization by applying different MVIC tasks. However, studies targeting optimization of EMG normalization are needed to further improve comparability of muscles or muscle regions. Furthermore, deeper components of the hamstrings cannot be examined with surface EMG. HD-EMG may be a good complement to mfMRI to study ST and BFlh muscles, with real-time recording, financial considerations, and time efficiency among the advantages of HD-EMG over mfMRI.

4.3 | NHE and SDL for injury prevention

The issue of which exercises can help to prevent hamstring injuries is under debate. Regarding the mechanism of injury, most running-type injuries seem to happen in the late swing phase of high-speed running,⁴² whereas the hamstring muscle-tendon unit is actively lengthened and subjected to the highest forces and strain within the step cycle.^{43,44} In this phase, hamstring muscles seem to be highly activated.⁴⁵ To mimic the injury mechanism, it is generally believed that eccentrically activating hamstring muscles at a longer muscle length would be optimal from an injury prevention perspective. Nonetheless, an increasing body of evidence suggests that the Nordic hamstring exercise, despite requiring a substantially shorter muscle operating length than in the late swing phase, has a preventive effect against hamstring injuries.^{7,9,46} In NHE, ST was more active than BFlh, but both ST and BFlh were highly activated, supporting the idea of positive architectural changes (ie, fascicle elongation) within BFlh in response to NHE intervention.³⁸ In SDL, hamstrings work at a longer muscle length than in NHE, and in this study, relative activity of BFlh compared to ST was higher than in NHE (with negligible difference between muscles). However, absolute activity was substantially lower in both muscles (ST = 72% vs 37%, BFlh = 64% vs 36%, of MVIC on average in NHE vs SDL). It should be noted that the load was not matched for the two exercises. Instead loads that are generally used in training were applied, making these exercises comparable from a practical point of view. Due to the relatively low activity level in the eccentric phase of SDL, we speculate that SDL alone may not be as effective as NHE to prevent hamstring injuries.

Within BFlh, we observed the lowest activity in the proximal region in NHE. This may be associated with higher strain close to the proximal muscle-tendon junction in BFlh during cyclic knee flexion-extension contractions.⁴⁷ Silder et al⁴⁷ also observed higher strain in previously injured BFlh, which may be associated with lower EMG activity in the BFlh of the injured limb.⁴⁸ The association between strain magnitude and EMG activity level should be further studied to reveal whether HD-EMG may be a useful tool for hamstring injury risk management.

Compared to the knee-dominant NHE, based on mfMRI studies,^{14,24} we expected relatively higher activity in the proximal BFlh compared to the distal region in the hip-dominant SDL. Instead, similar

proximal-distal activity patterns were observed in NHE and SDL. Even although the most proximal region cannot be measured with surface EMG, this study suggests that hip-dominant exercises do not necessarily activate proximal BFlh preferentially. Future studies should further examine whether the relative activity of muscle regions can be modulated with different exercises.

5 | PERSPECTIVE

Future studies should use HD-EMG to examine whether regional differences can be observed in other exercises and high-injury risk tasks. It should be emphasized that although proximal-distal differences seem to be significant, the clinical relevance of this phenomenon is yet to be clarified. In previously strained BFlh, decreased EMG activity,^{48,49} slower EMG onset,⁵⁰ and decreased metabolic activity¹⁵ were observed. It may be of value to examine whether inhibition is region-specific and whether exercise selection could affect the relative activity of muscle regions. This knowledge will help to identify the most suitable exercises for interventions and improve the restoration of neuromuscular function following a hamstring injury.

REFERENCES

1. Opar DA, Drezner J, Shield A, et al. Acute hamstring strain injury in track-and-field athletes: a 3-year observational study at the Penn Relay Carnival. *Scand J Med Sci Sports*. 2014;24:e254-e259. PMID: 24330073.
2. Orchard JW, Seward H, Orchard JJ. Results of 2 decades of injury surveillance and public release of data in the Australian Football League. *Am J Sports Med*. 2013;41:734-741.
3. Roe M, Murphy JC, Gissane C, Blake C. Hamstring injuries in elite Gaelic football: an 8-year investigation to identify injury rates, time-loss patterns and players at increased risk. *Br J Sports Med*. Published Online First: 20 October 2016. doi: 10.1136/bjsports-2016-096401. PMID: 27797729.
4. Opar DA, Williams MD, Timmins RG, Hickey J, Duhig SJ, Shield AJ. Eccentric hamstring strength and hamstring injury risk in Australian footballers. *Med Sci Sports Exerc*. 2015;47:857-865.
5. Thorborg K. What are the most important risk factors for hamstring muscle injury? *Clin J Sport Med*. 2014;24:160-161.
6. Timmins RG, Bourne MN, Shield AJ, Williams MD, Lorenzen C, Opar DA. Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *Br J Sports Med*. 2016;50:1524-1535.
7. Arnason A, Andersen TE, Holme I, Engebretsen L, Bahr R. Prevention of hamstring strains in elite soccer: an intervention study. *Scand J Med Sci Sports*. 2008;18:40-48.
8. Askling CM, Tengvar M, Tarassova O, Thorstensson A. Acute hamstring injuries in Swedish elite sprinters and jumpers: a prospective randomised controlled clinical trial comparing two rehabilitation protocols. *Br J Sports Med*. 2014;48:532-539.
9. Petersen J, Thorborg K, Nielsen MB, Budtz-Jorgensen E, Holmich P. Preventive effect of eccentric training on acute hamstring injuries in men's soccer: a cluster-randomized controlled trial. *Am J Sports Med*. 2011;39:2296-2303.
10. van der Horst N, Smits DW, Petersen J, Goedhart EA, Backx FJ. The preventive effect of the nordic hamstring exercise on hamstring injuries in amateur soccer players: a randomized controlled trial. *Am J Sports Med*. 2015;43:1316-1323.

11. Ekstrand J, Walden M, Hagglund M. Hamstring injuries have increased by 4% annually in men's professional football, since 2001: a 13-year longitudinal analysis of the UEFA Elite Club injury study. *Br J Sports Med.* 2016;50:731-737.
12. Flaxman TE, Alkjaer T, Simonsen EB, Krogsgaard MR, Benoit DL. Predicting the functional roles of knee joint muscles from internal joint moments. *Med Sci Sports Exerc.* 2017;49:527-537.
13. Kubota J, Ono T, Araki M, Torii S, Okuwaki T, Fukubayashi T. Non-uniform changes in magnetic resonance measurements of the semitendinosus muscle following intensive eccentric exercise. *Eur J Appl Physiol.* 2007;101:713-720.
14. Mendez-Villanueva A, Suarez-Arrones L, Rodas G, et al. MRI-based regional muscle use during hamstring strengthening exercises in elite soccer players. *PLoS One.* 2016;11:e0161356.
15. Bourne MN, Opar DA, Williams MD, Al Najjar A, Shield AJ. Muscle activation patterns in the Nordic hamstring exercise: Impact of prior strain injury. *Scand J Med Sci Sports.* 2016;26:666-674.
16. Bourne MN, Williams MD, Opar DA, Al Najjar A, Kerr GK, Shield AJ. Impact of exercise selection on hamstring muscle activation. *Br J Sports Med.* 2017;51(13):1021-1028. PMID: 27467123.
17. Jonhagen S, Halvorsen K, Benoit DL. Muscle activation and length changes during two lunge exercises: implications for rehabilitation. *Scand J Med Sci Sports.* 2009;19:561-568.
18. Zebis MK, Skotte J, Andersen CH, et al. Kettlebell swing targets semitendinosus and supine leg curl targets biceps femoris: an EMG study with rehabilitation implications. *Br J Sports Med.* 2013;47:1192-1198.
19. Drost G, Stegeman DF, van Engelen BG, Zwarts MJ. Clinical applications of high-density surface EMG: a systematic review. *J Electromyogr Kinesiol.* 2006;16:586-602.
20. Cronin NJ, Kumpulainen S, Joutjärvi T, Finni T, Piitulainen H. Spatial variability of muscle activity during human walking: the effects of different EMG normalization approaches. *Neuroscience.* 2015;300:19-28.
21. Gallina A, Ivanova TD, Garland SJ. Regional activation within the vastus medialis in stimulated and voluntary contractions. *J Appl Physiol.* 2016;121:466-474.
22. Wakeling JM. The recruitment of different compartments within a muscle depends on the mechanics of the movement. *Biol Lett.* 2009;5:30-34.
23. Watanabe K, Kouzaki M, Moritani T. Regional neuromuscular regulation within human rectus femoris muscle during gait. *J Biomech.* 2014;47:3502-3508.

24. Mendiguchia J, Garrues MA, Cronin JB, et al. Nonuniform changes in MRI measurements of the thigh muscles after two hamstring strengthening exercises. *J Strength Cond Res.* 2013;27:574-581.
25. Baechle TR, Earle RW. *Essentials of Strength Training and Conditioning*. Champaign, IL: Human Kinetics; 2008:395-396.
26. Komi PV, Linnamo V, Silventoinen P, Sillanpaa M. Force and EMG power spectrum during eccentric and concentric actions. *Med Sci Sports Exerc.* 2000;32:1757-1762.
27. Opar DA, Piatkowski T, Williams MD, Shield AJ. A novel device using the Nordic hamstring exercise to assess eccentric knee flexor strength: a reliability and retrospective injury study. *J Orthop Sports Phys.* 2013;43:636-640.
28. Woodley SJ, Mercer SR. Hamstring muscles: architecture and innervation. *Cells Tissues Organs.* 2005;179:125-141.
29. Hawkins D, Hull ML. A method for determining lower extremity muscle-tendon lengths during flexion/extension movements. *J Biomech.* 1990;23:487-494.
30. Hopkins WG. Spreadsheets for analysis of controlled trials with adjustment for a predictor. *Sportscience.* 2006;10:46-50.
31. Batterham AM, Hopkins WG. Making meaningful inferences about magnitudes. *Int J Sports Physiol Perform.* 2006;1:50-57.
32. Mendiguchia J, Arcos AL, Garrues MA, Myer GD, Yanci J, Idoate F. The use of MRI to evaluate posterior thigh muscle activity and damage during nordic hamstring exercise. *J Strength Cond Res.* 2013;27:3426-3435.
33. Seidel PM, Seidel GK, Gans BM, Dijkers M. Precise localization of the motor nerve branches to the hamstring muscles: an aid to the conduct of neurolytic procedures. *Arch Phys Med Rehabil.* 1996;77:1157-1160.
34. Wakahara T, Fukutani A, Kawakami Y, Yanai T. Nonuniform muscle hypertrophy: its relation to muscle activation in training session. *Med Sci Sports Exerc.* 2013;45:2158-2165.
35. Tosovic D, Muirhead JC, Brown JM, Woodley SJ. Anatomy of the long head of biceps femoris: an ultrasound study. *Clin Anat.* 2016;29:738-745.
36. Kellis E, Galanis N, Natsis K, Kapetanios G. Muscle architecture variations along the human semitendinosus and biceps femoris (long head) length. *J Electromyogr Kinesiol.* 2010;20:1237-1243.

37. Kellis E, Galanis N, Natsis K, Kapetanios G. Validity of architectural properties of the hamstring muscles: correlation of ultrasound findings with cadaveric dissection. *J Biomech.* 2009;42:2549-2554.
38. Bourne MN, Williams MD, Opar DA, Al Najjar A, Kerr GK, Shield AJ. Impact of the Nordic hamstring and hip extension exercises on hamstring architecture and morphology: implications for injury prevention. *Br J Sports Med.* 2017;51:469-477.
39. Higashihara A, Ono T, Kubota J, Fukubayashi T. Differences in the electromyographic activity of the hamstring muscles during maximal eccentric knee flexion. *Eur J Appl Physiol.* 2010;108:355-362.
40. Ono T, Higashihara A, Fukubayashi T. Hamstring functions during hip-extension exercise assessed with electromyography and magnetic resonance imaging. *Res Sports Med.* 2011;19:42-52.
41. De Luca CJ, Kuznetsov M, Gilmore LD, Roy SH. Interelectrode spacing of surface EMG sensors: reduction of crosstalk contamination during voluntary contractions. *J Biomech.* 2012;45:555-561.
42. Heiderscheit BC, Hoerth DM, Chumanov ES, Swanson SC, Thelen BJ, Thelen DG. Identifying the time of occurrence of a hamstring strain injury during treadmill running: a case study. *Clin Biomech.* 2005;20:1072-1078.
43. Schache AG, Dorn TW, Blanch PD, Brown NA, Pandy MG. Mechanics of the human hamstring muscles during sprinting. *Med Sci Sports Exerc.* 2012;44:647-658.
44. Thelen DG, Chumanov ES, Sherry MA, Heiderscheit BC. Neuromusculoskeletal models provide insights into the mechanisms and rehabilitation of hamstring strains. *Exerc Sport Sci Rev.* 2006;34:135-141.
45. Higashihara A, Nagano Y, Ono T, Fukubayashi T. Differences in activation properties of the hamstring muscles during overground sprinting. *Gait Posture.* 2015;42:360-364.
46. Mjolsnes R, Arnason A, Osthaugen T, Raastad T, Bahr R. A, 10- week randomized trial comparing eccentric vs. concentric hamstring strength training in well-trained soccer players. *Scand J Med Sci Sports.* 2004;14:311-317.
47. Silder A, Reeder SB, Thelen DG. The influence of prior hamstring injury on lengthening muscle tissue mechanics. *J Biomech.* 2010;43:2254-2260.
48. Opar DA, Williams MD, Timmins RG, Dear NM, Shield AJ. Knee flexor strength and bicep femoris electromyographical activity is lower in previously strained hamstrings. *J Electromyogr Kinesiol.* 2013;23:696-703.

49. Opar DA, Williams MD, Timmins RG, Dear NM, Shield AJ. Rate of torque and electromyographic development during anticipated eccentric contraction is lower in previously strained hamstrings. *Am J Sports Med.* 2013;41:116-125.

50. Sole G, Milosavljevic S, Nicholson H, Sullivan SJ. Altered muscle activation following hamstring injuries. *Br J Sports Med.* 2012;46:118-123.