



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document, This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Horswill, C., Manica, A., Daunt, F., Newell, M., Wanless, S., Wood, M. and Matthiopoulos, J. (2021), Improving assessments of data-limited populations using life-history theory. *J Appl Ecol.* Accepted Author Manuscript. <https://doi.org/10.1111/1365-2664.13863> and is licensed under Creative Commons: Attribution 4.0 license:

Horswill, C, Manica, A, Daunt, F, Newell, Marie-Louise, Wanless, S, Wood, Matt J ORCID logo
ORCID: <https://orcid.org/0000-0003-0920-8396> and Matthiopoulos, J (2021) Improving assessments of data-limited populations using life-history theory. *Journal of Applied Ecology*, 58 (6). pp. 1225-1236. doi:10.1111/1365-2664.13863

Official URL: <http://doi.org/10.1111/1365-2664.13863>

DOI: <http://dx.doi.org/10.1111/1365-2664.13863>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/9481>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Journal of Applied Ecology

DR CAT HORSWILL (Orcid ID : 0000-0002-1795-0753)

Article type : Research Article

Handling Editor - Bret Elder

Improving assessments of data-limited populations using life-history theory

C. Horswill^{1,2,3,4*}, A. Manica², F. Daunt⁵, M. Newell⁵, S. Wanless⁵, M. Wood⁶, and J. Matthiopoulos¹

¹Institute of Biodiversity, Animal Health & Comparative Medicine, University of Glasgow, Glasgow, G12 8QQ

²Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3DT

³ZSL Institute of Zoology, Regent's Park, London, NW1 4RY, UK

⁴Centre for Biodiversity and Environmental Research, Department of Genetics, Evolution and Environment, University College London, Gower Street, London, WC1E 6BT, UK

⁵UK Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB

⁶School of Natural and Social Sciences, University of Gloucestershire, Francis Close Hall, Cheltenham, GL50 4AZ

*Corresponding author: catrsw@gmail.com

Keywords: black-legged kittiwake, data-limited, environmental impact assessment, fecundity, marine renewables, missing data, population assessment, seabird, survival

Running title: Assessments of data-limited populations

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/1365-2664.13863](https://doi.org/10.1111/1365-2664.13863)

This article is protected by copyright. All rights reserved

Abstract

1. Predicting how populations may respond to climate change and anthropogenic pressures requires detailed knowledge of demographic traits, such as survival and reproduction. However, the availability of these data varies greatly across space and taxa. Therefore, it is common practice to conduct population assessments by filling in missing values from surrogate species or other populations of the same species. However, using these independent surrogate values concurrently with observed data neglects the life-history trade-offs that connect the different aspects of a population's demography, thus introducing biases that could ultimately lead to erroneous management decisions.
2. We use a Bayesian hierarchical approach to combine fragmented multi-population data with established life-history theory and reconstruct population-specific demographic data across a substantial part of a species breeding range. We apply our analysis to a long-lived colonial species, the black-legged kittiwake *Rissa tridactyla*, that is classified as globally Vulnerable and is highly threatened by increasing anthropogenic pressures, such as offshore renewable energy development. We then use a projection analysis to examine how the reconstructed demographic parameters may improve population assessments, compared to models that combine observed data with independent surrogate values.
3. Reconstructed demographic parameters can be utilised in a range of population modelling approaches. They can also be used as reference estimates to assess whether independent surrogate values are likely to over or underestimate missing demographic parameters. We show that surrogate values from independent sources are often used to fill in missing parameters that have large potential demographic impact, and that resulting biases can be driven in unpredictable directions thus precluding assessments from being consistently precautionary.
4. *Synthesis and applications:* Our study dramatically increases the spatial coverage of population-specific demographic data for black-legged kittiwakes. The reconstructed demographic parameters presented can also be used immediately to reduce uncertainty in the consenting process for offshore wind development in the UK and Ireland. More broadly, we show that the reconstruction approach used here provides a new avenue for improving evidence-based management and policy action for animal and plant populations with fragmented and error prone demographic data.

Introduction

Predicting how vulnerable populations are likely to respond to future threats requires detailed knowledge of life-history traits, including survival and reproduction (Pearson et al., 2014; Salguero-Gómez et al., 2016; Winemiller, 2005). However, our ability to monitor individual traits can differ greatly between traits, such that the availability of data is highly unbalanced across space and taxa (Horswill et al., 2019). This problem is so widespread that even the most optimistic data collection programme is not sufficient as a single solution to fill in all missing data. Therefore, identifying reliable ways to assess population status, even when

demographic data are missing, has been highlighted as a key ecological problem facing policy makers (Kindsvater et al., 2018).

A typical approach for assessing populations with missing demographic data involves filling in gaps with values from surrogate species or populations of the same species (Githiru, Matthysen, Lens, & Bennun, 2007; Hernández-Camacho, Bakker, Aurióles-Gamboa, Laake, & Gerber, 2015; Schtickzelle, WallisDeVries, & Baguette, 2005). However, combining independent surrogate estimates with observed demographic data neglects the physiological compromises that connect the different aspects of a population's demography (Stearns, 1992). Covariation among life-history traits is widely recognised across animal and plant groups (Bakewell, Davis, Freckleton, Isaac, & Mayhew, 2020; Healy, Ezard, Jones, Salguero-gómez, & Buckley, 2019; Jeschke & Kokko, 2009; Salguero-Gómez et al., 2016), and reflects the core relationships that connect fitness-related traits, such as body growth, reproduction and survival (Bennett & Owens, 2002; Lande, 1982; Stearns, 1992). At the extremes, species grow slowly, mature late, have low fecundity and high survival, or the reverse pattern. Similar patterns of covariation among life-history traits also exist across populations of the same species (Devenish-Nelson, Harris, Soulsbury, Richards, & Stephens, 2013; Frederiksen, Harris, & Wanless, 2005; Horswill et al., 2019). Therefore, population assessments that fill in missing values with demographic parameters from independent surrogate sources will incorporate biases and imprecisions that may ultimately lead to population mismanagement (Devenish-Nelson et al., 2013; Hernández-Camacho et al., 2015; Johnson, Mills, Stephenson, & Wehausen, 2010).

Predictive approaches that combine available data with established life-history trade-offs between demographic traits offer a new avenue for addressing taxonomic, spatial and temporal imbalances in demographic data (Horswill et al., 2019; Kindsvater et al., 2018; Thorson, Munch, Cope, & Gao, 2017). This method considers multiple life-history traits, populations, and species simultaneously in a hierarchical framework to reconstruct missing values based on the trade-offs that determine an organism's life-history strategy. The approach provides estimates for traits that are poorly monitored or difficult to measure empirically, and supports the estimation of parameter uncertainty based on measurement errors and data availability (Horswill et al., 2019). Consequently, we can use this technique to account for life-history trade-offs when filling in missing demographic data, as well as estimate parameter uncertainty that can be propagated through to population predictions to provide precautionary estimates of population change.

In this study, we use a predictive method based on life-history trade-offs (Horswill et al., 2019) to reconstruct population-specific demographic information across a substantial part of a species breeding range. The reconstruction framework used here can be applied across the plant and animal kingdom and we apply our analysis to an intensively monitored species of seabird, black-legged kittiwake *Rissa tridactyla*, where the spatial coverage of demographic monitoring is highly unbalanced between traits (Fig. 1). Black-legged kittiwakes are the most numerous gull in the world (Coulson, 2011); however, long-term

population declines mean that this species is currently categorised as globally Vulnerable (Birdlife International, 2019). A lack of population-specific demographic information for black-legged kittiwakes has also recently been highlighted as a key contributing factor to uncertainty in the consenting process for offshore wind development in the UK (Ruffino, Thompson, & O'Brien, 2020). We conduct a projection analysis based on age-structured models to examine how reconstructed (i.e., correlated) demographic parameters may improve population assessments in data-limited situations, compared to assessments that fill in missing data with independent surrogate values.

Materials and Methods

Data

We considered two demographic parameters for populations of black-legged kittiwakes breeding in the UK and Ireland: adult survival and fecundity (i.e., reproduction). The number of black-legged kittiwake populations in the UK and Ireland with empirical estimates for rates of fecundity ($n=68$) far exceeds the number of populations with empirical estimates for rates of adult survival ($n=5$, Fig. 1, supp. info. Table S1). For the populations that have empirical estimates for both adult survival and fecundity ($n=5$), we restricted the datasets to years when both demographic rates were monitored (supp. info. Table S1). For the survival dataset, we limited our analysis to studies using mark-recapture modelling that had at least eight years of data. The ability to estimate adult survival rates dramatically increases when mark-recapture time series are between five and ten years in duration (Horswill, Humphreys, & Robinson, 2018). For the wider fecundity dataset, we limited our analysis to populations and study plots in the Seabird Monitoring Programme Database (SMP 2020) that had more than 50 breeding pairs and at least five years of data since 2009. Fecundity can be estimated with considerably shorter time series than are required to estimate rates of survival, and we selected this threshold for population size in order to minimise the influence of ecological processes that differentially affect small populations, such as demographic stochasticity (Lande, Engen, & Sæther, 2003) and compensatory regulation (Horswill, O'Brien, & Robinson, 2017). The field protocols for collecting data at each site followed international guidelines for seabird monitoring (Walsh et al., 1995).

Potential demographic impact of vital rates

We used elasticity analysis to examine the potential demographic impact of adult survival and fecundity to population growth. We applied this analysis to the five populations with observed data for both demographic rates (Fig. 1). The projection matrices (A) followed a pre-breeding census (Caswell, 2001), and we used the mean observed rates of adult survival (Φ_i) and fecundity (F_i) for each population (Eqn. 1). The matrices were structured by age and the dimensions matched the mean age of first reproduction for black-legged kittiwake, i.e., four years (Horswill & Robinson, 2015). We calculated the elasticity of each demographic rate using the statistical package "popbio" (v. 2.7) (Stubben, Milligan, & Nantel, 2016) in Program R (v. 4.0.2) (R Core Team, 2020). This analysis estimates the proportional change in population growth given a proportional perturbation in each vital rate (Caswell, 2001).

$$A_i = \begin{pmatrix} 0 & 0 & 0 & J_i \frac{F_i}{2} \\ I_i & 0 & 0 & 0 \\ 0 & \Phi_i & 0 & 0 \\ 0 & 0 & \Phi_i & \Phi_i \end{pmatrix} \quad (1)$$

Like many seabirds, black-legged kittiwakes are largely unobservable during the first years of life. Therefore, population-specific estimates of juvenile (i.e., from fledging to age 1 year) and immature (i.e., before the age of first breeding) survival rates are limited (Horswill & Robinson, 2015). Available estimates for black-legged kittiwakes breeding in France report rates of juvenile survival (J_i) to be 75% of adult survival, immature rates of survival between age 1 and 2 years to be 87.5% of adult survival (I_i), and birds to have survival rates comparable to adults from age 2 onwards (Cam, Cooch, & Monnat, 2005). Therefore, we assumed an additive relationship between age-specific survival rates (Cam et al., 2005; Desprez, Pradel, Cam, Monnat, & Gimenez, 2011; Horswill et al., 2014, 2016), and estimated juvenile and immature survival rates to be 75% and 87.5% of the reported population-specific rates of adult survival, respectively. We assumed a 1:1 sex ratio in the population and halved fecundity to model female numbers only.

Reconstructing demographic parameters for populations of black-legged kittiwakes

We constructed a Bayesian hierarchical model (hereafter “correlated model”; Kindsvater et al. 2018; Horswill et al. 2019) to reconstruct mean rates of adult survival and fecundity for 68 populations of black-legged kittiwake breeding in the UK and Ireland. This represents approximately 37% of the UK population and 3% of the global breeding population. The correlated model uses stochastic functions to quantify the life-history trade-off connecting rates of adult survival to fecundity and account for sampling variability in these traits. Incorporating these sources of variance and covariance into the statistical imputation process means that the reconstructed (i.e., correlated) demographic parameters provide more biologically plausible replacements of missing values, compared to independent surrogate data from similar species or other populations of the same species (e.g. Horswill et al., 2019; Kindsvater et al., 2018).

The demographic component of the correlated model (Equation 2) contained linear functions for adult survival (ϕ) and fecundity (f):

$$\begin{aligned} \text{logit}(\phi_i) &= \gamma_{\phi,i} \\ \log(f_i) &= \gamma_{f,i} \end{aligned} \quad (2)$$

Here, we use lower-case notation to denote reconstructed demographic parameters. Subscript i denotes population, and γ is a multivariate normal (MVN) random effect that hierarchically connects rates of

adult survival to rates of fecundity. The covariance structure (Σ) of the random effect allows the relationship (i.e., life-history trade-off) between demographic parameters to emerge during model fitting:

$$(\gamma_{*i}) \sim MVN(\mu, \Sigma), \mu = (\mu_\phi, \mu_f), \mu_\phi \sim N(1.41, 0.03), \mu_f \sim N(-0.43, 0.03) \quad (3)$$

The mean values of the multivariate normal distribution (μ) represent the national mean values for each vital rate, and were resolved on the scale of the linear predictor. Therefore, we assigned these parameters from normal prior distributions (N). We set the expectation of these prior distributions using estimates of black-legged kittiwake demographic rates in France (Cap Sizun); i.e., adult survival = $\text{logit}(0.804)$, and fecundity = $\log(0.651)$ (cited by Frederiksen et al., 2005), and set the variance to allow imputation of the mean value within a realistic range for black-legged kittiwakes in the UK and Ireland (i.e., 0.03). The observed range of population-specific survival estimates was 0.81 to 0.86, and the 95% confidence interval of the prior distribution for the mean value was 0.69 to 0.89 (supp. info. Fig. S1A). Likewise, the observed range of population-specific fecundity estimates was 0.24 to 1.25, and the 95% confidence interval of the prior distribution for the mean value was 0.35 to 1.16 (supp. info. Fig. S1B) (see Fig. S2 for a comparison between the prior and posterior distributions of these parameters on the scale of the linear predictor).

We parameterised the variance–covariance matrix of the multivariate normal distribution (Σ , Equation 4) using an inverse precision matrix. We set the range of the variance priors (σ_ϕ, σ_f) to reflect predictions for long-lived species (such as black-legged kittiwakes) based on life-history theory, i.e., low variance in rates of adult survival and higher variance in rates of fecundity. The limited number of data points for adult survival required the shape of the prior distribution for variance in this trait to be more constrained than the equivalent prior distribution for fecundity, i.e., we used a truncated (T) normal distribution (N) for survival and uniform distribution (U) for fecundity (Eqn. 4). The correlation parameter was drawn from a normal distribution, where the expectation was informed by the life-history trade-off between adult survival and fecundity reported for black-legged kittiwakes by Frederiksen et al. (2005); i.e., -0.62. We kept the variance of this prior distribution relatively broad (i.e., 0.05) because we wanted the data to determine the strength of the trade-off (Eqn. 4, see supp. info. Fig. S3 for a comparison between the prior and posterior distributions of this parameter). We truncated the correlation parameter at -0.9 and 0.9 to prevent unrealistically strong correlations.

$$\Sigma = \text{Inv} \begin{pmatrix} \sigma_\phi^2 & \rho\sigma_\phi\sigma_f \\ \rho\sigma_\phi\sigma_f & \sigma_f^2 \end{pmatrix} \quad (4)$$

$$\sigma_\phi \sim N(0.25, 0.05)T(0.1, 1)$$

$$\sigma_f \sim U(0.1, 1)$$

$$\rho \sim N(-0.62, 0.05)T(-0.9, 0.9)$$

We linked the empirical data for survival and fecundity with the reconstructed (i.e., correlated) demographic parameters through an observation component that reflected sampling variability. Survival values were

drawn from a beta distribution and fecundity values from a truncated gamma distribution. We selected these distributions to bound inference within a plausible range on the observed scale, i.e., 0-1 for survival, and 0-3 for fecundity. We truncated the gamma distribution at three to match the maximum clutch size for black-legged kittiwakes.

$$\begin{aligned}\Phi_i &\sim B(\alpha_i, \beta_i) \\ F_i &\sim \text{Gamma}(r_i, \mu_i)T(0, 3)\end{aligned}\tag{5}$$

Here, upper-case notation denotes empirical data for survival (Φ_i) and fecundity (F_i) on the observed scale. We defined the shape parameters of these distributions ($\alpha_i, \beta_i, r_i, \mu_i$) so that the population values reconstructed in Equation 2 (ϕ_i, f_i) provided the expectation, and published measures of sampling variability for black-legged kittiwakes determined the variance; i.e., 6×10^{-4} for survival and 3×10^{-3} for fecundity (Frederiksen, Wanless, Harris, Rothery, & Wilson, 2004) (supp. info. Appendix 1). This stochastic node acts on data points with and without observed data to provide population-specific values with estimates of uncertainty for all survival and fecundity parameters.

To assess the influence of using informative prior distributions for certain parameters in our correlated model, we replaced these with uniform prior distributions that had biologically or theoretically driven bounds and re-ran our analysis (supp. info. Appendix 3) (Hobbs & Hooten, 2015). Using a bounded uniform prior distribution to assign the expectation for survival, fecundity, and the correlation parameter in the multivariate normal distribution, did not prevent model convergence but decreased the minimum effective sample size, and increased the posterior credible intervals, of the reconstructed demographic rates (supp. info. Appendix 3). By contrast, using a bounded uniform prior distribution to assign the variance parameter for survival in the variance-covariance matrix (i.e., replicating the prior distribution used to assign the variance parameter for fecundity, Eqn. 4) prevented model convergence (supp. info. Appendix 3).

We implemented all models using JAGS (v. 4.3.0) via the “jagsUI” library (v 1.5.1) (Kellner, 2016) for program R (v. 4.0.2). We fitted the models by running three Monte Carlo Markov chains (MCMC) for 2×10^5 iterations and retaining every 100th step to minimise autocorrelation in the MCMC sampling. We initialised each chain at different points in the parameter space. To confirm convergence of the chains we used the Brooks–Gelman–Rubin diagnostic tool (all values $\hat{r} < 1.01$) and the effective sample size of each parameter (all values > 950). Convergence occurred before 5000 MCMC draws, and therefore we removed the first 5000 draws as burn-in. We provide a guide on how to assess model convergence in the supplementary material (supp. info. Appendix 2).

Population assessments using independent and correlated demographic parameters

We conducted a projection analysis to examine the potential biases and imprecisions generated by assessing populations with independent (i.e., combining observed population-specific data with

independent surrogate values) and correlated demographic parameters (i.e., values reconstructed using the correlated model). In long-lived species, such as seabirds, the expectation is that rates of adult survival will experience low levels of variability, such that mean values should be relatively constant between populations. Therefore, it is common practice to use the national mean in place of missing adult survival data to conduct impact assessments on data limited seabird populations (e.g. O'Brien, Cook, & Robinson, 2017). The life-history trade-off (i.e., covariance) between rates of survival and fecundity means that the difference between the national mean rate of adult survival and the true population-specific value will be greatest for populations with the highest and lowest rates of fecundity. Consequently, we restricted the projection analysis to the ten populations with the highest and lowest rates of fecundity (Fig. 3B), thereby examining the worst-case scenario for introducing bias into population assessments.

We conducted the projection analysis under four scenarios (Table 1). We ran each projection scenario for 25 years and 1000 iterations. In the first and second projection scenarios, we held the mean demographic rates constant across all iterations. For the first scenario we projected populations using independent demographic parameters by combining population-specific rates of fecundity with the national mean rate of adult survival. To estimate the national mean rate of adult survival for black-legged kittiwakes we used the five study populations with available data (Fig. 1, national mean adult survival=0.84, range=0.81-0.86). In the second projection scenario we projected populations using correlated demographic parameters that incorporated life-history trade-offs between traits, i.e., we use the median posterior values reconstructed from our correlated model.

For the third and fourth projection scenarios we propagated parameter uncertainty through to the population predictions. In these models, each iteration received a different set of baseline demographic parameters. For the third scenario, we added sampling variability that was uncorrelated between the two demographic traits. To do this, we randomly selected survival values from a beta distribution and fecundity values from an independent truncated gamma distribution. The expectation of these distributions was set to the national mean for rates of adult survival, and the population-specific mean for rates of fecundity, and we set the variance to match the trait-specific variance used in the correlated model (Eqn. 5). Finally, for the fourth projection scenario, we randomly selected correlated values of adult survival and fecundity from the complete joint posterior distribution imputed by our correlated model (Eqns. 2-3). Thus, in this final scenario, we maintained covariance between traits whilst accounting for uncertainty associated with sampling variability and confidence in the reconstructed life-history trade-off between traits.

To quantitatively compare the four projection scenarios, we examined the distribution of absolute (i.e., final) population sizes generated and measured vulnerability as the percentage of projections that resulted in a population size of less than 50 breeding females. We used this threshold because compensatory regulation associated with increased predation at low density differentially affects individual populations of black-legged kittiwakes (Horswill, O'Brien, et al., 2017) and it is not possible to identify which populations

experience this regulatory pressure. We used pairwise comparisons between the outputs of our four projection scenarios to examine how independent and correlated demographic parameters and their associated uncertainty influence population assessments. For example, by comparing the outcomes of projection scenarios 1 and 2, we examined how independent demographic parameters influence population assessments, compared to correlated values (Table 1). By comparing the outcomes of projection scenarios 1 and 3, as well as scenarios 2 and 4, we examined how incorporating parameter uncertainty can influence population assessments, and by comparing the outcomes of projection scenarios 2 and 4, we examined the effect of incorporating uncertainty that also accounts for observed levels of covariance between demographic parameters (Table 1).

For each projection scenario (p) we used an age-structured model to predict the size (S) of each population (i) at time $t + 1$ as a function of population size at time t and constant demographic parameters (Eqn. 6). All projection scenarios also included demographic stochasticity using binomial and Poisson distributions for survival and fecundity events, respectively. We structured the projection matrices for these deterministic models following the same principles that we used to construct the projection matrices in the elasticity analysis (Eqn. 1).

$$S_{i,t+1,p}^r = \begin{pmatrix} 0 & 0 & 0 & J_{i,p} \frac{f_{i,p}}{2} \\ I_{i,p} & 0 & 0 & 0 \\ 0 & \phi_{i,p} & 0 & 0 \\ 0 & 0 & \phi_{i,p} & \phi_{i,p} \end{pmatrix} S_{i,t,p} \quad (6)$$

We initiated each population with 1200 females across a stable age distribution derived from the dominant eigenvector of the projection matrix (Eqn. 6) (Caswell, 2001). Following other assessments of black-legged kittiwake populations that use an age-structured matrix modelling approach, we applied a maximum population size to prevent exponential population growth ($S_{\max} = 1600$, Miller et al. 2019).

Results

Potential demographic impact of vital rates

As expected for long-lived species, we found that elasticity estimates for black-legged kittiwake demographic rates were consistently higher for rates of adult survival (mean=0.61, SD=0.04), compared to rates of survival from hatching to age 1-year (mean=0.10, SD=0.01). The much greater potential demographic impact of adult survival on population growth highlights the importance of using accurate estimates of this demographic parameter in population assessments.

Reconstructing demographic parameters for populations of black-legged kittiwakes

We detail the reconstructed (i.e., correlated) rates of survival and fecundity for 68 populations of black-legged kittiwakes in the supp. info. Table S1 (including 63 populations unmonitored for rates of adult survival). The correlation (i.e., life-history trade-off) between rates of adult survival and fecundity was -0.57 ($SD=0.18$) (Fig. 2, supp. info. Fig. S3). Incorporating sampling variability and life-history trade-offs into the analytical framework enabled the imputation of posterior distributions for both missing and observed data points, i.e., all survival and fecundity values. All observed data points fell within the 95% credible intervals of these posterior distributions (Fig. 3).

Population assessments using independent and correlated demographic parameters

Using independent demographic parameters resulted in faster rates of decline in the low fecundity group, and slower rates of decline in the high fecundity group, compared to assessments using correlated demographic parameters (Table 2; Fig. 4A-B). Therefore, the difference between the predicted absolute population sizes for populations with high and low rates of fecundity was larger when we used independent demographic parameters, as opposed to correlated values. The percentage of model simulations that resulted in vulnerable populations was considerably larger for the low fecundity group when we used independent demographic parameters, rather than correlated values. Including parameter uncertainty increased the predicted vulnerability for both high and low fecundity populations assessed using correlated demographic parameters (Table 2; Fig. 4C-D).

Discussion

In this study we use a formal hierarchical Bayesian approach to quantify the trade-offs connecting different demographic rates and reconstruct missing values. Life-history trade-offs constrain the range of possible life-history strategies that can evolve across the tree of life (Lande, 1982; Stearns, 1992). Accordingly, the correlated model used in this study can be applied across the plant and animal kingdom to provide potentially more precise estimates of life-history traits with missing data, compared to values selected from independent surrogate species or populations. We used the reconstructed (i.e., correlated) demographic data to parameterise age-structured models; however, the values can be applied to any analysis of population dynamics based on demographic rates, including population viability analysis, integrated population models, integral projection models and partial differential and ordinary differential equations.

Using the correlated model, we predicted values of adult survival rates for 63 populations of black-legged kittiwakes that are unmonitored for this trait. This output dramatically increases the spatial coverage of populations that have localised estimates for both rates of adult survival and fecundity, especially in areas without intensive monitoring such as the East of England and West Scotland (Fig. 1). Using elasticity analysis, we show that rates of adult survival in this species have a considerably larger demographic impact than rates of fecundity and juvenile survival. This result agrees with life-history theory and highlights the importance of having robust estimates of adult survival rates when conducting impact assessments on

long-lived species, such as seabirds. It may be less critical to use correlated values to fill in missing demographic parameters with low elasticity if assessments are deterministic. However, environmental stochasticity can increase the demographic impact of vital rates with low elasticity (Gaillard, Festa-Bianchet, & Yoccoz, 1998), such that reliable estimates of mean values will become more important if assessment models include demographic rates that vary temporally.

In data-limited circumstances, it is common practise to adopt a pre-cautionary approach to population assessment, and favour the method that provides the maximum estimate of potential mortality (Horswill, O'Brien, et al., 2017). We show that filling in missing data with independent surrogate values does not offer a consistently precautionary approach. Using an independent value that underestimates the missing demographic parameter will overestimate rates of population decline. By contrast, using an independent value that overestimates the missing demographic parameter could seriously underestimate rates of population decline. At present it is not possible for practitioners to ascertain whether independent surrogate values will over or underestimate missing values. However, correlated vital rates reconstructed using life-history trade-offs can provide reference estimates for assessing potential biases.

Population-specific data on other life-history traits, such as age of recruitment and body growth rates, are limited in seabirds. Therefore, we conducted our analysis using two vital rates, i.e., adult survival and fecundity, and a one-dimensional parameter space to represent the life-history trade-off (i.e., covariance between traits). As a result, populations with rates of fecundity close to the national average will have reconstructed rates of adult survival that are also close to the national average. In accordance, national averages could be used to fill in missing data for populations that have life-history strategies very close to the average. However, this will change if the modelling framework is extended to incorporate more than two traits, such that inference becomes multi-dimensional. Multivariate extensions of the correlated model can be easily implemented and are likely to improve the reconstruction of highly fragmented data sets, and where the correlation with other traits is unknown based on life-history theory (e.g. traits that are difficult to empirically measure, such as indeterminate fecundity in batch spawning fish, Horswill et al., 2019). Future reviews and syntheses of how different life-history traits relate to each other by species and groups is likely to be a valuable resource for specifying these models.

In Bayesian statistics, prior distributions can be used to summarise our understanding of how the world works to obtain meaningful inference from small and fragmented data sets (Hobbs & Hooten, 2015). In our correlated model, we used scientific knowledge on the biology of the study species and from life-history theory to inform the shape of the prior distributions assigning certain parameters. In the absence of reference studies, it is still possible to incorporate existing scientific knowledge when defining prior distributions by using biologically or theoretically driven bounds; for example, assigning survival from a uniform distribution bound by 0 and 1, or the correlation parameter from a uniform distribution bound between -1 and 1 (supp. info. Appendix 3). If the direction of the correlation between traits can be predicted

based on life-history theory, the bounds of the prior distribution can be refined accordingly. Using uniform prior distributions with biologically and theoretically driven bounds to assign the expectation of demographic rates and the correlation parameter did not prevent model convergence in our study, but instead increased the posterior credible intervals for 65% of the reconstructed demographic values (supp. info. Appendix 3). By contrast, assigning the variance of adult survival rates in the variance–covariance matrix of the multivariate normal distribution from a uniform prior distribution prevented model convergence. Empirical estimates of adult survival rates were highly limited in our original dataset ($n=5$, Fig. 1), and evidently lacked the information required to quantify variance in this trait. Consequently, correlated models incorporating highly fragmented data are likely to benefit from formulating expectations based on existing data and expertise, i.e., by using informative prior distributions on the relevant parameters. We recommend defining these prior distributions to reflect predictions from life-history theory. For example, amongst populations of a fast-lived species, rates of fecundity are expected to exhibit lower variance than rates of survival (Lande, 1982; Stearns, 1989).

We incorporated demographic stochasticity into the projection analysis. In large populations, environmental stochasticity is usually a dominant source of demographic variation, however in populations of less than 100 individuals, demographic variation is mostly driven by demographic stochasticity (Lande et al., 2003). Instead of explicitly modelling demographic stochasticity, deterministic thresholds are often used to predict extinction, i.e., critical densities beyond which a population can no longer persist (Bascompte, 2003). Extinction thresholds are difficult to define because they will vary by taxon, mating system and local environmental pressures (Engen et al., 2003; Lande et al., 2003; Sæther et al., 2013). Consequently, incorporating demographic stochasticity into population assessments provides a more mechanistic approach for modelling processes at low density, thus reducing uncertainty in how small or declining populations will respond to environmental pressures.

Other rule-based methods of impact assessment, such as Potential Biological Removal (PBR), have lower data requirements than matrix (e.g. age-structured) models, making them quite appealing in data-limited situations (Dillingham & Fletcher, 2008; Wade, 1998). However, the algorithms underpinning rule-based methods often rely on implicit assumptions that limit their application (Niel & Lebreton, 2005). Consequently, rule-based methods are considered unsuitable for environmental impact assessments, especially on seabirds (Green, Langston, McCluskie, Sutherland, & Wilson, 2016; Miller et al., 2019; O'Brien et al., 2017). Alternative approaches for overcoming data limitations in population assessments include using phylogenetic and spatial distance to classify species with similar traits and conservation status (Jetz & Freckleton, 2015). Phylogeny may not be a strong predictor of species-level demographic parameters (Che-Castaldo, Che-Castaldo, & Neel, 2018), and populations in close proximity may not share similar vital rates if environmental pressures differ, such as density dependence or predation (supp. info. Table S1). Furthermore, phylogeny becomes less informative when considering population, as opposed to species-level strategies, and when examining closely related species within the same genus.

Including density-dependent regulation in population assessment models remains debated (Green et al., 2016; Horswill, O'Brien, et al., 2017; O'Brien et al., 2017). In our study, including compensatory density-dependence would have slowed the rate of population decline, but the relative differences between simulations based on independent and correlated demographic parameters would not have changed. Therefore, whilst compensatory and depensatory mechanisms may be important for determining how a specific species or population will respond to environmental pressures, they will not influence a comparison between correlated and independent demographic parameters. We also used baseline (i.e., mean) demographic parameters in our projection analysis. This approach is common for conducting population and environmental impact assessments (Hernández-Camacho et al., 2015; Johnson et al., 2010; O'Brien et al., 2017; Peck, Faulquier, Pinet, Jaquemet, & Le Corre, 2008), however temporal variance and covariance between demographic rates can contribute substantially to population growth (Coulson, Gaillard, & Festa-Bianchet, 2005; Doak, Morris, Pfister, Kendall, & Bruna, 2005). Positive temporal covariance is reported in seabird demographic rates, whereby higher fecundity and survival reinforce each other within a given year (Davis, Nager, & Furness, 2005; Orzack, Steiner, Tuljapurkar, & Thompson, 2011). Here, survival and fecundity are either elevated in an environmentally favourable year, when there are no detectable survival costs of reproduction, or reduced in an environmentally adverse year, when reproduction incurs a survival cost. The correlated model used in this study could be extended to reconstruct full time series of demographic rates, as opposed to just mean values. Achieving this would require separate covariance structures to estimate temporal covariance and life-history trade-offs between traits, similar to the structures previously developed to account for differences in the causal factors influencing life-history trade-offs at the species and population level (Horswill et al., 2019). If data are available, separate covariance structures may also be required to account for age-specific differences in carryover effects associated with adverse environmental conditions (Horswill et al., 2014; Horswill, Trathan, & Ratcliffe, 2017).

In addition to supporting internationally important breeding populations of seabirds (Mitchell, Newton, Ratcliffe, & Dunn, 2004), North-West Europe also provides the best wind resources in the region (Wind Europe, 2020). Extensive development of offshore wind farms is planned to take place within European waters over the next decade (Corbetta, Ho, & Pineda, 2015; OGL, 2019), and mortality due to turbine collision has been identified as a serious potential threat to many species of seabird, including black-legged kittiwakes (Bradbury et al., 2014; Dias et al., 2019; Furness, Wade, & Masden, 2013). International legislation requires that most European countries, granting or refusing consent for the construction of wind farms, conduct a full assessment of the possible impacts to local and regional bird populations (EU Birds Directive 2009/147/EC, EU Habitat Directive 92/43/EEC, EU EIA Directive 85/337/EEC). However, at present, large amounts of missing demographic data are contributing to uncertainty in these impact assessments (Ruffino et al., 2020). Whilst it may be possible to estimate a population mean for rates of fecundity using relatively short time series of data, collecting enough mark-recapture data to robustly estimate rates of adult survival may not be attainable within the consenting process of a wind farm (Horswill

et al., 2018). Therefore, the correlated demographic parameters we detail for black-legged kittiwakes have immediate application for improving environmental management decisions relating to the construction of wind farms in UK and Irish waters. We recommend that impact assessments on data-limited populations evaluate potential future risk by comparing predictions based on independent and correlated demographic parameters, as well as models that propagate correlated parameter uncertainty through to the population predictions. The reconstruction technique applied in this study can easily be adapted to include additional populations. It can also easily be refocused to another species. It is important to highlight that applications to more data limited species may require stronger prior distributions (Appendix 3). For example, we recommend using the posterior distribution of the correlation parameter reported in this study to set the corresponding prior distribution in correlation models of seabird species with fewer estimates of adult survival. Being able to benefit from several sources of information is a key strength of the reconstruction approach applied in this study, which also highlights the importance of maintaining an extensive network of long-term monitoring data sets to support future applications.

Authors' Contributions

CH, JM and AM contributed to the design and planning of this study; FD, MN, SW, MW contributed data; CH conducted all analyses and wrote the first draft of the manuscript, and all authors contributed substantially to revisions and gave final approval for publication.

Acknowledgments

This work was partially funded by Research England to CH, Marine Alliance for Science and Technology Scotland grant SG411 to CH; UK Natural Environmental Research Council grant NE/P004180/1 to JM and CH, UK Natural Environmental Research Council grant NE/R016429/1 as part of the UK-SCaPE programme delivering National Capability to FD, funding from the UK Joint Nature Conservation Committee (DEFRA) to MW and FD. We thank the organisations and individuals who have supported, developed and directed the long-term data collection of black-legged kittiwakes, including but not limited to the Seabird Monitoring Programme, the Wildlife Trust of South and West Wales and NatureScot. We also extend thanks to Dr Julie Miller for discussions regarding impact assessments on black-legged kittiwakes, Dr Sue O'Brien for providing comments on an earlier version of the manuscript, as well as Professor Philip Stephens, Dr Bret Elder, Dr William H. Morgan and one anonymous reviewer for providing comments during the review process.

Data Availability Statement

Input data and the script for the correlated model available via the Dryad Digital Repository <https://doi.org/10.5061/dryad.qnk98sfg0> (Horswill, et al., 2021). All data and sources are also listed in the Supplementary Information Table S1.

References

- Bakewell, A. T., Davis, K. E., Freckleton, R. P., Isaac, N. J. B., & Mayhew, P. J. (2020). Comparing life histories across taxonomic groups in multiple dimensions: How mammal-like are insects? *American Naturalist*, 195(1), 70–81. doi: 10.1086/706195
- Bascompte, J. (2003). Extinction thresholds: insights from simple models. *Annales Zoologici Fennici*, 40(2), 99–114.
- Bennett, P. M., & Owens, I. P. F. (2002). *Evolutionary ecology of birds: Life histories, mating systems and extinction*. Oxford: Oxford University Press.
- Birdlife International. (2019). *Rissa tridactyla*. The IUCN Red List of Threatened Species 2019. Downloaded on 8 May 2020.
- Bradbury, G., Trinder, M., Furness, B., Banks, A. N., Caldow, R. W. G., & Hume, D. (2014). Mapping seabird sensitivity to offshore wind farms. *PLoS ONE*, 9(9), 1–17. doi: 10.1371/journal.pone.0106366
- Cam, E., Cooch, E. G., & Monnat, J.-Y. (2005). Earlier recruitment or earlier death? On the assumption of equal survival in recruitment studies. *Ecological Monographs*, 75(3), 419–434.
- Caswell, H. (2001). *Matrix population models: construction, analysis and interpretation*. Sunderland: Sinauer Associates Inc.
- Che-Castaldo, J., Che-Castaldo, C., & Neel, M. C. (2018). Predictability of demographic rates based on phylogeny and biological similarity. *Conservation Biology*, 32(6), 1290–1300. doi: 10.1111/cobi.13135
- Corbetta, G., Ho, A., & Pineda, I. (2015). *Wind energy scenarios for 2030*. Brussels: Wind Europe (European Wind Energy Association).
- Coulson, J. C. (2011). *The Kittiwake*. London: T & AD Poyser.
- Coulson, T., Gaillard, J. M., & Festa-Bianchet, M. (2005). Decomposing the variation in population growth into contributions from multiple demographic rates. *Journal of Animal Ecology*, 74(4), 789–801. doi: 10.1111/j.1365-2656.2005.00975.x
- Davis, S. E., Nager, R. G., & Furness, R. W. (2005). Food availability affects adult survival as well as breeding success of parasitic jaegers. *Ecology*, 86(4), 1047–1056.
- Desprez, M., Pradel, R., Cam, E., Monnat, J. Y., & Gimenez, O. (2011). Now you see him, now you don't: Experience, not age, is related to reproduction in kittiwakes. *Proceedings of the Royal Society B: Biological Sciences*, 278(1721), 3060–3066. doi: 10.1098/rspb.2011.0189
- Devenish-Nelson, E. S., Harris, S., Soulsbury, C. D., Richards, S. A., & Stephens, P. A. (2013). Demography of a carnivore, the red fox, *Vulpes vulpes*: What have we learnt from 70 years of published studies? *Oikos*, 122(5), 705–716. doi: 10.1111/j.1600-0706.2012.20706.x
- Dias, M. P., Martin, R., Pearmain, E. J., Burfield, I. J., Small, C., Phillips, R. A., ... Croxall, J. P. (2019). Threats to seabirds: A global assessment. *Biological Conservation*, 273, 525–537. doi: 10.1016/j.biocon.2019.06.033

- Dillingham, P. W., & Fletcher, D. (2008). Estimating the ability of birds to sustain additional human-caused mortalities using a simple decision rule and allometric relationships. *Biological Conservation*, 141(7), 1783–1792. doi: 10.1016/j.biocon.2008.04.022
- Doak, D. F., Morris, W. F., Pfister, C., Kendall, B. E., & Bruna, E. M. (2005). Correctly estimating how environmental stochasticity influences fitness and population growth. *The American Naturalist*, 166(1), E14–E21. doi: 10.1086/430642
- Engen, S., Lande, R., & Sæther, B.-E. (2003). Demographic stochasticity and Allee effects in populations with two sexes. *Ecology*, 84, 2378–2386.
- Frederiksen, M., Harris, M. P., & Wanless, S. (2005). Inter-population variation in demographic parameters: a neglected subject? *Oikos*, 111(2), 209–214.
- Frederiksen, M., Wanless, S., Harris, M. P., Rothery, P., & Wilson, L. J. (2004). The role of industrial fisheries and oceanographic change in the decline of North Sea black-legged kittiwakes. *Journal of Applied Ecology*, 41(6), 1129–1139.
- Furness, R. W., Wade, H. M., & Masden, E. A. (2013). Assessing vulnerability of marine bird populations to offshore wind farms. *Journal of Environmental Management*, 119, 56–66. doi: 10.1016/j.jenvman.2013.01.025
- Gaillard, J. M., Festa-Bianchet, M., & Yoccoz, N. G. (1998). Population dynamics of large herbivores: Variable recruitment with constant adult survival. *Trends in Ecology and Evolution*, 13(2), 58–63. doi: 10.1016/S0169-5347(97)01237-8
- Githiru, M., Matthysen, E., Lens, L., & Bennun, L. A. (2007). Can a common bird species be used as a surrogate to draw insights for the conservation of a rare species? A case study from the fragmented Taita Hills, Kenya. *Oryx*, 41, 239–246.
- Green, R. E., Langston, R. H. W., McCluskie, A., Sutherland, R., & Wilson, J. D. (2016). Lack of sound science in assessing wind-farm impacts on seabirds. *Journal of Applied Ecology*, 53(6), 1635–1641. doi: 10.1111/1365-2664.12731
- Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-gómez, R., & Buckley, Y. M. (2019). Animal life history is shaped by the pace of life and the distribution of age-specific mortality and reproduction. *Nature Ecology and Evolution*, 3, 1217–1224. doi: 10.1038/s41559-019-0938-7
- Hernández-Camacho, C. J., Bakker, V. J., Aurióles-Gamboa, D., Laake, J., & Gerber, L. R. (2015). Use of surrogate data in demographic population viability analysis: A case study of California sea lions. *PLoS ONE*, 10(9), 1–16. doi: 10.1371/journal.pone.0139158
- Hobbs, N. T., & Hooten, M. B. (2015). Informative priors. In *Bayesian Models: A Statistical Primer for Ecologists* (STU-Stud, pp. 97–101). Princeton; Oxford: Princeton University Press.
- Horswill, C., Dulvy, N. K., Juan-Jordá, M. J., Kindsvater, H. K., Mangel, M., & Matthiopoulos, J. (2019). Global reconstruction of life-history strategies: A case study using tunas. *Journal of Applied Ecology*, 56, 855–865. doi: 10.1111/1365-2664.13327

- Horswill, C., Humphreys, E. M., & Robinson, R. A. (2018). When is enough...enough? Effective sampling protocols for estimating the survival rates of seabirds with mark-recapture techniques. *Bird Study*, 65(3), 290–298. doi: 10.1080/00063657.2018.1516191
- Horswill, C., Matthiopoulos, J., Green, J. A., Meredith, M. P., Forcada, J., Peat, H. J., ... Ratcliffe, N. (2014). Survival in macaroni penguins and the relative importance of different drivers: individual traits, predation pressure and environmental variability. *Journal of Animal Ecology*, 83(5), 1057–1067. doi: 10.1111/1365-2656.12229
- Horswill, C., O'Brien, S. H., & Robinson, R. A. (2017). Density dependence and marine bird populations: are wind farm assessments precautionary? *Journal of Applied Ecology*, 54(5), 1406–1414. doi: 10.1111/1365-2664.12841
- Horswill, C., Manica, A., Daunt, F., Newell, M., Wanless, S., Wood, M., Matthiopoulos, J. (2021) Data from: Improving assessments of data-limited populations using life-history theory. Dryad Digital Repository. <https://doi.org/10.5061/dryad.qnk98sfg0>
- Horswill, C., Ratcliffe, N., Green, J. A., Phillips, R. A., Trathan, P. N., & Matthiopoulos, J. (2016). Unravelling the relative roles of top-down and bottom-up forces driving population change in an oceanic predator. *Ecology*, 97(8), 1919–1928. doi: 10.1002/ecy.1452
- Horswill, C., & Robinson, R. A. (2015). Review of seabird demographic rates and density dependence. JNCC Report No. 552. Peterborough: Joint Nature Conservation Committee.
- Horswill, C., Trathan, P. N., & Ratcliffe, N. (2017). Linking extreme interannual changes in prey availability to foraging behaviour and breeding investment in a marine predator, the macaroni penguin. *PLoS ONE*, 12(9), 1–12. doi: 10.1371/journal.pone.0184114
- Jeschke, J. M., & Kokko, H. (2009). The roles of body size and phylogeny in fast and slow life histories. *Evolutionary Ecology*, 23(6), 867–878. doi: 10.1007/s10682-008-9276-y
- Jetz, W., & Freckleton, R. P. (2015). Towards a general framework for predicting threat status of data-deficient species from phylogenetic, spatial and environmental information. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370(1662), 20140016–20140016. doi: 10.1098/rstb.2014.0016
- Johnson, H. E., Mills, L. S., Stephenson, T. R., & Wehausen, J. D. (2010). Population-specific vital rate contributions influence management of an endangered ungulate. *Ecological Applications*, 20(6), 1753–1765. doi: 10.1890/09-1107.1
- Kellner, K. (2016). Package 'jagsUI' v 1.4.4. CRAN.
- Kindsvater, H. K., Dulvy, N. K., Horswill, C., Juan-Jordá, M.-J., Mangel, M., & Matthiopoulos, J. (2018). Overcoming the data crisis in biodiversity conservation. *Trends in Ecology and Evolution*, 33, 676–688. doi: 10.1016/j.tree.2018.06.004
- Lande, R., Engen, S., & Sæther, B.-E. (2003). *Stochastic population dynamics in ecology and conservation*. Oxford: Oxford University Press.
- Lande, Russell. (1982). A quantitative genetic theory of life history evolution. *Ecology*, 63(3), 607–615.

- Miller, J. A. O., Furness, R. W., Trinder, M., & Matthiopoulos, J. (2019). The sensitivity of seabird populations to density - dependence, environmental stochasticity and anthropogenic mortality. *Journal of Applied Ecology*, (56), 2118–2130. doi: 10.1111/1365-2664.13448
- Mitchell, P. I., Newton, S. F., Ratcliffe, N., & Dunn, T. E. (2004). *Seabird populations of Britain and Ireland*. London: T. & A.D. Poyser.
- Niel, C., & Lebreton, J. D. (2005). Using demographic invariants to detect overharvested bird populations from incomplete data. *Conservation Biology*, 19(3), 826–835. doi: 10.1111/j.1523-1739.2005.00310.x
- O'Brien, S. H., Cook, A. S. C. P., & Robinson, R. A. (2017). Implicit assumptions underlying simple harvest models of marine bird populations can mislead environmental management decisions. *Journal of Environmental Management*, 201, 163–171. doi: 10.1016/j.jenvman.2017.06.037
- OGL. (2019). *Industrial strategy: Offshore Wind Sector Deal*. London.
- Orzack, S. H., Steiner, U. K., Tuljapurkar, S., & Thompson, P. (2011). Static and dynamic expression of life history traits in the northern fulmar *Fulmarus glacialis*. *Oikos*, 120(3), 369–380. doi: 10.1111/j.1600-0706.2010.17996.x
- Pearson, R. G., Stanton, J. C., Shoemaker, K. T., Aiello-Lammens, M. E., Ersts, P. J., Horning, N., ... Akçakaya, H. R. (2014). Life history and spatial traits predict extinction risk due to climate change. *Nature Climate Change*, 4(3), 217–221. doi: 10.1038/nclimate2113
- Peck, D. R., Faulquier, L., Pinet, P., Jaquemet, S., & Le Corre, M. (2008). Feral cat diet and impact on sooty terns at Juan de Nova Island, Mozambique Channel. *Animal Conservation*, 11(1), 65–74. doi: 10.1111/j.1469-1795.2007.00153.x
- R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Ruffino, L., Thompson, D., & O'Brien, S. H. (2020). *Black-legged kittiwake population dynamics and drivers of population change in the context of offshore wind development*. Peterborough: Joint Nature Conservation Committee.
- Sæther, B.-E., Coulson, T., Grøtan, V., Engen, S., Altwegg, R., Armitage, K. B., ... Weimerskirch, H. (2013). How life history influences population dynamics in fluctuating environments. *The American Naturalist*, 182(6), 743–759. doi: 10.1086/673497
- Salguero-Gómez, R., Jones, O. R., Jongejans, E., Blomberg, S. P., Hodgson, D. J., Mbeau-Ache, C., ... Buckley, Y. M. (2016). Fast–slow continuum and reproductive strategies structure plant life-history variation worldwide. *Proceedings of the National Academy of Sciences*, 113(1), 230–235. doi: 10.1073/pnas.1506215112
- Schtickzelle, N., WallisDeVries, M. F., & Baguette, M. (2005). Using surrogate data in population viability analysis: The case of the critically endangered cranberry fritillary butterfly. *Oikos*, 109(1), 89–100. doi: 10.1111/j.0030-1299.2005.13745.x
- SMP. (2020). These data were extracted from the Seabird Monitoring Programme Database (<https://app.bto.org/seabirds>). Data have been provided to the SMP by the generous contributions of nature conservation and research organisations, and many volunteers throughout Britain and Ireland.

- Accepted Article
- Stearns, S. C. (1989). Trade-offs in life-history evolution. *Functional Ecology*, 3(3), 259–268. doi: 10.2307/2389364
- Stearns, S. C. (1992). *The Evolution of Life Histories*. New York: Oxford University Press.
- Stubben, C., Milligan, B., & Nantel, P. (2016). *popbio: Construction and Analysis of Matrix Population Models*. CRAN.
- Thorson, J. T., Munch, S. B., Cope, J. M., & Gao, J. (2017). Predicting life history parameters for all fishes worldwide. *Ecological Applications*, 27, 2262–2276. doi: 10.1002/eap.1606
- Wade, P. R. (1998). Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Marine Mammal Science*, 14(1), 1–37. doi: 10.1111/j.1748-7692.1998.tb00688.x
- Walsh, P. M., Halley, D. J., Harris, M. P., Del Nevo, A., Sim, I. M. W., & Tasker, M. L. (1995). *Seabird monitoring handbook for Britain and Ireland. A compilation of methods for survey and monitoring of breeding seabirds*. Peterborough: Joint Nature Conservation Committee.
- Wind Europe. (2020). *Offshore wind in Europe: Key trends and statistics 2019*. Brussels.
- Winemiller, K. O. (2005). Life history strategies, population regulation, and implications for fisheries management. *Canadian Journal of Fisheries and Aquatic Sciences*, 62(4), 872–885. doi: 10.1139/f05-040

Tables

Table 1. The four projection scenarios (PS) used to examine the potential biases and imprecisions generated by assessing populations with independent and correlated demographic parameters.

	Independent demographic parameters	Correlated demographic parameters
Without parameter uncertainty	PS 1	PS 2
With parameter uncertainty	PS 3	PS 4

Table 2. Percentage of model iterations from the four projection scenarios that resulted in vulnerability or populations increasing from the initial size.

Projection scenario	Vulnerability		Increase	
	Low fecundity	High fecundity	Low fecundity	High fecundity
1. Independent demographic parameters	37.6	0.0	0.0	55.6
2. Correlated demographic parameters	0.0	0.0	0.0	0.0
3. Independent demographic parameters with parameter uncertainty	38.4	0.0	0.0	47.4
4. Correlated demographic parameters with parameter uncertainty	9.0	4.9	0.4	14.9

Figures



Figure 1. The UK and Ireland has considerably fewer sites where populations of black-legged kittiwakes are monitored for rates of adult survival and fecundity (black triangles), compared to sites where only rates of fecundity are monitored (red circles). Projection WGS84.




Figure 2. The cumulative posterior distribution of the correlated demographic parameters. Median posterior point estimates for reconstructed (Rec, i.e., correlated) rates of adult survival and fecundity shown on the scale of the linear predictor. Populations with empirical data for survival and fecundity are black triangles, and populations with empirical data for fecundity only are red circles. Median posterior values align with the imputed correlation between the two demographic rates.




Figure 3. A) Correlation between observed population-specific values of mean adult survival rates and the median reconstructed values from our correlated model. B) Correlation between observed population-specific values of mean fecundity and median reconstructed values from our correlated model. For reference, we show the 95% credible intervals for all reconstructed (i.e., correlated) values, and the 1:1 relationship between observed and reconstructed values as a grey diagonal line. We detail the ten populations selected for the population assessment projections (Table 1) as open symbols in Panel B. Populations with empirical data for survival and fecundity are black triangles, and populations with empirical data for fecundity only are red circles.




Figure 4. A) The final population sizes obtained from the projection analysis with different rates of fecundity strongly differed when using independent demographic parameters (projection scenario 1, Table 1). B) This difference was smaller when assessments were based on median posterior values from our correlated model (projection scenario 2, Table 1). (C) Propagating parameter uncertainty that was independent between rates of adult survival and fecundity (projection scenario 3, Table 1) decreased predicted vulnerability in the low fecundity group, compared to projection scenario 1, and produced a large range of possible outcomes for the high fecundity group. (D) Propagating parameter uncertainty that maintained observed levels of covariance between traits increased the range of possible outcomes and increased predicted vulnerability in both the low and high fecundity groups, compared to projection scenario 2 (projection scenario 4, Table 1). Initial population size for simulations show as horizontal grey dashed lines.