

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document, © 2022 -IOS Press. All rights reserved. and is licensed under All Rights Reserved license:

Ryan, Jennifer M., Lavelle, Grace, Noorkoiv, Marika, Kiernan, Damien, Theis, Nicola ORCID logoORCID: https://orcid.org/0000-0002-0775-1355, Little, Úna, Shortland, Adam and Malone, Ailish (2022) Associations between gait kinematics, gross motor function and physical activity among young people with cerebral palsy: A cross sectional study. Physiotherapy Practice and Research, 43 (1). pp. 71-80. doi:10.3233/PPR-200411

Official URL: http://doi.org/10.3233/PPR-200411 DOI: http://dx.doi.org/10.3233/PPR-200411 EPrint URI: https://eprints.glos.ac.uk/id/eprint/11286

## Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

# Associations between gait kinematics, gross motor function and physical activity among young people with cerebral palsy: A cross sectional study

Jennifer M. Ryan, Grace Lavelle, Marika Noorkoiv, Damien Kiernan, Nicola Theis, Una Little, Adam Shortland, and Ailish Malone

#### Abstract.

**INTRODUCTION:** The aim of this study was to investigate the association between gait parameters, gross motor function and physical activity (PA) in young people with cerebral palsy (CP).

**METHODS:** Thirty-eight adolescents aged between 10-19 years with spastic CP in GMFCS levels I-ill (mean [standard deviation] age 13.7 [2.4] yr; 53% female) were included in this cross-sectional study. Hip, knee and ankle joint excursion and stance time was assessed using 3D gait analysis. Self-selected walking speed was assessed during a timed 10 m overground walk and treadmill walking. Gross motor function was assessed using dimensions D and E of the Gross Motor Function Measure (GMFM-66). Moderate-to-vigorous PA, light PA and step-count were assessed using an accelerometer. Linear regression was used to examine associations.

**RESULTS:** After adjusting for age, sex and GMFCS level, percentage stance time was associated with dimension E of the GMFM-66 (/3 =-0.29, 95% CI-0.54 to -0.05). There was no evidence that any other gait parameters were associated with GMFM-66 dimensions D or E. There was also no evidence that gait parameters or GMFM-66 dimensions D or E were associated with step-count or time in PA after adjusting for age, sex and GMFCS level.

**DISCUSSION:** The findings provide an insight into the complexity of the relationship between gait quality or ability at the impairment level, function as measured in a controlled environment, and the performance of habitual PA, which is essential for health among children with CP.

Keywords: Developmental neurology and neurodisability, functional performance, gait, gait analysis, mobility limitation, paediatrics

## 1. Introduction

Cerebral palsy (CP) is a permanent, non-progressive disorder associated with an impairment of the developing fetal or infant brain [1]. It is a common cause of childhood physical disability worldwide and is characterised by abnormal fine and gross motor functioning [1]. In addition to abnormal tone, people with CP experience reduced muscle strength and aerobic fitness, musculoskeletal disorders, and often participate in low levels of physical activity (PA) [1-3]. Management of CP encompasses medical, surgical and therapeutic interventions, which typically aim to improve functioning as described by the World Health Organisation's International Classification of Functioning, Disability and Health (ICF). According to the ICF, functioning refers to body structures and functions, activity and participation, where activity is described as the execution of a task by an individual and participation is involvement in a life situation [4].

Improving mobility is often a primary therapeutic goal for people with CP [5]. Many interventions to improve mobility target impairments in body structures and functions in order to optimise activity and participation. However, the effects of such treatments are often evaluated on only one level of the ICF [6, 7]. For example, the effect of orthopaedic surgery on mobility is often examined by assessing gait parameters, as a measure of body structures and functions [7]. Whereas the effect of exercise on mobility is often examined by assessing gross motor function, as a measure of activity [6]. Further, the effect of interventions on mobility in the context of a person's usual environment is rarely examined [6, 7]. As functioning of an individual is context-dependent [4], a person's mobility in a controlled environment (i.e., activity capacity) may differ to their mobility in their daily environment (i.e., activity performance) [8]. Guide-lines in the United Kingdom recommend that children should participate in an average of at least 60 minutes of moderate-to-vigorous intensity physical activity (MVPA) per day across the week [9]. Although PA of any intensity can provide health benefits, MVPA is needed to achieve the maximum health benefits [9]. As many children with CP do not achieve sufficient MVPA [3], understanding the potential association between gait parameters, function and MVPA warrants further consideration.

The literature investigating the associations between gait parameters, gross motor function, and PA performance in a person's daily environment is relatively limited. To date, there has been no investigation of the relationship between gait kinematics and both gross motor function and PA in the same cohort of children with CP. In relation to the association between gait parameters and gross motor function, Robinson found that the Gait Profile Score (GPS) differed between children in Gross Motor Function Classification System (GMFCS) levels I, II and III [10], with more impaired gait kinematics seen in those with poorer functional mobility. Molloy found a strong correlation (r = 0. 70) between the Gait Deviation Index and total score on the Gross Motor Function Measure (GMFM) among children aged 4-17 yr in GMFCS levels I-IV [ 11]. Damiano and Abel found cadence, normalized velocity, hip knee excursion and percentage single support were associated with score on the GMFM among children aged [12]. Two studies have examined the association between gait kinematics and PA. Guinet and Desailly reported a correlation of 0.41 (Spearman 's rho) between the Gait Deviation Index (GDI) and step count in 25 adolescents with CP in GMFCS level I-II [13]. Wilson reported a stronger correlation of 0.58 between the GDI and step count in a larger sample of 55 children and adolescents aged 6 to 18 years [14]. However, neither study found an association between GDI and time in MVPA. Guinet and Desailly reported fair correlations between MVPA and key kinematic parameters at heel strike and toe off (spearman's rho 0.3 to 0.33) [13]. However, these specific gait cycle points give a limited snapshot of an individual's gait deviation.

Given that the ICF places an emphasis on the interaction between body functions and activity, understanding the association between gait parameters, gross motor function and habitual PA is important for awareness of the impact of CP on the individual and identification of appropriate interventions. However, there is a lack of evidence regarding the association between gait and gross motor function, and gait and habitual PA, respectively, in a single cohort of people with CP.

The aim of this study was to investigate the association between gait parameters, gross motor function and PA in young people with CP.

#### 2. Methods

Data for this cross-sectional study were obtained from the baseline assessment of a randomised controlled trial that aimed to investigate the effects of strength training for adolescents with CP [15]. Recruitment took place between August 2015 and May 2017. Participants were recruited throughout England from eight National Health Service (NHS) trusts, a special education needs school, a University, a primary care organisation, national organisations for people with disabilities, and by word of mouth. Inclusion criteria were a diagnosis of spastic CP, aged 10-19 years, ability to walk independently with or without a mobility aid (i.e., GMFCS levels I-III), and an ability to activate the ankle plantarflexors as determined by palpation. Adolescents were excluded if they had orthopaedic surgery of the lower limbs in the past 12 months, had botulinum toxin type A injections or serial casting in the past 6 months, or had insufficient cognition to comply with assessment procedures and the training programme delivered as part of the trial. Participants aged 16 years and over gave written informed consent. Participants under 16 years of age gave written informed assent and a parent or guardian provided written informed consent. The trial was approved by Brunel University Lon-don's College of Health and Life Sciences Research Ethics Committee and the Surrey Borders Research Ethics Committee (ref: 15/LO/0843).

#### 2.1. Gait parameters

The gait parameters examined in this study were total sagittal plane excursion of the hip, knee and ankle (i.e., the difference between maximal and minimal angles over one full gait cycle), percentage stance time, and self-selected walking speed. These parameters were selected in order to provide clinical information relevant to joint movement during gait. Reduced sagittal excursion at the hip, knee and ankle has been shown in pathological gaits in CP, for example, at the knee and ankle in crouch [16] and at the hip, knee and ankle in children with hemiplegia who also show excessive muscle coactivation [17]. Variation in gait patterns was expected among participants, given the inclusion of adolescents in GMFCS levels I, II and II. Therefore, analysis of total sagittal plane excursion was preferred over joint angles at single gait cycle points, which could differ considerably depending on the participant's pattern:

Kinematic data during treadmill walking were collected using a computerized motion capture system (Motion Analysis, Motion Analysis Corporation, Santa Rosa, CA, USA) with 8 infrared cameras. Two researchers with over 5 years of experience each in 3D motion capture research and PhDs in closely related areas collected the data. The Motion Analysis soft-ware Cortex was used for the processing of kinematic data (150 Hz).

Participants were asked to walk on a fully instrumented treadmill (Bertec Corporation, Columbus, Ohio) at their preferred walking speed. Prior to testing, participants performed a familiarisation session to ensure they were comfortable walking on the treadmill, and to establish a comfortable preferred treadmill walking speed as previously described [18]. Following a mandatory two minute rest after familiarization, participants were asked to walk on the treadmill at their preferred walking speed for two minutes to ensure a minimum of 30 gait cycles were collected for further analysis [19].

Knee, ankle and hip joint kinematics were measured from a body motion analysis marker set. For this purpose, reflective markers were placed on the anterior superior iliac spine, sacrum, the greater trochanters, mid-thigh, medial and lateral femoral epicondyles, on each tibia (midway between the ankle and knee), the medial and lateral malleoli, the heads of the first and fifth metatarsals and the calcaneus. Kinematics from the most affected leg, as reported by the participant, were calculated in Visual 3D software using the conventional gait model [20], with a modification at the foot. Where the participant reported that both sides were equally affected, data from the right leg were used. Joint angles were computed as the angles between the proximal and distal segment of the relevant joint within the primary axis of the foot segment defined as the line between the ankle joint centre (rather than the calcaneus) and the mid-point of the first and fifth metatarsal heads, as previously described [21.]. The foot was calculated in this way due to intermittent occlusion of the calcaneus marker by the treadmill apparatus, but the use of the ankle joint centre did not affect total ankle joint range.

In addition to gait parameters collected during treadmill walking, self-selected overground walking speed was measured as this is potentially a more feasible method of assessing gait during routine clinical practice. Preferred walking speed during two minutes of treadmill walking was recorded. Participants were instructed to walk at a self-selected speed over approximately 15 metres. The time taken to walk 10 metres was recorded. Participants repeated this three times and the average speed of the three trials was used in analysis. Walking speed normalised to height was used in analyses.

#### 2.2 Gross motorfunction

Gross motor function was assessed using dimensions D and E of the GMFM-66, which were administered by two physiotherapists and video-recorded. A specialist paediatric physiotherapist, with training and experience of scoring the GMFM-66, scored performance. The GMFM-66 is valid and reliable in children with CP [22, 23]. Dimension D evaluates activities in standing. Dimension E evaluates activities in walking, running and jumping. A higher score indicates better gross motor function.

#### 2.3 Physical activity

Daily light PA, moderate-to-vigorous PA and step-count were measured using an Actigraph wGT3X accelerometer (Actigraph, USA) worn on the waist above the right hip or least affected side in the case of significant asymmetry, in the midaxillary line. Participants were asked to wear the accelerometer for 7 consecutive days. Participants with at least 2 days of monitoring were included in the analysis as two days is necessary to achieve a reliability coefficient of 0. 70 for adolescents with CP [24]. Data were analysed using Actilife Software. Non-wear time was identified using an algorithm developed by Choi [25]. Non-wear time was identified as a period of 90 minutes of no movement with a spike tolerance of two minutes. Cut-points validated in children and adolescents with CP were applied to

#### 2.4 Statistical analysis

identify time spent in light physical activity (LPA) and MVPA [26].

Data were analysed using Stata version 15.0 (StatCorp LLC, TX, USA). Distribution of data was explored using histograms and Q-Q plots. We removed GMFM-66 D score for one participant with an outlying value, 5 standard deviations from the mean. Continuous data were summarised as mean and standard deviation or median and interquartile range if data were not normally distributed. Categorical variables were summarised as frequency counts. Linear regression was used to explore associations between gait parameters (i.e., hip, knee and joint excursion, percentage stance time, normalised over-ground and treadmill walking speed), gross motor function (i.e., GMFM-66 dimension D and E), and PA (i.e. time in MVPA and LPA, and step- count). Uni-variable models (i.e., linear regression models with one dependent variable and one independent variable) were fitted to examine separate associations between each gait parameter (i.e., independent variable) and GMFM-66 D score,

GMFM-66 E score, time in LPA, time in MVPA, and step-count (i.e., dependent variables). Multivariable models (i.e., linear regression models with one dependent variable and more than one independent variable) were fitted to examine the associations between each independent variable and each dependent variable after adjusting for age, sex and GMFCS level. We explored the effect of additionally adjusting for distribution but chose not to include it in final models because: there was no evidence that it changed the coefficient for the independent variable of interest by> 10% after age, sex and GMFCS level had been adjusted for. To explore if gross motor function and gait parameters jointly explained variation in PA, we fitted three linear regression models with GMFM-66 D and E score, hip, knee and ankle joint angle, normalised overground walking speed, and percentage stance time as independent variables and (1) MVPA, (2) LPA and (3) step-count as dependent variables. For each model, F-tests were used to test the null hypothesis that gait parameters and gross motor function were not jointly associated with PA.

Assumptions of linear regression, namely normally distributed residuals, homoscedasticity, and a linear relationship between each independent variable and dependent variable conditional on the other independent variables in the model, were checked using appropriate plots. In models where GMFM-66 D and E score were dependent variables, there was some evidence that residuals were not normally distributed. Therefore, for these models a bootstrapping procedure was used. The bootstrap provides an alternative way to estimate valid standard errors and confidence intervals without relying on assumptions about distributions [27]. It involves calculating  $\beta$  in multiple "bootstrap samples" that are sampled with replacement from the original sample [27]. Bias corrected and accelerated bootstrap confidence intervals (Cls) were calculated from 2,000 replicates [28]. As exact p values are not calculated when a bootstrap-ping procedure is used, the p value associated with each effect estimate was infer ed from the confidence interval (Cl) (i.e., p < 0.05 or p < 0.01 where the 95% Cl or 99% Cl, respectively, did not include the null value).

### 3. Results

Thirty-eight adolescents with complete data on gait parameters were included in the study. Participant characteristics are described in Table 1.

Hip, knee and ankle joint excursion, percentage stance time, self-selected walking speed during over-ground walking and treadmill walking, GMFM-66 D score, GMFM-66 E score, light PA, moderate-to-vigorous PA and stepcount are described in Table 2. Figure I shows hip, knee and ankle kinematics over the gait cycle in the sagittal plane.

Associations between gait parameters and gross motor function are presented in Table 3. In unadjusted analyses, normalised overground walking speed and percentage stance time were associated with dimensions D and E of the GMFM-66, and normalised treadmill walking speed was associated with dimension D of the GMFM-

66. However, after adjusting for age, sex and GMFCS level, only percentage stance time remained associated with dimension E ( $\beta$ =-0.29, 95% CI -0.54 to -0:05). Hip, knee and ankle joint excursion were not associated with dimension D or E in unadjusted or adjusted analysis.

Associations between gait parameters and PA are presented in Table 4. There was no evidence that hip, knee or ankle joint excursion were associated with LPA, MVPA or step-count. There was also no evidence that percentage stance time or normalised treadmill walking speed were associated with LPA, MVPA or step-count. Normalised overground walking speed was associated with daily step-count and time in MVPA, but not LPA, in unadjusted analyses. However, after adjusting for age, sex and GMFCS level there was no evidence that normalised over-ground walking speed was associated with LPA, MVPA or step-count.

When all gait parnmeters, GMFM-66 D and GMFM-66 E score were entered into a regression model as independent variables, hip joint excursion was associated with step-count (p = 0.046; Table 5). A one degree increase in hip excursion was associated with an increase of 133 steps per day. However, there was no evidence that gait parameters and gross motor function were jointly associated with step-count (p = 0.133), MVPA (0.209) or LPA (p = 0.406). In com-bination, gait parameters and gross motor function explained 33% of the variance in step-count (R2 = 0.325), 29% of the variance in MVPA (R 2 = 0.289) and 22% of the variance in LPA (R 2 = 0.225).

P	Table I	
Pa	articipant characteristics	
	n (%)	Mean (SD)
Age, yr	38	13.7 (2.4)
Gender		
Female	20 (53)	
Male	18 (47)	
Height, cm	38	154.5 (13.4)
Mass, kg	38	47.9 (12.9)
Distribution		
Unilateral	22 (58)	
Bilateral	16 (42)	
<b>GMFCS</b> level		
I	19 (50)	
II	17 (45)	
III	2 (5)	
SD: standard deviation	n.	

Table 2 Description of kinematic variables, walking speed, gross motor function and physical activity

	n	Mean (SD)	Range
Hip joint excursion, deg	38	36.2 (9.2)	22.7 to 56.8
Knee joint excursion, deg	38	48.4 (11.2)	15.7 to 77.0
Ankle joint excursion, deg	38	20.6 (7.6)	7.9 to 42.9
Overground walking speed, m.s <sup>-1</sup>	38	1.13 (0.19)	0.72 to 1.47
Normalised overground walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	38	0.75 (0.12)	0.54 to 0.95
Treadmill walking speed, m.s <sup>-1</sup>	37	0.44 (0.13)	0.11 to 0.80
Normalised treadmill walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	37	0.28 (0.09)	0.07 to 0.56
Stance time, %	38	70.5 (6.2)	55.0 to 85.0
GMFM-66 Dimension Da	37	36.2 (3.2)	22.0 to 39.0
GMFM-66 Dimension E <sup>a</sup>	38	67.5 (7.0)	12.0 to 72.0
Light physical activity, min.day <sup>-1</sup>	35	200.4 (52.0)	96.4 to 354.7
Moderate-to-vigorous physical activity, min.day-1	35	59.6 (22.7)	10.0 to 118.6
Step count, steps.day <sup>-1</sup>	35	6365.5 (2123.6)	2218.6.to 11562.8

SD: standard deviation. <sup>a</sup>Data presented as median (interquartile range); <sup>b</sup>normalised to height.

#### 4. Discussion

The results of this study indicated percentage time spent in stance is negatively associated with gross motor function, specifically activities relating to walking, running and jumping. After controlling for age, sex and GMFCS level, a 1% increase in stance time is associated with on average a -0.29 point reduction on GMFM-66 dimension E score. Gait parameters and function were not associated with step-count or time in PA after adjusting for age, sex and GMFCS level. Similarly, in combination, gait parameters and function were not predictive of step-count or time in PA and explained a relatively small proportion of the variance in these outcomes.

Few studies have examined associations between gait parameters, gross motor function and PA. Molloy et al. reported a strong association between the GDI and total score on the GMFM in children and adolescents in GMFCS levels I-IV [11]. However, potentially confounding variables such as age, sex and GMCS level were not adjusting for when examining this association. Similarly to this study, Damiano and Abel reported unadjusted associations between percentage single support, normalised velocity and score on the GMFM among 32 children aged 3-18 years [12]. After adjusting for age, sex and GMFCS level, we found that walking speed did not remain associated with GMFM-66 dimensions D or E. Damiano and Abel also found that hip knee excursion was associated with GMFM score. Lack of agreement between our findings may be due to dif-erences in the age range of the sample and the use of the full GMFM score. The variance of full GMFM score and the variance of hip joint excursion are likely bigger among children aged 3-18 years compared to children aged 10-19 years, and may explain why an association was observed by Damiano and Abel.

We did not find that gait parameters or function were associated with step-count or time in MVPA after adjusting for age, sex and GMFCS level. However, when all gait parameters and GMFM- 66 dimensions D and E were included together in multiple linear regression, mp joint excursion was independently associated with step-count. A one degree increase in mp excursion was associated with an average increase of 133 steps per day. A previous study, examining associations between a set of 54 kinematic and spatio-temporal parameters, step-count and MVPA among adolescents with CP, found that hip flexion at toe off, knee flexion at heel strike and ankle flexion at heel strike, were weakly and negatively correlated with MVPA [13]. The same study found GDI and stance were weakly correlated with step-count but were not correlated with MVPA. A moderate correlation between GDI and step-count has also been observed among children with CP [14]. These findings suggest that gait parameters may be weakly correlated with performance of steps in daily life but not general activity.

However, gait parameters and function still explained relatively little of the variance in step-count. This, in combination with the limited number of associations we observed between gait impairments, function, and PA, emphasise the important contribution of environmental and personal factors to the interaction between body structures and functions and activity, as outlined by the ICF [4]. Associations between impairments and activity limitations, either in a controlled environment or a person's usual environment, are influenced by contextual factors. Motivation and self-efficacy play an important role in participation in PA among adolescents with CP [29, 30], which may be unrelated to impairments. Other barriers include lack of access to appropriate equipment, inadequate staffing within schools, transport to activities, and lack of inclusive sport opportunities [29, 30].



Fig. I. Additional kinematic graphs.

Table 3	
Unadjusted and adjusted associations between gait parameters and gross motor function	

.

	n	(95% CI)	p value <sup>a</sup>	R <sup>2</sup>	Adjusted <sup>a</sup> β (95% CI)	p value"	R <sup>2</sup>
Dependent variable: GMFM-66 Dimension D							
Hip joint excursion, deg	37	-0.06 (-0.33 to 0.09)	> 0.05	0.022	-0.11 (-0.41 to 0.07)	>0.05	0.396
Knee joint excursion, deg	37	-0.002 (-0.15 to 0.07)	>0.05	0.000	-0.03 (-0.15 to 0.03)	>0.05	0.345
Ankle joint excursion, deg	37	0.02 (-0.05 to 0.11)	> 0.05	0.003	0.07 (-0.01 to 0.24)	>0.05	0.360
Normalised overground walking speed, c m.s-1.m-1	37	7.34 (0.10 to 16.74)	< 0.05	0.074	5.38 (-2.14 to 17.99)	>0.05	0.370
Normalised treadmill walking speed, c m.s-1.m-1	36	11.95 (3.03 to 28.82)	< 0.01	0.085	4.14 (-7.87 to 15.61)	> 0.05	0.311
Stance time, %	37	-0.11 (-0.28 to -0.01)	< 0.05	0.047	-0.04 (-0.19, 0.06)	> 0.05	0.342
Dependent variable: GMFM-66 Dimension E							
Hip joint excursion, deg	38	-0.31 (-1.12 to 0.34)	>0.05	0.048	-0.23 (-0.80 to 0.07)	>0.05	0.822
Knee joint excursion, deg	38	0.11 (-0.32 to 0.52)	>0.05	0.009	0.02 (-0.19 to 0.22)	>0.05	0.804
Ankle joint excursion, deg	38	0.04 (-0.49 to 0.34)	>0.05	0.000	0.18 (-0.04 to 0.52)	>0.05	0.812
Normalised overground walking speed, c m.s-1.m-1	38	47.16 (13.44 to 100.46)	< 0.01	0.185	14.16 (-0.28 to 39.66)	> 0.05	0.817
Normalised treadmill walking speed, c m.s-1.m-1	37	-31.48 (-133.81 to 35.57)	>0.05	0.061	18.13 (-4.51 to 49.08)	> 0.05	0.752
Stance time, %	38	-0.84 (-1.80 to -0.33)	< 0.01	0.156	-0.29 (-0.54 to -0.05)	< 0.05	0.819

<sup>a</sup> adjusted for age, sex and GMFCS level; <sup>b</sup>exact p value not provided as bootstrapping procedure was used to obtain  $\beta$  and associated confidence interval; <sup>c</sup>normalised to height. CI: confidence interval. Bold text indicates p < 0.05.

Olladj	usicu	and adjusted associations betwee	an gant pa	amerci	is and physical activity		
	n	Unadjusted B (95% CI)	<i>p</i> value	R <sup>2</sup>	Adjusted <sup>a</sup> β (95% CI)	p value	R <sup>2</sup>
Dependent variable: Step-coun	t, step	ps.day <sup>-1</sup>					
Hip joint excursion, deg	35	19.05 (-62.14 to 100.23)	0.636	0.007	35.20 (-51.26 to 121.67)	0.412	0.266
Knee joint excursion, deg	35	-9.81 (-83.16 to 63.53)	0.787	0.002	-25.12 (-97.23 to 46.99)	0.482	0.262
Ankle joint excursion, deg	35	-13.45 (-113.88 to 86.97)	0.787	0.002	13.98 (-86.43 to 114.39)	0.778	0.251
Normalised overground walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	35	6186.31 (536.37 to 11836.24)	0.033	0.131	4216.62 (-1933.31 to 10366.55)	0.171	0.297
Normalised treadmill walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	34	-1252.14 (-9106.30 to 6602.01)	0.747	0.003	-1738.47 (-12267.74 to 8790.80)	0.738	0.159
Stance time, %	35	-96.87 (-209.98 to 16.23)	0.091	0.084	-64.41 (-181.70 to 52.87)	0.271	0.280
Dependent variable: Moderate-	to-vi	gorous physical activity, min.day	y <sup>-1</sup>				
Hip joint excursion, deg	35	-0.09 (-0.96 to 0.78)	0.837	0.001	-0.04 (-0.87 to 0.79)	0.922	0.410
Knee joint excursion, deg	35	0.02 (-0.77 to 0.80)	0.968	0.000	-0.22 (-0.91 to 0.46)	0.513	0.418
Ankle joint excursion, deg	35	-0.27 (-1.34 to 0.80)	0.612	0.008	0.06 (-0.89 to 1.01)	0.899	0.410
Normalised overground walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	35	71.52 (11.81 to 131.22)	0.020	0.153	46.86 (-10.76 to 104.48)	0.107	0.416
Normalised treadmill walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	34	-34.36 (-116.25 to 47.53)	0.399	0.022	-43.51 (-142.40 to 55.37)	0.375	0.331
Stance time, %	35	-0.90 (-2.12 to 0.32)	0.144	0.064	-0.55 (-1.67 to 0.57)	0.325	0,429
Dependent variable: Light phys	ical a	activity, min.day <sup>-1</sup>					
Hip joint excursion, deg	35	-0.02 (-2.01 to 1.97)	0.984	0.000	-1.02 (-2.94 to 0.91)	0.290	0.391
Knee joint excursion, deg	35	0.07 (-1.73 to 1.87)	0.940	0.000	-0.76 (-2.37 to 0.85)	0.344	0.386
Ankle joint excursion, deg	35	0.25 (-2.21 to 2.71)	0.839	0.001	0.66 (-1.59 to 2.90)	0.554	0.374
Normalised overground walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	35	80.11 (-65.44 to 225.66)	0.271	0.037	44.23 (-97.58 to 186.04)	0.529	0.375
Normalised treadmill walking speed, <sup>b</sup> m.s <sup>-1</sup> .m <sup>-1</sup>	34	50.02 (-153.87 to 253.92)	0.621	0.008	-62.64 (-296.77 to 171.50)	0.588	0.386
Stance time, %	35	1.99 (-0.81 to 4.80)	0.158	0.060	2.49 (-0.04 to 5.01)	0.053	0.445

Table 4 Unadjusted and adjusted associations between gait parameters and physical activity

<sup>a</sup> adjusted for age, sex and GMFCS level; <sup>b</sup>normalised to height. CI: confidence interval. Bold text indicates p < 0.05.

		Associations	between gai	t paramete	ers, gross motor function and	physical acti	ivity	
Dependent va	ariable	Model 1: Step-count, step	s.day <sup><math>-1</math></sup> ( $n =$	34)	Model 2: MVPA, m	$n^{-1}$ (n =	=34)	Mc
Independent	variables	β (95% CI)	p value	$\mathbb{R}^2$	β (95% CI)	p value	$\mathbb{R}^2$	β
Hip joint exc	ursion, deg	132.82 (2.56 to 263.08)	0.046	0.325	0.62 (-0.78 to 2.02)	0.373	0.289	0.58 (
Knee joint ex	cursion, deg	-90.29 (-194.74 to 14.17)	0.087		-0.30 (-1.42 to 0.82)	0.587		0.47 (
Ankle joint e	xcursion, deg	-23.37 (-121.40 to 74.67)	0.628		-0.37 (-1.43 to 0.68)	0.477		-0.25
Normalised c	verground walking	962.69 (-5769.03 to 7694.40)	0.771		21.74 (-50.69 to 94.18)	0.543		76.11 (-)
speed," m.s	-m-1-2							
Stance time,	0/0	-42.49 (-164.86 to 79.88)	0.482		-0.12 (-1.44 to 1.19)	01 00 10		2.84 (
D SCI	ore	156.35 (-171.61 to 484.31)	0.336		1.64 (-1.89 to 5.17)	00 "< M 1C		7.06 (-
	ore	37.66 (-74.38 to 149.69)	0.496		0.49 (-0.72 to 1.70)	". <l` 1C</l` 		-1.16
enc	e interval; LPA: light p	physical activity; MVPA: moderat	e-to-vigorou	s activity.	_			

odel 3: LPA, min.day<sup>-1</sup> (n = 34)

Table 5

#### 4.1 Study limitations

Limitations of this study include the relatively small sample and lack of applicability of findings to adolescents with moderate motor impairment, given that only two participants were in GMFCS level III. A strength of the study is that we adjusted for age, sex and GMFCS level, which confounded a number of associations between gait parameters, function and PA as evidenced by differences between unadjusted and adjusted f3 coefficients. These confounders were not controlled for in previous studies. We calculated separate scores for GMFM-66 dimension D and E, rather than calculating a score for the full GMFM- 66, as we hypothesised that associations between gait parameters, physical activity and function may differ for function relating to standing (i.e., GMFM dimension D) and function relating to walking running and jumping (i.e., GMFM dimension E). We also believed these specific associations are of interest to clinicians. However, it should be noted that these scores on the individual dimensions of the GMFM-66 may not be reflective of a person's full GMFM-66 score. While abbreviated approaches for estimating the full GMFM-66 score are accurate at a single time point, they are less accurate at estimating change in the full GMFM-66 over time [32].

#### 5. Conclusion

This study found percentage time spent in stance is negatively associated with function assessed in a con-trolled environment, specifically activities relating to walking, running and jumping. Gait parameters and function were not associated with step-count or time in PA after adjusting for age, sex and GMFCS level. The findings provide an insight into the complexity of the relationship between gait quality or ability at the impairment level, function as measured in a controlled environment, and the performance of habitual PA, which is essential for health among children with CP.

#### Acknowledgments

Action Medical Research and the Chartered Society of Physiotherapy Charitable Trust have jointly funded this project, and it is supported by a generous grant from The Henry Smith Charity (GN2340).

#### **Conflict of interest**

The authors have no conflict of interest to report.

#### References

[I] Rosenbaum P, Paneth N, Leviton A, et al. A report: the definition and classification of cerebral palsy April 2006. Dev Med Child Neural Suppl. 2007;109:8-14.

[2] Verschuren O, Smorenburg ARP, Luiking Y, et al. Deter- minants of muscle preservation in individuals with cerebral palsy across the lifespan: a narrative review of the litera- ture. J Cachexia Sarcopenia Muscle. 2018;9:453-64. doi: 10.1002/jcsm.12287.

[3] Ryan JM, Allen E, Gormley J, et al. The risk, burden, and management of non-conunuicable diseases among peo- ple with cerebral palsy: a scoping review. Dev Med Child Neural. 2018;60:753-64 . doi: 10. l II I/dmcn.13737.

[4] World Health Organization. Towards a common language for functioning, disability and health: !CF. Geneva: 2002.

[5] Vargus-Adams JN, Martin LK. Domains of importance for parents, medical professionals and youth with cerebral palsy considering treatment outcomes. Child Care Health Dev. 2011;37:276-81. doi: 10.1 | J.J.1365-2214.2010.0l121.x.

[6] Ryan JM, Cassidy EE, Noorduyn SG, et al. Exercise inter- ventions for cerebral palsy. Cochrane Database Syst Rev 2017;6:CD011660. doi: 10.1002/14651858.CD011660.

pub2.

[7] Amirmudin NA, Lavelle G, Theologis T, et al. Multilevel Surgery for Children With Cerebral Palsy: A Metaanalysis. Pediatrics. 2019;143:e20183390. doi: 10.1542/peds.2018-

3390.

[8] Holsbeeke L, Ketelaar M, Schoemaker MM, et al. Capacity, capability, and performance: different constructs or three of a kind? Arch Phys Med Rehabil. 2009;90:849-55. doi: 10.1016/j.apnu.2008.11.015.

[9] Department of Health and Social Care. UK Chief Medical Officers' physical activity guidelines. 2019. Available at: https://assets.publishing.service.gov.uk/government/ uploads/system/uploads/attachment,data/file/832868/uk- chiefmedical-officers-physical-activity-guidelines.pdf [accessed 23 June 2021]

(10] Robinson LW, Clement N, Fullarton M, et al. The relation- ship betweei1 the Edinburgh Visual Gait Score, the Gait Pro- file Score and GMFCS levels I-III. Gait Posture. 2015;4 I: 741-3. doi: 10.1016/j.gaitpost.2015.01.022

(11] Molloy M, McDowell BC, Kerr C, et al:Further evidence of validity of the Gait Deviation Index. Gait Posture. 201 0; 31:479-82. doi: 10.1016/j.gaitpost.2010.01.025.

(12] Damiano DL, Abel MF. Relation of gait analysis to gross motor function in cerebral palsy. Dev Med Child Neural. 1996;38:389-96.

[13] Guinet AL, Desailly E. Is physical activity of children with cerebral palsy cor'elated with clinical gait analysis or physical examination parameters? Comput Methods Biomech Biomed Engin. 2017;20:99-100. doi: 10.1080/10255842. 2017.1382880.

(14] Wilson NC, Signal N, Naude Y, et al. Gait Deviation Index Correlates With Daily Step Activity in Children With Cerebral Palsy. Arch Phys Med Rehabil. 2015;96:1924-7. doi: 10.1016/j.apmr.2015.05.024

[15] Ryan JM, Theis N, Kilbride C, et al. Strength Training for Adolescents \Vith cerebral palsy (STAR): study protocol of a randomised controlled\_trial to determine the feasibility, acceptability and efficacy of resistance training for adoles- cents with cerebral palsy. BMJ Open. 2016;6:e012839. doi: I0.1136/bmjopen-2016-012839.

[16] Kiernan D, O'Sullivan R. The influence of crouch gait on sagittal trunk position and lower lumbar spinal loading in children with cerebral palsy. Gait Posture. 2019;67:65-70.

[17] Gross R, Leboeuf F, Hardouin JB, Perrouin-Verbe B, Brochard S, Remy-Neris O. Does muscle coactivation influ- ence joint excursions during gait in children with and without hemiplegic cerebral palsy? Relationship between muscle coactivation and joint kinematics. Clin Biomech. 2015;30(10):1088-93.

[18] Dingwell JB, Marin LC. Kinematic variability and local dynamic stability of upper body motions when walking at differentspeeds. J Biomech. 2006;39:444-52.

[19] Galna B, Lord S, Rochester L. Is gait variability reliable in older adults and Parkinson's disease? Towards an optimal testing protocol. Gait Posture.2013;37:580-s'. doi: 10.1016/j.gaitpost.2012.09.025.

[20] Davis R, Ounpuu S, Tyburski D, et al. A gait analysis data collection and reduction technique. Hum Mov Sci. 1991; I 0: 575-87.

(21] Noorkoiv M, 'Lavelle G, Theis N, et al. Predictors of Walking Efficiency in Children With Cerebral Palsy: Lower-Body Joint Angles, Moments, and Power. Phys Ther. 2019; 99:711-20. doi: 10.1093/ptj/pzz041.

[22] Nordmark E, Hagglund G, Jarnlo GB. Reliability of the gross motor function measure in cerebral palsy. Scand J Rehabi'! Med. 1997;29:25-8.

[23] Russell DJ, Rosenbaum PL, Cadman DT, et al. The gross motor function measure: a means to evaluate the effects of physical therapy. Dev Med Child Neural. 1989;3 I:341-52.

(24] Mitchell LE, Ziviani J, Boyd RN. Variability in measuring physical activity in children with cerebral palsy. Med Sci Sports Exer. 2015;47:194-200. doi: IO. I 249/MSS.00000 000000374.

[25] Choi L, Liu Z, Matthews CE, et al. Validation of accelerom- eter wear and nonwear time classification algorithm. Med Sci Sports Exer. 2011 ;43:357-64. doi: I 0.1249/MSS.0b013 e3181ed6la3.

[26] Trost SG, Fragala-Pinkham M, Lennon N, et al. Decision Trees for Detection of Activity Intensity in Youth with Cerebral Palsy. Med Sci Sports Exer. 2016;48:958-66. doi:.

IO. I 249/MSS.0000000000842.

[27] Bland JM, Altman DG. Statistics Notes: Bootstrap resam- pling methods. BMJ. 2015;350:112622.

[28] Carpenter J, Bithell J. Bootsu-ap confidence intervals: when, wliich, what? A practical guide for medical statisticians. Stat Med. 2000; 19(9):1141-64.

(29] Cleary SL, Taylor NF, Dodd KJ, et al. Barriers to and facili- tators of physical activity for children with cerebral palsy in special education. Dev Med Child Neural. 2019;61:1408-

15. doi: 10. IIII/dmcn.14263.

(30) Wright A, Roberts R, Bowman G, el al. BalTiers and facil- itators lo physical activity participation for children with physical disability: comparing and contrasting the views of children, young people, and their clinicians. Disabil Rehabil. 2019;41:1499-507. doi: I 0.1080/09638288.2018. 1432702.

(31) Schwartz MH, Rozumalski A. The Gait Deviation Index: a new comprehensive index of gait pathology. Gait Posture. 2008;28:351-7. doi: 10. l 016/j.gaitpost.2008.05.001.

(32] Avery LM, Russell DJ, Rosenbaum PL. Criterion validity of the GMFM-66 item set and the GMFM-66 basal and ceiling approaches for estimating GMFM-66 scores. Dev Med Child Neural. 2013;55(6):534-8.