



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document and is licensed under All Rights Reserved license:

Chambers, Frank M ORCID: 0000-0002-0998-2093, Brain, Sally A, Mauquoy, Dmitri, McCarroll, Julia and Daley, Tim (2014) The 'Little Ice Age' in the Southern Hemisphere in the context of the last 3000 years: Peat-based proxy-climate data from Tierra del Fuego. *Holocene*, 24 (12). pp. 1649-1656. doi:10.1177/0959683614551232

Official URL: <https://doi.org/10.1177/0959683614551232>

DOI: <http://dx.doi.org/10.1177/0959683614551232>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/1146>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

This is a pre-print (final draft pre-refereeing) version of the following published document:

**Chambers, Frank M and Brain, Sally A and Mauquoy, Dmitri and McCarroll, Julia and Daley, Tim (2015).
The 'Little Ice Age' in the Southern Hemisphere in
the context of the last 3000 years: Peat-based proxy-
climate data from Tierra del Fuego. The Holocene.**

Published in The Holocene, and available online at:

<http://hol.sagepub.com/content/early/2014/10/01/0959683614551232.abstract>

We recommend you cite the published (post-print) version.

The URL for the published version is <http://dx.doi.org/10/01/0959683614551232>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

The 'Little Ice Age' in the Southern Hemisphere in the context of the last 3000 years: peat-based proxy-climate data from Tierra del Fuego

Journal:	<i>The Holocene</i>
Manuscript ID:	HOL-14-0152.R1
Manuscript Type:	Fast-Track Report
Date Submitted by the Author:	28-Jul-2014
Complete List of Authors:	Chambers, Frank; University of Gloucestershire, Centre for Environmental Change and Quaternary Research, School of Natural and Social Sciences Brain, Sally; University of Gloucestershire, Centre for Environmental Change and Quaternary Research, School of Natural and Social Sciences Mauquoy, Dmitri; University of Aberdeen, School of Geosciences McCarroll, Julia; University of Gloucestershire, Centre for Environmental Change and Quaternary Research, School of Natural and Social Sciences Daley, Tim; University of Plymouth, Institute for Sustainability Solutions Research
Keywords:	Little Ice Age, proxy-climate data, Tierra del Fuego, solar forcing, peat humification, Sphagnum, Southern Hemisphere, climate change, late Holocene
Abstract:	The so-called 'Little Ice Age' (LIA) of c. the 15th to 19th centuries AD is well-attested from much of Europe and from some other parts of the Northern Hemisphere. It has been attributed to solar forcing, associated with reduced solar activity, notably during the Spörer, Maunder and Dalton solar minima, although other causes have also been proposed and feature strongly in recent papers. Detection of the LIA in some proxy-climate records from the Southern Hemisphere is less clear, leading to suggestions that the LIA was perhaps not a global phenomenon. Resolving this issue requires more data from the Southern Hemisphere. We present proxy-climate data (plant macrofossils; peat humification) covering the past three millennia from an ombrotrophic mire (peat bog) in Tierra del Fuego, southern South America, but focus our discussion on the period traditionally associated with the LIA. During parts of this time the mire surface was apparently relatively dry compared with much of its 3000-year record. It was reported earlier that a particularly dry episode in the mire coincided with the 2800 cal. BP 'solar' event (since identified as a Grand Solar Minimum), which was attributed to solar-driven changes in atmospheric circulation, and more specifically to a shift in position of the Westerlies. Parts of the LIA record show a similar shift to dryness, and we invoke a similar cause. The shifts to and from dry episodes are abrupt. These new data support the concept of a global LIA, and for at least the intense dry episodes might reinforce the claim for solar forcing of parts of the LIA climate.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



SCHOLARONE™
Manuscripts

For Peer Review

The 'Little Ice Age' in the Southern Hemisphere in the context of the last 3000 years: peat-based proxy-climate data from Tierra del Fuego

Frank M. Chambers¹, Sally A. Brain¹, Dmitri Mauquoy², Julia McCarroll¹ and Tim Daley³

¹Centre for Environmental Change and Quaternary Research, School of Natural and Social Sciences, University of Gloucestershire, Francis Close Hall, Swindon Rd, Cheltenham GL50 4AZ, United Kingdom

²School of Geosciences, University of Aberdeen, Elphinstone Road, Aberdeen AB24 3UF, United Kingdom

³Institute for Sustainability Solutions Research, A522, Portland Square, Plymouth University, Drake Circus, Plymouth PL4 8AA, United Kingdom

Abstract

The so-called 'Little Ice Age' (LIA) of c. the 15th to 19th centuries AD is well-attested from much of Europe and from some other parts of the Northern Hemisphere. It has been attributed to solar forcing, associated with reduced solar activity, notably during the Spörer, Maunder and Dalton solar minima, although other causes have also been proposed and feature strongly in recent papers. Detection of the LIA in some proxy-climate records from the Southern Hemisphere is less clear, leading to suggestions that the LIA was perhaps not a global phenomenon. Resolving this issue requires more data from the Southern Hemisphere. We present proxy-climate data (plant macrofossils; peat humification) covering the past three millennia from an ombrotrophic mire (peat bog) in Tierra del Fuego, southern South America, but focus our discussion on the period traditionally associated with the LIA. During parts of this time the mire surface was apparently relatively dry compared with much of its 3000-year record. It was reported earlier that a particularly dry episode in the mire coincided with the 2800 cal. BP 'solar' event (since identified as a Grand Solar Minimum), which was attributed to solar-driven changes in atmospheric circulation, and more specifically to a shift in position of the Westerlies. Parts of the LIA record show a similar shift to dryness, and we invoke a similar cause. The shifts to and from dry episodes are abrupt. These new data support the concept of a global LIA, and for at least the intense dry episodes might reinforce the claim for solar forcing of parts of the LIA climate.

Keywords: Little Ice Age; proxy-climate data; Tierra del Fuego; solar forcing; peat humification; *Sphagnum*

Introduction

While the timing of its onset varies both geographically and between archives, the so-called 'Little Ice Age' (LIA) from the 15th to 19th centuries AD has long been characterised in Europe as a period of advancing glaciers (Le Roi Ladurie, 1971), relatively cold winters (Grove, 1988) and, at times, failing harvests (Lamb, 1995). Often characterised as having intensely cold winters in Northwest Europe under anticyclonic conditions, the late autumn/winter weather in the LIA was much more varied than this and included a particularly severe storm event in England in late November AD 1703, classically reported by Defoe (1704).

With glaciers in Southern Norway reaching their maximum Holocene extent variously from the early 18th to the late 19th Century AD (Nesje, 2009), the LIA has long been regarded in Europe as the most severe prolonged downturn in climate warmth since the Late-glacial. However, it was not as deep as

1
2
3 the centennial-scale downturn of the so-called 8.2 ka event of the early Holocene, which is now
4 regarded as the most severe climate shift in the post-glacial (Kobashi et al., 2007; Daley et al., 2011).
5 A pronounced putative global climate shift c. 2800 cal. BP (van Geel et al., 1996; Chambers et al.,
6 2007) and a decade of relative coolness in the AD 540s, recorded especially in Europe but also
7 elsewhere (Baillie 2001), are other candidates for the second-most severe downturn since the Late-
8 glacial. In those instances, the inferred climate changes are abrupt, but short-lived. By contrast, the
9 LIA is of much longer duration than any other inferred reduction in climate warmth during the
10 Holocene, although the climate within the LIA was variable and was not uniformly colder, either
11 temporally or geographically.
12

13 While proxy-climate data from individual sites show that the period from c. AD 1420 to c. AD 1850
14 was not cold continuously, it can be viewed as a multi-centennial period in which temperatures in the
15 Northern Hemisphere were on average some 0.8°C lower than the 1961–90 reference period (Bradley
16 and Jones, 1993; Mann et al., 1998; Mann, 2002), although some archival datasets imply far greater
17 temperature reductions in specific locations (e.g. in northern Norway: Lauritzen and Lundberg, 1999).
18 More recent examination of tree-ring data over the last millennium in northern Scandinavia by Esper
19 et al. (2012) suggests some reappraisal of the magnitude of temperature changes is required.
20

21 In Europe and North America, a wide range of proxy-climate methods has been used to infer the
22 climate over the LIA. Detection of the LIA elsewhere has been patchy. Mann (2002) reviewed some of
23 this evidence, which was largely terrestrially based, and noted rather subdued cooler overall
24 temperatures in eastern China, more pronounced cooling for a time in the tropical Andes and
25 evidence of glacial advances in New Zealand, but that there were large regional variations (Jones and
26 Mann, 2004). Indeed, subsequently, Schaeffer et al. (2009) reported glacier fluctuations in New
27 Zealand that are not fully in accord with those in the Northern Hemisphere.
28

29 Whether the LIA can be regarded as a 'global' event has been disputed (compare, for example, views
30 expressed in Jones et al., 2009; Rhodes et al., 2012). Part of the reason for caution is the paucity of
31 LIA proxy-climate records from southern latitudes, which contrasts with the far greater concentration
32 of proxy-climate studies of the LIA in the Northern Hemisphere. This is attributable in part to the large
33 expanse of ocean compared with land in the Southern Hemisphere (SH), which restricts the
34 availability of suitable SH terrestrial sites to provide land-based records. However, new data from the
35 Antarctic Peninsula suggest the period of lowest ice melt and coolest temperatures in the last
36 millennium was c. AD 1410 and 1460, "when mean temperature was 1.6°C lower than that of 1981–
37 2000" (Abram et al., 2013, p. 404). Ocean-margin ice-core records point to significant LIA climate
38 changes in the SH, averaging 1.6 ± 1.4 °C cooler than during the last 150 years (Rhodes et al., 2012);
39 these authors emphasise that other SH proxy-climate records are required to enrich the LIA database.
40
41

42 The depth, geographical extent and duration of the LIA has a bearing on expectations of future
43 climate, because both the trajectory of recent climate changes and computer projections for future
44 climate are made against a background of the proxy-climate of the past millennium (Mann et al.,
45 1998; Mann, 2007) or past two millennia (e.g. Mann et al., 1999; 2008; IPCC, 2013), in which both the
46 Medieval Climate Anomaly and LIA feature—more prominently in some reconstructions than in others
47 (Mann et al., 2009; cf. Esper et al., 2012). The importance of being able to reconstruct climates of
48 past recent millennia was stressed by McCarroll (2010, p. 1661), who stated: "The wider the range of
49 climate parameters that can be reconstructed, and the wider the range of locations, the more
50 constraint will be provided".
51

52 Here, in this Research Report, we present proxy-climate data from a peat bog in Tierra del Fuego,
53 southern South America, for the last 3000 years. The methods used to derive a proxy-climate record
54 from this southerly site are identical to those that have been used widely in northern and central
55 Europe and in parts of North America (Chambers et al., 2012) and for which a set of field and
56 laboratory protocols has been established (De Vleeschouwer et al., 2010a). We focus our discussion
57 on the period of time commonly associated with the LIA, and draw comparisons within the 3000-yr
58
59
60

1
2
3 record, including a previously reported, particularly intense but short-lived climate shift in that record,
4 dated to c. 2800 cal. BP (Chambers et al., 2007), which corresponded in time with one of opposite
5 direction, reported from Europe (van Geel and Renssen, 1998). Our proxy-climate record is based
6 largely on bog surface wetness (*sensu* Barber et al., 1994), rather than temperature. Although there
7 may be a temperature component in the data, this is unlikely to be a major component at such
8 southerly cool latitudes; indeed, on the basis of changes within the period of meteorological records it
9 seems more likely that precipitation changes in response to variation in the Westerlies would have the
10 strongest influence on bog surface wetness (Daley et al., 2012).

11
12 Our data can be seen in the context of claims made recently by Moreno et al. (2014, p.1) based on
13 stratigraphic data from a lake in southwest Patagonia that there is “in-phase inter-hemispheric
14 coupling of palaeoclimate over the last 3000 years through atmospheric teleconnections”.

15 16 **Study Site and Field Methods**

17
18 The field location for analyses reported here is a *Sphagnum*-dominated raised mire, located c. 6 km
19 north-east of Ushuaia, in the broad Andorra valley (Fig. 1), Southern Argentina. Details of the mire
20 vegetation are provided in Mauquoy et al. (2004), in which results of pollen analysis on a separate
21 core of peat were reported. The two-metre core of peat (labelled AND-2) for the present study was
22 extracted later by DM using a series of 10×10×50 cm stainless steel boxes, hammered into a cleaned
23 peat face exposed during the creation of ditches for peatland harvesting (De Vleeschouwer et al.,
24 2010b), and sub-sampled into 1 cm slices, which were bagged and transported back to the UK, for
25 laboratory analysis.
26

27 **Laboratory Methods**

28
29 Two standard methods were used for producing a proxy-climate record from peats: plant macrofossil
30 analysis (Mauquoy et al., 2010), conducted by DM in Uppsala, and determination of peat humification
31 (Blackford and Chambers, 1993), the latter method modified (as in Chambers et al., 2011) and
32 conducted by SB at Cheltenham. Radiocarbon dates for the core were requested by FMC from the
33 Natural Environment Research Council, UK, in two submissions: a rangefinder series that focused on
34 major shifts in the peat humification data, followed by groups of samples that amplified the original
35 rangefinder data (see Table 1), to facilitate possible wiggle-matching (van Geel and Mook, 1989;
36 Piotrowska et al., 2011). With one exception (an *Ericales* macrofossil, close to the base of the core),
37 all dates were carried out on *Sphagnum* leaf macrofossils, separated assiduously from the peat by
38 SB, following the methods used in the Northern Hemisphere ACCROTELM project (Chambers, 2006).
39 An attempt to date the upper part of the core by ²¹⁰Pb was unsuccessful, probably owing to low radon
40 source in this region.
41

42
43 Initial deposition-rate curves were produced using a precursor of the ‘Bacon’ method of Blaauw and
44 Christen (2011); limitations of this method for extrapolation when the basal age is unknown were
45 exposed by initial lack of dating control beyond the base of the core, and so further samples were
46 dated close to the base, to anchor the basal age by wiggle-matching. Dated samples in this basal-age
47 wiggle match fitted the calibration curve very closely (Fig. 2; see also Fig. 3 in Chambers et al., 2007),
48 and so permitted precise and accurate dating of the base of the sequence. The closely-spaced sets of
49 radiocarbon dated samples at intervals along the core have ensured precision and accuracy in dating
50 the proxy-climate records, except in the uppermost layers, for which the sampling date provides a
51 topmost dating-anchor point.
52

53 **Results**

54
55 Proxy-climate data for the past 3000 years are presented in a composite diagram (Fig. 3), showing
56 peat humification (expressed inversely as percentage light transmission) and the proportions of
57 *Sphagnum* leaves and *Empetrum*/*Ericaceae* roots in plant macrofossils recorded from the mire.
58
59
60

1
2
3 These data show that for most of the past 3000 years the principal peat former has been *Sphagnum*,
4 normally with over 70% and typically 75–95% of plant macrofossils being *Sphagnum* remains.
5 Overwhelmingly, the dominant species has been *S. magellanicum*, with some 80 to 100% of the
6 *Sphagnum* leaves attributable to this species. Owing to this dominance, it was not necessary to apply
7 k-value corrections to the humification data (cf. Hughes et al., 2012). **While original above-ground**
8 **macrofossils (e.g. *Sphagnum* leaves) and below-surface remains (e.g. *Empetrum*/Ericaceae roots) in**
9 **the same horizon may not be exactly contemporaneous, the latter are relatively shallow-rooted in**
10 **ombrotrophic peats.**

11
12 There are notable exceptions to the *Sphagnum* dominance: (i) a short-lived but pronounced episode
13 c. 2800 cal. BP, in which *Sphagnum* leaves are below 30% of plant macrofossils, while
14 *Empetrum*/Ericaceae macrofossils were more abundant at >50%; (ii) a short-lived episode c. 750 cal.
15 BP (c. cal. AD 1200) in which total *Sphagnum* remains dip below 70%; and (iii) a longer period
16 towards the top the core, estimated between 275 cal. BP and 200 cal. BP (i.e., cal. AD 1675 to cal.
17 AD 1750) in which *Sphagnum* remains are below 50%, within which they twice dip below 10%, and
18 over which *Empetrum*/Ericaceae remains are generally higher than for most of the time period
19 covered by the core. In each of these low points in the *Sphagnum* curve, the light transmission data
20 also dip.
21

22
23 Peat humification data from peat bogs often show an overall trend from higher humification values
24 (i.e., lower light transmission) at the base to lower humification (or higher light transmission) towards
25 the surface. Part of the reason for this is undecayed plant material in the upper peat layer (acrotelm),
26 but the overall trend relates to continued slow decomposition over time in the catotelm below (Clymo
27 1984). Removal of that overall trend (Fig. 4) **by linear regression** reveals the low points in the light
28 transmission curve, to be c. 2800 cal. BP, 1735–1710 cal BP, c. 950 cal. BP, c. 490 cal. BP and from
29 250–180 cal. BP. These are equivalent to c. 850 cal. BC, c. cal. AD 215–240, c. cal. AD 1000, c. cal.
30 AD 1460 and c. cal. AD 1700–1770.
31

32 The ages cited above are based on the new calibration curve (Fig. 2) using the Bacon method of
33 Blaauw and Christen (2011), with the estimated ages then rounded to the nearest 5 years.
34

35 Interpretation and Discussion

36 *Plant Macrofossils*

37
38 The overall plant macrofossil record for the past 3000 years, shown in Fig. 3, indicates a mire
39 dominated by *Sphagnum magellanicum* for most this period, but with some short interludes in which
40 the mire has increased proportions of *Empetrum*/Ericaceae when *Sphagnum* is less abundant. This
41 general dominance of *S. magellanicum* for millennia is reminiscent of northwest European bogs in
42 which *S. imbricatum* (syn. *S. austinii*) has overwhelming dominance as the major peat former.
43 Interestingly, in parts of Northwest Europe, notably in Britain, *S. imbricatum* becomes extinct at many
44 sites during the last millennium, possibly related to airborne eutrophication (McClymont et al., 2008);
45 at many sites it is replaced by *S. magellanicum*, apparently occupying the same ecological niche,
46 leading to its record being interpreted as having the same climate-indicator value in data analysis,
47 with the data from the two species amalgamated (Barber et al., 1994). As *S. imbricatum* is a Northern
48 Hemisphere species, it is not native to Patagonia. No major moss species change took place in this
49 core, and so no species amalgamation has been necessary for interpreting the *Sphagnum* record.
50
51

52 The longest period in which the proportion of *Sphagnum* is lower, while *Empetrum*/Ericaceae remains
53 are more abundant, though not as prominent as c. 2800 cal. BP, is that between c. cal. 275 and c. cal.
54 200 cal. BP (i.e., c. cal. AD 1675 to 1750). This period can therefore be regarded as a significant
55 deviation from an overwhelming dominance by *Sphagnum magellanicum* during most of the past
56 three millennia.
57
58
59
60

Peat Humification

There is generally close agreement in direction, if not in magnitude, between the plant macrofossil record of a generally wet site (indicated by high percentages of *Sphagnum* remains) interrupted by short, dry episodes (shown by lower *Sphagnum* values and increased representation of *Empetrum*/Ericaceae remains), and the light transmission data, which indicate a mainly wet bog surface (shown by values >40%) interposed by drier episodes (when values dip below 40%). The most extreme dry episode is short-lived, at c. 2800 cal. BP (reported by Chambers et al., 2007). A short dry episode is suggested subsequently at c. cal. AD 1460, while a longer dry episode is indicated c. cal. AD 1700–1770.

Combining the proxy-climate records

When the two proxy-climate records are combined, there is evidence mainly from a pronounced low in the light transmission curve for a dry episode at c. cal. AD 1460, while a combination of plant macrofossil and light transmission values suggests a particularly dry episode from c. cal. AD 1675 to c. cal. AD 1770. These data compare with those from Rhodes et al. (2012, p. 1223), who inferred from a coastal ice-core record in the Ross Sea, Antarctica, that there were “three 12–30 yr intervals of cooler temperatures” within the LIA, centred on AD 1690, 1770 and 1840. The time-period c. cal. AD 1675–1770 encompasses two of those three intervals, and noticeably has a somewhat less dry period in between. The overall time-period corresponds with the most severe climatic episodes within the LIA in terms of cold climate in Northwest Europe, matched only by the early 15th Century (Lamb, 1995).

Relationship to Solar Activity?

Short-term climate changes in the Holocene have been attributed to volcanism (Robock, 2000), changes in solar activity (Beer and van Geel, 2008) or other extra-terrestrial causes (Baillie 2007), whereas the Little Ice Age has in the past been attributed to reduced solar activity (Grove, 1988), notably around the Maunder (Eddy, 1976) and Dalton minima. However, other factors, including volcanism (Robock, 2000) and ocean circulation, are claimed to play a part (Mann 2002), while its onset has recently been claimed to be caused by four, large, sulphur-rich explosive eruptions in a 50-year period, and its continuance to be sustained by sea ice/ocean feedback “during a hemispheric insolation minimum” (Miller et al. 2012, p. 1), rather than requiring a reduction in solar activity. In contrast, Hunt (2006, p. 677) used only internal forcing in the CSIRO mark 2 coupled global climate model and concluded that both the ‘Medieval Warm Period’ (i.e. the MCA) and LIA could not be explained by natural (i.e., internal) processes, and that “external forcing must be involved”. This conclusion was earlier exemplified by Mauquoy et al. (2002; 2008), who used proxy-climate techniques on mires in Denmark and northern England, and identified solar-forcing signals during the Little Ice Age

It is notable that the dry episode we infer from the Valle de Andorra mire at c. cal. AD 1460 is between a putative start of the LIA at AD 1420, according to Lamb (1995), and the lowest point in the solar-activity Spörer Minimum (at AD 1490), while the dry episode of c. cal. AD 1675–1770 is very close to the Maunder Minimum of AD 1645 to AD 1715. Indeed, the period c. cal. AD 1675–1770 presents as the driest in this mire record since the inferred solar-driven ‘global’ climate change c. 2800 cal. BP (Chambers et al., 2007; see also van Geel et al., 1996).

Suck-in and Smear

The deposition-rate curve from this site (Fig.2) contains some of the best examples of wiggle-matched dates from peat sequences, but this tight chronology only serves to emphasise the lesser chronological control in recent centuries. Baillie (1991) cautioned archaeologists about the dangers of both ‘suck-in’—in which unrelated events, apparently closely dated in time, might be interpreted falsely as presenting evidence for a single event—and ‘smear’, in which the dating of a single event

1
2
3 at different sites may cover a wide dating range, owing to inaccuracy and/or imprecision in dating
4 techniques. Interpretations made in the present study are subject to that cautionary advice. For
5 example, although the dry period from c. AD 1675–1770 is close both in time and duration to that of
6 the Maunder Minimum (AD 1645–1715), limitations of the method preclude more precise dating of the
7 mire data, because the dating of the peat that accumulated during recent centuries is compromised
8 by the acknowledged ‘fuzziness’ of the calibration (see Fig. 2) and the failure of ^{210}Pb dating at this
9 site. Indeed, it is unfortunate that some other methods to help date the top of the core, which can be
10 used in Europe (van der Plicht et al., 2013), could not be applied here. Nevertheless, the closeness of
11 fit of the driest episodes in the mire with cold episodes noted elsewhere and with solar minima, is
12 striking, even allowing for imprecision in calibrated ages (and therefore possible inaccuracy in dating)
13 in this part of the core.
14

15 *Relationship to previous work*

16
17 In previous work at the Valle de Andorra bog, using climate proxies on a separate peat core,
18 Mauquoy et al. (2004) did not find clear evidence for the LIA. However, this was under a prevailing
19 mindset that expected the mire to respond in the same way as those in Northwest Europe—to show
20 evidence of cooler, wetter conditions—and these were not evident. In this more recent detailed work,
21 the opposite tendency is shown for the LIA. Mauquoy et al. (2004) identified two notably wet periods:
22 from c. AD 1030–1100 and from c. AD 1800 to c. AD 1930. It is interesting to note that the former is
23 close to the peak warmth of the Medieval Climate Anomaly (a time-period associated in Europe with
24 warmer drier conditions and a relatively ice-free North Atlantic: Lamb, 1995), while the latter
25 encompasses the global temperature recovery from the Little Ice Age and the warming trend in the
26 early 20th Century (cf. Folland et al., 2002). Taken at face value, these data might imply that when
27 conditions in Northwest Europe, and arguably also globally, become noticeably warmer, conditions in
28 this South American mire become wetter. This is the counterpoint to the drier-than-normal episodes
29 recorded in the present study during the LIA and from this same core at 2800 cal. BP. Even so, the
30 same cautionary remarks concerning ‘suck-in’ must apply here, and this *post hoc* interpretation did
31 not feature in the 2004 paper.
32
33

34 Large-scale controls on the Patagonian climate have been explored recently using results from a 30-
35 yr numerical simulation: Garreaud et al. (2013, p. 215) conclude that variation in zonal winds account
36 for much of the rainfall variability in Patagonia “at synoptic and interannual timescales” and note
37 somewhat opposite responses to the west of the Andes (recent drying trend) compared with the
38 southern tip (modest increase). In general a West–East contrast was noted, with a positive/negative
39 correlation between the 850-hPa zonal wind speed and monthly precipitation for sites located to the
40 west/east of the Andes ridge. It is difficult to relate their findings to our interpretations above, because
41 their data covered only a limited time period; it also would be necessary to decide into which of two
42 regions the mire falls. Geographically it is East for direct precipitation; but in terms of original moisture
43 source for the river in the valley, then arguably it could be West, because it originates there from
44 glaciers and snowpacks. Hypothetically, during times of stronger Westerlies, and so with a more
45 pronounced föhn effect, there could be more snow melt, causing a higher river discharge, which in
46 turn could impede efflux from the mire; so, local water tables rise and appear to give a ‘West’
47 response. However, the overall scale of change suggested in the 3000-yr Andorra bog record is
48 greater than encompassed in that 30-yr simulation, while the abrupt episodes of LIA dryness are
49 pervasive for at least a few decades. This implies a more significant mechanism for parts of the LIA
50 than minor variations in strength of the Westerlies, such as for example a major shift in their
51 geographical position (cf. Moreno et al., 2009).
52
53

54 Varma et al. (2011) suggest that during times of reduced solar forcing (as for example, during parts of
55 the LIA), the Westerlies moved north; whereas, during times of higher solar activity (for example,
56 Medieval Climate Anomaly; late 20th Century), the westerly belt was more southerly. This contrasts
57 with an earlier view of Moy et al. (2008), who suggested a poleward shift of the Westerlies during the
58
59
60

1
2
3 LIA, based on data from a lake site in Chile. Also to the west of the Andes, from Lake Puyehue in
4 Chile, Bertrand et al. (2006) report a humid period from AD 1490–1700, which they interpret as
5 coinciding with the LIA; whereas, our record, well to the southeast of the Andes, suggests dryness in
6 the latter part of that period. Note that the dating does not coincide exactly, because after AD 1770,
7 Bertrand et al. (2006) report (in the west) a drying trend; our record shows that a putative
8 countervailing wetter period (to the east) did not start until after c. cal. AD 1770. Our data are,
9 however, seemingly in accord with Neukom et al. (2010) whose reconstructions show dry summers
10 for the 17th century in southern South America, compared with those for the mid-late 20th century,
11 which are noticeably wetter. They have only sparse data points for southernmost South America,
12 whereas our reconstructions extend the evidence for (late) 17th Century dryness, further south into
13 Tierra del Fuego. Daley et al. (2012) noted that Ushuaia (see Fig. 1) has in recent decades been
14 getting wetter over the instrumental period in response to intensification and southward movement of
15 the Westerlies. In that context, it may be hypothesized that a LIA drying of the surface of the Andorra
16 bog is not inconsistent with the Westerlies moving northward, which itself would be consistent with the
17 notion of reduced Hadley cell circulation that would be expected with reduced solar activity.

19 *Limitations to the proxy-climate measures*

21 We used two proxy-climate measures, rather than a single method, on this core. However, the
22 measures are not completely independent, because there may be a species effect in humification
23 data (Chambers et al., 1997; Yeloff and Mauquoy, 2006). The peat-humification technique has also
24 been criticised for not fully capturing decay processes (Hansson et al., 2013). In a limited comparative
25 study of methods to assess peat decay at a single site in Europe by Biester et al. (2014), techniques
26 other than the colorimetric technique could be considered to have performed better; several of these
27 other methods have been used in three pristine mires elsewhere in Patagonia (Broder et al., 2012).
28 Nevertheless, the datasets reported here indicate that the greatest inferred climatic deviations fit well
29 with climate excursions elsewhere, even though it is assumed that the principal change in the mire
30 relates to bog surface wetness, rather than to temperature. Further work might be directed to
31 investigating deuterium excess, as suggested in Daley et al. (2012), which could help to show
32 whether shifting Westerlies were indeed the mechanism for inferred changes in bog surface wetness
33 at this site.

36 **Conclusions**

38 Proxy-climate data from a mire in Tierra del Fuego, Southern Argentina, using the techniques of plant
39 macrofossil analysis and determination of peat humification, show that the main climate changes
40 recorded during the past 3000 years were (i) at 2800 cal. BP and (ii) during parts of the time period
41 conventionally associated with the Little Ice Age. These changes manifest as vegetation changes and
42 changes in peat humification, both indicative of changes in bog surface wetness; they present as
43 significantly drier-than-normal interludes. Although there may well be a temperature component in the
44 records, no temperature-change equivalent can at present be calculated from these data.

46 The 2800 cal. BP shift is sharp, short-lived, and coincides with a claimed abrupt 'global' climate
47 change attributed to a temporary decrease in solar activity during a 'Grand Solar Minimum'. That drier
48 episode was interpreted by Chambers et al (2007) as indicating a shift in the position of the moisture-
49 bearing Westerlies. Subsequent short-lived climate shifts, of lesser magnitude, seemingly occurred at
50 c. cal. AD 215–240 and c. cal. AD 1000. Later, a more pronounced short-lived shift occurred c. cal.
51 AD 1460, close to the putative claimed start of the LIA, while a twin-peak episode of more comparable
52 magnitude to that of c. 2800 cal. BP, lasted from c. cal. AD 1675–1770, coinciding with the most
53 severe interludes of the Little Ice Age in Europe. A similar mechanism to the pronounced shift at c.
54 2800 cal. BP might be invoked to explain these shifts to dryness in the last millennium: namely a
55 change in the position of the Westerlies.

1
2
3 While the proximal cause of major changes in bog surface wetness may be shifts in the southerly jet
4 stream, the data from this site suggest (i) that the LIA is recorded in Southern Argentina; (ii) at its
5 peak intensities it is comparable in magnitude to the 2800 cal. BP climate perturbation; (iii) the data
6 support Moreno et al's (2014) claim of "inter-hemispheric [temporal] symmetry"; (iv) these data are not
7 inconsistent with the long-standing interpretation of the more severe multi-decadal episodes within the
8 LIA being associated with decreased solar activity; and (v) that it may be hypothesized that decreased
9 solar activity caused an equatorward shift of the Westerlies, resulting in a drier mire surface at the
10 Andorra bog in Tierra del Fuego c. 2800 cal. BP (which while temporally coincident with a major
11 change to wetness recorded in mires in northwest continental Europe is nevertheless an opposite
12 response of the bog-surface here) and again during the more intense phases of the LIA; this
13 hypothesis remains to be tested at other sites.
14

15 Acknowledgements

16
17 Radiocarbon dating was provided by the Natural Environment Research Council, UK, under Allocation
18 number 1012.1002.
19

20 References

21
22 Abram NJ, Mulvaney R, Wolff EW et al. (2013) Acceleration of snow melt in an Antarctic Peninsula ice
23 core during the twentieth century. *Nature Geoscience* 6: 404–411.

24
25 Baillie MGL (1991) Suck in and smear—two related chronological problems for the 90s. *Journal of*
26 *Theoretical Archaeology* 2: 12–16.

27
28 Baillie, MGL (2001) The AD 540 event. *Current Archaeology* 15: 266–269.

29
30 Baillie MGL (2007) The case for significant numbers of extraterrestrial impacts through the late
31 Holocene. *Journal of Quaternary Science* 22: 101–109.

32
33 Barber KE, Chambers FM, Maddy D et al. (1994) A sensitive high-resolution record of late-Holocene
34 climatic change from a raised bog in northern England. *The Holocene* 4: 200–207.

35
36 Beer J and van Geel B (2008) Holocene climate change and the evidence for solar and other forcings.
37 In: Battarbee RW and Binney HA (eds) *Natural Climate Variability and Global Warming: A Holocene*
38 *Perspective*. Chichester: Wiley–Blackwell, pp. 138–162.

39
40 Bertrand S, Boes X, Castiaux J et al. (2006) Temporal evolution of sediment supply in Lago Puyehue
41 (Southern Chile) during the last 600 yr and its climatic significance. *Quaternary Research* 64: 163–
42 175.

43
44 Biester H, Knorr K-H, Schellekens J et al. (2014) Comparison of different methods to determine the
45 degree of decomposition in peat bogs. *Biogeosciences* (in press)

46
47 Blaauw M and Christen JA (2011) Flexible palaeoclimate age-depth models using an autoregressive
48 gamma process. *Bayesian Analysis* 6: 457–474.

49
50 Blackford JJ and Chambers FM (1993) Determining the degree of peat decomposition for peat-based
51 palaeoclimatic studies. *International Peat Journal* 5: 7–24.

52
53 Bradley RS and Jones PD (1993) 'Little Ice Age' summer temperature variations: their nature and
54 relevance to recent global warming trends. *The Holocene* 3: 367–376.

55
56 Broder T, Blodau C, Biester, H et al. (2012) Peat decomposition records in three pristine ombrotrophic
57 bogs in southern Patagonia. *Biogeosciences* 9: 1479–1491.
58
59
60

1
2
3 Chambers FM (2006) *ACCROTELM: Abrupt Climate Changes Recorded Over The European Land*
4 *Mass*. Final Report to European Commission, Contract no. EVK2-CT-2002-00166. 117 pp.

5
6 Chambers FM, Barber KE, Maddy D et al. (1997) A 5500-year proxy-climate and vegetational record
7 from blanket mire at Talla Moss, Peebleshire, Scotland. *The Holocene* 7: 391–399.

8
9 Chambers FM, Mauquoy D, Brain SA et al. (2007) Globally synchronous climate change 2800 years
10 ago: Proxy data from peat in South America. *Earth and Planetary Science Letters* 253: 439–444.

11
12 Chambers FM, Booth RK, De Vleeschouwer F et al. (2012) Development and refinement of proxy-
13 climate indicators from peats. *Quaternary International* 268: 21–33.

14
15 Chambers FM, Bielman DW and Yu ZC (2011) Methods for determining peat humification and for
16 quantifying peat bulk density, organic matter and carbon content for palaeostudies of climate and
17 peatland carbon dynamics. *Mires and Peat* 7: Article 07, 1–10.

18
19 Clymo RS (1984) The limits to peat bog growth. *Philosophical Transactions of the Royal Society of*
20 *London, Series B* 303: 605–654.

21
22 Daley, T.J., Thomas, E.R., Holmes, J.A., Street-Perrott, F.A., Chapman, M.R., Tindall, J.C., Valdes,
23 P.J., Loader, N.J., Marshall, J.D., Wolff, E.W., Hopley, P.J., Atkinson, T., Barber, K.E., Fisher, E.H.,
24 Robertson, I., Hughes, P.D.M., Roberts, C.N., (2011) The 8200 yr BP cold event in stable isotope
25 records from the North Atlantic region, *Global and Planetary Change*, 79, pp. 288-302

26
27 Daley TJ, Mauquoy D, Chambers, FM et al. (2012) Investigating late Holocene variations in
28 hydroclimate and the stable isotope composition of precipitation using southern South American
29 peatlands: an hypothesis. *Climate of the Past* 8: 1457–1471.

30
31 Defoe D (1704) *The Storm*. London: John Nutt.

32
33 De Vleeschouwer F, Hughes PDM, Nichols JE et al. (eds) (2010a) Special volume: a review of
34 protocols in peat palaeoenvironmental studies. *Mires and Peat* 7: Article 00, 1.

35
36 De Vleeschouwer F, Chambers FM and Swindles GT (2010b) Coring and sub-sampling of peatlands
37 for palaeoenvironmental research. *Mires and Peat* 7: Article 01, 1–10.

38
39 Eddy JA (1976) The Maunder Minimum. *Science* 192: 1189–1192.

40
41 Esper J, Frank DC, Timonen M et al. (2012) Orbital forcing of tree-ring data. *Nature Climate Change*
42 2: 862–866.

43
44 Folland CK, Rayner NA, Brown SJ et al. (2002) Global temperature change and its uncertainties since
45 1861. *Geophysical Research Letters* 28: 2621–2624.

46
47 Garreaud R, Lopez P, Minvielle M et al. (2013) Large-scale control on the Patagonian climate. *Journal*
48 *of Climate* 26: 215–230.

49
50 Grove JM (1988) *The Little Ice Age*. London: Methuen.

51
52 Hansson, SV, Rydberg, J, Kylander M et al. (2013) Evaluating paleoproxies for peat decomposition
53 and their relationship to peat geochemistry. *The Holocene* 23: 1666–1671.

54
55 Hughes PDM, Mallon G, Essex HJ et al. (2012) The use of k-values to examine plant 'species signals'
56 in a peat humification record from Newfoundland. *Quaternary International* 268: 156–165.

57
58 Hunt BG (2006) The Medieval Warm Period, Little Ice Age and simulated climatic variability. *Climate*
59 *Dynamics* 27: 677–694.
60

1
2
3 IPCC (2013) *Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the*
4 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: CUP.

5
6 Jones P D and Mann ME (2004) Climate over past millennia, *Reviews of Geophysics* 42: RG2002, 1–
7 42. doi:10.1029/2003RG000143.

8
9 Jones PD, Briffa KR, Osborn TJ et al. (2009) High-resolution palaeoclimatology of the last millennium:
10 a review of current status and future prospects. *The Holocene* 19: 3–49.

11
12 Kobashi T, Severinghaus JP, Brook EJ et al. (2007) Precise timing and characterization of abrupt
13 climate change 8200 years ago from air trapped in polar ice. *Quaternary Science Reviews* 26: 1212–
14 1222.

15
16 Lamb HH (1995) *Climate, History and the Modern World, 2nd ed*. London: Routledge.

17
18 Lauritzen S-E. and Lundberg J (1999) Calibration of speleothem delta function: an absolute
19 temperature record for the Holocene in northern Norway. *The Holocene* 9: 659–669.

20
21 Le Roy Ladurie E (1971) *Histoire du climat depuis l'an mil*. Paris: Flammarion, 1967. Translated by
22 Barbara Bray as *Times of Feast, Times of Famine: A History of Climate since the year 1000*. Revised
23 and updated. Garden City: Doubleday.

24
25 Mann ME (2002) Little Ice Age. In: MacCracken, MC and Perry JS (eds) *Encyclopedia of Global*
26 *Environmental Change*. Chichester: Wiley.

27
28 Mann ME (2007) Climate over the past two millennia. *Annual Review of Earth and Planetary Sciences*
29 35: 111–136.

30
31 Mann ME, Bradley RS and Hughes MK (1998) Global-scale temperature patterns and climate forcing
32 over the past six centuries. *Nature* 392: 779–787.

33
34 Mann ME, Bradley RS and Hughes MK (1999) Northern hemisphere temperatures during the past
35 millennium: inferences, uncertainties, and limitations. *Geophysical Research Letters* 26: 759–762.

36
37 Mann ME, Zhang ZH, Hughes MK, Bradley RS et al. (2008) Proxy-based reconstructions of
38 hemispheric and global surface temperature variations over the past two millennia. *Proceedings of the*
39 *National Academy of Sciences of the United States of America* 105: 13252–13257.

40
41 Mann ME, Zhang ZH, Rutherford S. et al. (2009) Global signatures and dynamical origins of the Little
42 Ice Age and Medieval Climate Anomaly. *Science* 326: 1256–1260.

43
44 Mauquoy D, Van Geel B, Blaauw M et al. (2002) Evidence from North-West European bogs shows
45 'Little Ice Age' climatic changes driven by variations in solar activity. *The Holocene* 12: 1–6.

46
47 Mauquoy D, Blaauw M, Van Geel B et al. (2004) Late Holocene climatic changes in Tierra del Fuego
48 based on multi-proxy analyses of peat deposits. *Quaternary Research* 61: 148–158.

49
50 Mauquoy D, Yeloff D, Van Geel B et al. (2008). Two decadal resolved records from north-west
51 European peat bogs show rapid climate changes associated with solar variability during the mid-late
52 Holocene. *Journal of Quaternary Science* 23: 745–763.

53
54 Mauquoy D, Hughes PDM and van Geel B (2010): A protocol for plant macrofossil analysis of peat
55 deposits. *Mires and Peat* 7: Art. 6, 1–5.

56
57 McCarroll, D (2010) Future climate change and the British Quaternary research community.
58 *Quaternary Science Reviews* 29: 1661–1672.

1
2
3 McClymont EL, Mauquoy D, Yeloff D et al. (2008) The disappearance of *Sphagnum imbricatum* from
4 Butterburn Flow, UK. *The Holocene* 18: 991–1002.

5
6 Miller GH, Geirsdóttir Á, Zhong Y et al. (2012) Abrupt onset of the Little Ice Age triggered by
7 volcanism and sustained by sea-ice/ocean feedbacks, *Geophysical Research Letters* 39: L02708,
8 doi:[10.1029/2011GL050168](https://doi.org/10.1029/2011GL050168).

9
10 Moreno PI, François JP, Villa-Martínez et al. (2009) Millennial-scale variability in Southern
11 Hemisphere westerly wind activity over the last 5000 years in SW Patagonia. *Quaternary Science*
12 *Reviews* 28: 25-38.

13
14 Moreno PI, Vilanova I, Villa-Martínez R et al. (2014) Southern Annular Mode-like changes in
15 southwestern Patagonia at centennial timescales over the last three millennia. *Nature*
16 *Communications* 5: 4375 doi:10.1038/ncomms5375

17
18 Moy CM, Dunbar RB, Moreno PI et al. (2008) Isotopic evidence for hydrologic change related to the
19 westerlies in SW Patagonia, Chile, during the last millennium. *Quaternary Science Reviews* 27: 1335–
20 1349.

21
22 Nesje A (2009) Latest Pleistocene and Holocene glacier fluctuations in Scandinavia. *Quaternary*
23 *Science Reviews* 28: 2119–2136.

24
25 Neukom R, Luterbacher J, Villalber R et al. (2010) Multi-centennial summer and winter precipitation
26 variability in southern South America. *Geophysical Research Letters* 37: L14708.
27 doi:10.1029/2010GL043680

28
29 Piotrowska N, Blaauw M, Mauquoy D, et al. (2011) Constructing deposition chronologies for peat
30 deposits using radiocarbon dating. *Mires and Peat* 7: Article 10, 1–14.

31
32 Rhodes RH, Bertler NAN, Baker JA et al. (2012) Little Ice Age climate and oceanic conditions of the
33 Ross Sea, Antarctica from a coastal ice core record. *Climate of the Past* 8: 1223–1238.
34 doi:10.5194/cp-8-1223-2012

35
36 Robock A (2000) Volcanic eruptions and climate. *Reviews of Geophysics* 38: 191–219.

37
38 Schaefer JM, Denton GH, Kaplan M et al. 2009. High-Frequency Holocene Glacier Fluctuations in
39 New Zealand Differ from the Northern Signature. *Science* 324: 622–625.

40
41 van der Plicht J, Yeloff D, van der Linden M et al. (2013) Dating recent peat accumulation in
42 European ombrotrophic bogs. *Radiocarbon* 55: 1763–1778.

43
44 van Geel B, Mook WG (1989) High-resolution 14C dating of organic deposits using natural
45 atmospheric 14C variations. *Radiocarbon* 31: 151–155.

46
47 van Geel B and Renssen H (1998) Abrupt climate change around 2,650 BP in North-West Europe:
48 evidence for climatic teleconnections and a tentative explanation. In: Issar AS. and Brown N (eds)
49 *Water, Environment and Society in Times of Climatic Change*. Dordrecht: Kluwer, pp. 21–41.

50
51 van Geel B, Buurman J. and Waterbolk HT (1996) Archaeological and palaeoecological indications for
52 an abrupt climate change in The Netherlands and evidence for climatological teleconnections around
53 2650 BP. *Journal of Quaternary Science* 11: 451–460.

54
55 Varma V, Prange M, Lamy F et al. (2011) Solar-forced shifts of the Southern Hemisphere Westerlies
56 during the late Holocene. *Climate of the Past* 7: 339–347.

57
58 Yeloff D and Mauquoy D (2006) The influence of vegetation composition on peat humification:
59 implications for palaeoclimatic studies. *Boreas* 35: 662–673.
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Figure captions

Fig. 1 Map to show location of Valle de Andorra mire and Ushuaia, Tierra del Fuego.

Fig. 2. Radiocarbon-calibrated deposition-rate curve for the AND2 core, produced using the Bacon method of Blaauw and Christen (2011), showing groups of samples that assist with 'wiggle-matching' (van Geel and Mook, 1989), resulting in a tight chronology for most of the past 3000 years. Grey-scale plot indicates relative precision of dating; calibration based on SHCal13.

Fig. 3 Percentages of identifiable *Sphagnum* [showing proportion of *S. magellanicum*] leaves and stems, and percentage light transmission through alkali extract of the AND2 core peat [low values indicate dryness; high values indicate wetness] in contiguous 1 cm samples for the past three millennia, plotted against *Empetrum*/*Ericaceae* roots [indicating relative dryness]. Age estimates derive from SHCal04, to be consistent with data published in Chambers et al. (2007); there are minor differences from SHCal13 used for Fig. 2.

Fig. 4 Detrended peat humification data for the AND2 core, plotted using absorption data, for the past 3000 years (age estimates as per Fig. 3). Note the reverse vertical axis used when plotting these data, so as to create the same visual impression as in Figure 3: deep troughs indicate short-lived dry episodes.

Table Caption

Table 1. Radiocarbon dates for Core AND-2.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 1

For Peer Review




Fig. 2





Fig. 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Detrended peat humification data for the AND2 core, plotted using absorption data, for the past 3000 years (age estimates as per Fig. 3). Note the reverse vertical axis used when plotting these data, so as to create the same visual impression as in Figure 3: deep troughs indicate short-lived dry episodes.
254x190mm (96 x 96 DPI)

Review

Table 1. Radiocarbon dates for Core AND-2

Sample name	Material	Depth (cm)	Date (¹⁴ C BP)
SUERC-2468	Sphagnum	37	395±24
AA-54892	Sphagnum	40	468±38
SUERC-2469	Sphagnum	43	394±24
SUERC-2470	Sphagnum	46	448±20
SUERC-2471	Sphagnum	49	402±18
SUERC-2472	Sphagnum	52	565±24
SUERC-2473	Sphagnum	70	952±22
AA-54893	Sphagnum	73	1101±39
SUERC-2476	Sphagnum	76	961±22
SUERC-2477	Sphagnum	79	1050±25
SUERC-2478	Sphagnum	82	1029±20
SUERC-3033	Sphagnum	85	1131±20
SUERC-2481	Sphagnum	118	1721±25
SUERC-2482	Sphagnum	121	1722±22
SUERC-2483	Sphagnum	124	1747±25
SUERC-2486	Sphagnum	127	1813±22
SUERC-2487	Sphagnum	130	1796±25
AA-54894	Sphagnum	133	1810±38
SUERC-3877	Sphagnum	153	2196±25
SUERC-3878	Sphagnum	157	2236±25
SUERC-3881	Sphagnum	161	2256±19
SUERC-3882	Sphagnum	165	2255±25
SUERC-3883	Sphagnum	169	2238±25
SUERC-3884	Sphagnum	173	2241±21
SUERC-3886	Sphagnum	180	2398±23
SUERC-3887	Sphagnum	184	2418±25
SUERC-3888	Sphagnum	188	2465±21
SUERC-3891	Sphagnum	192	2508±26
SUERC-3892	Sphagnum	196	2577±23
SUERC-5570	Sphagnum	198	2717±30
SUERC-6312	Ericales wood	199	2704±22
SUERC-5573	Sphagnum	199	2805±30
AA-54895	Sphagnum	200	2866±41