



This is a peer-reviewed, final published version of the following document, © 2023 Hanson, Fryer, Higgins and Stoner. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms. and is licensed under Creative Commons: Attribution 4.0 license:

**Hanson, Erik D, Fryer, Simon M ORCID logoORCID:
<https://orcid.org/0000-0003-0376-0104>, Higgins, Simon and
Stoner, Lee (2023) Editorial: Sedentary behavior and health
outcomes. Origins, mechanisms, and policy. *Frontiers in
Physiology*, 14. Art 1197926. doi:10.3389/fphys.2023.1197926**

Official URL: <http://doi.org/10.3389/fphys.2023.1197926>

DOI: <http://dx.doi.org/10.3389/fphys.2023.1197926>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/12677>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.



OPEN ACCESS

EDITED AND REVIEWED BY
Giuseppe D'Antona,
University of Pavia, Italy

*CORRESPONDENCE
Erik D. Hanson,
✉ edhanson@email.unc.edu

RECEIVED 31 March 2023
ACCEPTED 04 April 2023
PUBLISHED 13 April 2023

CITATION
Hanson ED, Fryer S, Higgins S and
Stoner L (2023), Editorial: Sedentary
behavior and health outcomes. Origins,
mechanisms, and policy.
Front. Physiol. 14:1197926.
doi: 10.3389/fphys.2023.1197926

COPYRIGHT
© 2023 Hanson, Fryer, Higgins and
Stoner. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License](#)
(CC BY). The use, distribution or
reproduction in other forums is
permitted, provided the original author(s)
and the copyright owner(s) are credited
and that the original publication in this
journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Editorial: Sedentary behavior and health outcomes. Origins, mechanisms, and policy

Erik D. Hanson^{1,2*}, Simon Fryer³, Simon Higgins¹ and
Lee Stoner^{1,4,5}

¹Department of Exercise and Sport Science, University of North Carolina, Chapel Hill, NC, United States, ²Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, United States, ³School of Natural, Social and Sport Sciences, University of Gloucestershire, Gloucester, Gloucestershire, United Kingdom, ⁴Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, United States, ⁵Center for Health Promotion and Disease Prevention, University of North Carolina, Chapel Hill, NC, United States

KEYWORDS

sedentary behavior, physical inactivity, cardiometabolic health, prolonged sitting, cardiovascular disease

Editorial on the Research Topic

Sedentary behavior and health outcomes. Origins, mechanisms, and policy

1 Introduction

Sedentary behaviors (SB), defined as low-intensity activities in a seated, reclined, or supine positions, have become a ubiquitous aspect of contemporary societies. Accordingly, international public health authorities, such as the World Health Organization, now recognize the importance of SB as it relates to health outcomes, including but not limited to: cardiometabolic diseases, some cancers, quality of life, and mental health. As highlighted by [Higgins et al.](#), SB research is in its infancy and there is much work to be done before we as a research community can inform policy beyond the vague advice of “move more and sit less”.

Population-level increases in SB coupled with decreasing levels of physical activity (PA) have negative implications on cardiovascular disease (CVD) rates, which is the leading cause of death world-wide. CVD risk is multi-factorial and is influenced by many different lifestyle factors highlighted in this Research Topic, such as poor diet ([Fryer et al.](#)), reduced PA ([Vaara et al.](#); [Bowden Davies et al.](#)), and increased sitting time ([Bowden Davies et al.](#); [Zimmer et al.](#)). Lifestyle behaviors are complex and interrelated, with individual-, social-, and environmental-level determinants, but are also modifiable with improved diets and greater activity levels working to reduce low-grade inflammation that hastens chronic disease progression ([Huston](#)). However, by examining the origins of SB, we can begin to understand how these occurrences came to be and are better positioned to develop strategies to sustainably change behavior and inform policy ([Higgins et al.](#)).

This Research Topic examined SB in the context of its evolution, biological plausibility for associations with health outcomes, interruption interventions, and implementation through to policy. It includes 6 papers ([Vaara et al.](#); [Bowden Davies et al.](#); [Higgins et al.](#); [Huston](#); [Zimmer et al.](#); [Fryer et al.](#)). Within this Research Topic, cardiometabolic health outcomes ranged widely, but were consistently investigated within the context of SB and/or PA.

2 Health outcomes

Examining SB and health outcomes was a most common approach within this Research Topic. Previously, self-reported PA and SB have been linked with cardiorespiratory fitness, but such measures lack objectivity and can lead to over/under-reporting. Here, objective tracking devices were used to improve outcome reporting accuracy, with PA and SB being positively and negatively associated with cardiorespiratory fitness, respectively (Vaara et al.). Moreover, PA and SB were also associated with lower body fitness (standing long jump, maximal isometric force) in a similar manner, extending knowledge in this area. Next, 14 days of step reduction in habitually active individuals (>10,000 steps/day) decreased endothelial function (flow mediated dilation), insulin sensitivity, and cardiorespiratory fitness while increasing total body fat (Bowden Davies et al.). However, these impairments were all reversed upon resumption of normal activity levels. Finally, poor dietary choices and prolonged SB are often co-occurring behaviors. Meals with high fat content are a potentially important factor associated with increased CVD prevalence. Supporting this idea, an acute bout of 180 min of sitting coupled with a low-fat meal revealed negligible changes in carotid-femoral pulse wave velocity, though prolonged sitting coupled with a high fat meal led to significantly increased (worsened) arterial stiffness (Fryer et al.). Importantly, heel raises performed every 5 min abrogated the adverse effects of the high fat meal. Together, transitory periods of decreased PA that consequently increased SB led to acute negative changes in markers of cardiovascular health that can be reversed with resumption of normal activity.

3 SB interruption interventions

Effective SB interruption strategies was the second major theme, although this frequently intertwined with Health Outcomes as both Bowden Davies et al. (Bowden Davies et al.) and Fryer et al. incorporated interruption strategies to improve their targeted health outcomes. In addition, the context of SB is a growing area of investigation, with the impact of different types of SB on cardiometabolic outcomes receiving recent attention. However, amateur e-sports report no changes in energy expenditure, respiratory or cardiovascular parameters during 30 min of videogame play (Zimmer et al.), suggesting that whilst short bouts of video gaming may not cause acute cardiovascular dysfunction, they also do not provide the benefits of traditional PA and likely contribute to increased SB time. Collectively, regular muscle contractions appear to be a necessary factor leading to healthy endothelial function and maintaining arterial stiffness. However, identifying ways to integrate greater movement into current lifestyles along through the identification of barriers and facilitators SB along with practical disruption doses remain to be determined.

4 Sedentary behavior evolution and context

The remaining papers provided an evolutionary perspective and a network analysis. Higgins and others (Higgins et al.) discuss how

ancestral SB and PA likely differed from the modern world, including co-occurring behaviors (e.g., eating, screen time, mental stress) that may accelerate CVD risk. The authors highlight that the scientific literature has a long-standing emphasis on achieving adequate PA levels, yet ironically many of these same active individuals also engage in high volumes of SB. Continued efforts that improve awareness of SB are essential, with development of specific guidelines that are on par with sleep (7–9 h per night) and PA (>150 min/week) being a key towards policy changes. A network analysis provides a theory for how unhealthy lifestyle factors (high SB/low PA) lead to chronic disease (Huston). These included (i) the macroenvironment, (ii) social/culture factors, (iii) lifestyle, (iv) physiological systems, (v) the microenvironment and (vi) intracellular networks. Low-grade, sustained inflammation is argued to be a principal underlying factor within the body, stemming from poor diet, along with socioeconomic and education disparities that contribute to reduced PA and increased SB which are exacerbated by global instability.

5 Conclusion

This Research Topic has helped address some of the pressing research questions related to SB in the context of its evolution, associations with health outcomes, and potential interruption strategies. While the goal of implementation of specific policies to reduce SB remains unfulfilled, studies of these nature are necessary first steps. As the world emerges from the COVID-19 pandemic, there is an opportunity to progress development of guidelines that ultimately can be used to inform policy. From a practical standpoint, effective, evidence-based tools to sustainably reduce SB are needed to assist practitioners and policymakers in the development of population specific health education plans and reduce the increasing prevalence and severity of non-communicable diseases.

Author contributions

EH: design, writing, and topic editor. SF: proof reading and topic editor. SH: proof reading and topic editor. LS: proof reading and topic editor. All authors contributed to the article and approved the submitted version.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.