



This is a peer-reviewed, final published version of the following document and is licensed under Creative Commons: Attribution-Noncommercial 4.0 license:

**Walker, Daniel, Qureshi, Adam W., Marchant, David, Ford, Ben
ORCID logoORCID: <https://orcid.org/0000-0001-7502-7505> and
Balani, Alex Bahrami (2023) Providing a clearer insight into
how sport-related concussion and physical pain impact mental
health, cognition, and quality of life. Journal of Concussion, 7.
doi:10.1177/20597002221142379**

Official URL: <http://doi.org/10.1177/20597002221142379>

DOI: <http://dx.doi.org/10.1177/20597002221142379>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/13862>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Providing a clearer insight into how sport-related concussion and physical pain impact mental health, cognition, and quality of life

Journal of Concussion
Volume 7: 1–9
© The Author(s) 2023
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20597002221142379
journals.sagepub.com/home/ccn



Daniel Walker¹ , Adam W. Qureshi², David Marchant², Ben Ford^{2,3} and Alex Bahrami Balani²

Abstract

Sport-related concussion (SRC) and physical pain are both associated with poor mental health, impaired cognition, and reduced quality of life. Despite SRC and physical pain often co-occurring, there is little research that investigates these two factors together, and therefore it is difficult to conclude which of these contributes to the negative outcomes associated with them. Therefore, the present study aimed to investigate the effect of SRC and physical pain on mental health, cognitive ability, and quality of life. Depression was measured using the Center for Epidemiological Studies Depression Scale, anxiety was assessed using the State-Trait Anxiety Inventory while the SF-12 recorded health-related quality of life. A trail making task (TMT) assessed cognitive flexibility of participants. Analysis of 83 participants (43 concussed) revealed that SRC led to reduced accuracy on TMT(A) and (B), whereas physical pain was responsible for poorer mental health and reduced quality of life. This study highlights the influence that SRC has on cognitive ability and the impact that physical pain has on mental health and quality of life. With this information, we are better placed to predict the negative consequences of SRC and physical pain and therefore tailor support accordingly.

Keywords

Depression, anxiety, cognitive flexibility, quality of life, concussion, physical pain

Date received: 12 May 2022; accepted: 12 November 2022

Introduction

Sport-related concussion (SRC) has become a widely investigated area in recent years, with many sport-related activities involving high energy events that can expose athletes to direct and indirect traumas and increase their risk of injury.^{1,2} Consequences of SRC have been identified, with poorer mental health,^{3,4} impaired cognition,^{5,6} and reduced quality of life,^{7,8} among the most prevalent outcomes. Although physical pain often accompanies SRC,^{9,10} surprisingly few studies incorporate this factor into SRC research. Therefore, the relationship between SRC and physical pain and these three broad outcomes warrants investigation. By better understanding how SRC and physical pain relate to mental health, cognitive ability, and quality of life, we are better able to support athletes that have sustained SRC, experiencing physical pain, or both.

Depression is one of the most researched mental health disorders, likely due to its high prevalence, with 3.8% of the general population suffering from depression.¹¹ However, higher rates are reported in athletes that

have^{12,13} and have not sustained SRC.^{14,15} These elevated depressive symptoms are also found in student athletes who report higher levels of physical pain.^{14,16} Given the great comorbidity between depression and anxiety, it is predictable that elevated scores of anxiety are also often found in those that have sustained SRC¹⁷ and experiencing physical pain.^{18–20} Therefore, as SRC and physical pain are likely to co-occur^{9,10} and both are found to negatively impact depression and anxiety, it is important to understand which contributes more to these outcomes.

Cognitive flexibility involves the ability to adjust to change in task demands or priorities²¹ and is therefore

¹University of Bradford, West Yorkshire, UK

²Edge Hill University, Ormskirk, Lancashire, UK

³University of Bristol, Bristol, UK

Corresponding author:

Daniel Walker, University of Bradford, Richmond Road, Bradford, West Yorkshire, UK

Email: dwalker5@bradford.ac.uk



important for athletes with evidence that cognitive flexibility levels of team athletes are higher than those of individual athletes.²² Aslan²² suggests this could be due to individual sports requiring less cognitively than team sports. Team sports tend to be contact sports such as rugby, football, and hockey whereby SRC is likely,²³ and as SRC negatively impacts cognitive flexibility,^{24,25} this threatens optimal performance²² as well as exacerbating other negative consequences of SRC. Therefore, it is necessary to protect athletes from sustaining SRC and the impact this can have on cognitive flexibility. The role of physical pain on cognitive flexibility is still unclear with many studies examining chronic pain in humans^{26,27} and animals.²⁸ Despite the literature unclear on the role that physical pain has on cognitive flexibility, there are studies linking physical pain with impairments to other areas of cognition.^{5,6} As these studies did not account for SRC, it may be that impaired cognition was misattributed to physical pain, and therefore assessing both simultaneously is warranted.

Considering there is evidence that SRC and physical pain are associated poorer mental health^{12,13,17} and cognitive impairment,^{24,25} it is unsurprising that these factors are also associated with reduced quality of life.^{29,30} It is possible that the two broad outcomes could negatively impact quality of life. For example, impaired cognition leading to individuals struggling to complete daily tasks could negatively impact quality of life. Likewise, not being able to compete at a sporting level that one once could, can lead to poorer mental health and quality of life. However, even if these examples did explain reduced quality of life in athletes, it is still unclear whether this is due to SRC or physical pain, with evidence that both negatively impact quality of life,^{31–34} and therefore this warrants further exploration.

Although there is evidence that SRC negatively impacts mental health such as depression^{12,13} and anxiety,¹⁷ areas of cognition such as cognitive flexibility,^{24,25} and quality of life,^{29,31,34} the role that physical pain has on these outcomes is still ambiguous. This is due to evidence that physical pain can also lead to poorer mental health,^{14,16} impaired cognition,^{26,27} and reduced quality of life.^{32,33} Although SRC and pain are often experienced simultaneously,^{9,10} and athletes are among the most at-risk of sustaining both, the two contributing factors are rarely examined in conjunction in the literature. Therefore, the present study aims to address this gap in the literature, as learning which factor affects each outcome will allow better support for athletes that have sustained SRC, are experiencing physical pain, or both.

Method

Participants

Forty-three post-concussed participants were compared with 40 non-concussed participants. The total sample

completed questionnaires measuring depression and anxiety symptoms as well as quality of life, and a trail making task (TMT) designed to measure cognitive flexibility. Participants were recruited via the online departmental recruitment system within the university, advertisements on social media platforms LinkedIn and Twitter as well as word of mouth. Men and women over 18 years of age were welcomed to take part in the study. For those that had sustained concussion, a minimum of 28 days must have passed before taking part in this study to avoid exacerbating post-concussion symptoms due to the surveys being displayed on a computer screen. Those that had sustained concussion < 28 days were ineligible to take part until this period had elapsed. Post-concussed participants had sustained SRC between 1 and 48 months ($M = 21.38$, $SD = 14.92$) prior to participation with incidences ranging between 1 and 12 ($M = 3.88$, $SD = 3.01$). Further information of participants is presented in Table 1.

Sample size calculations. Sample size was calculated using G*Power 3.1.9.7. A priori calculations indicated that 84 participants were required (across the two groups) to achieve adequate power ($\beta = 0.95$) to detect an effect size of $f = 0.4$ (large effect size) at $\alpha = 0.05$. Therefore, the sample in this study falls one short to achieve this level of power. Post-hoc calculations were conducted to compute the actual power achieved with the sample obtained ($\beta = 0.949$) which highlighted the sample was not underpowered. A large effect size was used in calculations due to the time difference between variables (SRC history being 28+ days ago compared with physical pain in the past week).

Measures

General Information Questionnaire (GIQ): included data on sex, age, athlete status, concussion history, and level of physical pain experienced in the past week.

Numeric Rating Scale-11 (NRS-11)³⁵: The NRS-11 measured self-reported physical pain in the past week providing a score ranging 0–10. Downie et al.³⁵ suggests that a score of ‘0’ indicates no pain, ‘1–3’ mild pain, ‘4–6’ moderate pain, and ‘7–10’ severe pain. Therefore, data from NRS-11 can be used as continuous or categorical.

Center for Epidemiological Studies Depression Scale (CESD)³⁶: The CESD is a 20-item questionnaire that measures the depressive symptoms that participants have experienced in the past week. A four-point scale was used to rank the responses; 0 to 3. Zero indicated that participants had experienced that symptom ‘rarely or none of the time’ (less than once a week), 1 ‘some or a little of the time’ (1–2 days a week), 2 ‘occasionally or a moderate amount of time’ (3–4 days a week) and 3 ‘most or all of the time’ (5–7 days a week). Four items were reverse coded due to the nature of the question. A total score ≥ 16 of 60

Table 1. Participant characteristics.

		Age				N			
		n	M	SD	p	SRC	Non-SRC	Total	p
Sex	Male	50	24.22	4.85	.604	35	15	50	.001
	Female	33	23.55	6.98		8	25	33	
Athlete Status	Athlete	50	24.10	4.87	.775	38	12	50	.001
	Non-athlete	33	23.73	6.96		5	28	33	
SRC History	Yes	43	23.51	4.74	.474	43	-	43	-
	No	40	24.43	6.71		-	40	40	
Depressed Categorisation	Yes	43	23.23	4.66	.240	24	19	43	.455
	No	40	24.73	6.72		19	21	40	

highlights an individual may be experiencing some form of depression.³⁷ A higher total score reflected more severe depressive symptoms.

*State-Trait Anxiety Inventory (STAI)*³⁸: The STAI was used to assess state and trait-anxiety in participants. The 40-item questionnaire was presented as two surveys; Y-1 and Y-2, which measured state-anxiety and trait-anxiety, respectively. The Y-1 form evaluated how participants felt at present whereas the Y-2 form recorded how participants feel generally. The four responses for both forms were ranked 1 to 4. For the Y-1 form, 1 suggested 'not at all', 2 indicated 'somewhat', 3 indicated 'moderately so' and 4 indicated 'very much so'. For the Y-2 form, 1 suggested 'almost never', 2 indicated 'sometimes', 3 indicated 'often' and 4 indicated 'almost always.' This resulted in a scoring range of 20–80 for both State-Anxiety and Trait-Anxiety with greater scores indicating higher anxiety levels.

*Health Related Quality of Life: SF-12 (SF-12)*³⁹: The SF-12 contained 12 items which had a two, three, five and six-point Likert scale depending on the nature of the question. The sum of scores was calculated between 0 and 36, with greater scores indicating better quality of life.

*TMT: The Trail Making Task (TMT)*⁴⁰: a computerised version of TMT (Pavlovía⁴¹) was used. TMT is a popular switching task that can assess cognitive flexibility and consists of two parts (A & B) in which the participant connects 25 circles as quickly as possible while maintaining accuracy.⁴² The first part of the task, TMT-A, the circles are numbered 1–25, and the person should connect the numbers in ascending order. In part B, the circles include both number (1–13) and letters (A–L). Connecting dots 1–25 provides a baseline score whereas the second part, TMT-B, acts as the switching task. Accuracy scores and reaction times were recorded for both TMT (A) and TMT (B).

Reliability statistics

Reliability statistics was calculated by conducting a Cronbach's Alpha. Analyses revealed internal consistency scores of $\alpha = .91$ for CESD, $\alpha = .95$ for STAI (state), $\alpha =$

.94 for STAI (trait) which are all considered excellent.⁴³ The SF-12 bore an internal consistency score of $\alpha = .81$ which Sharma⁴³ describes as a good score.

Procedure

Participants completed the questionnaires on the online survey platform Qualtrics⁴⁴ (Qualtrics, Provo, UT) and were fully informed about the study with their consent obtained prior to participation. This was followed by the participants completing the study questionnaires (GIQ, CESD, STAI, SF-12). After completing the last questionnaire, participants were directed to the TMT on Pavlovía.

Ethics

British Psychological Society (BPS) ethical guidelines were adhered to with data collection commencing after ethical approval was obtained from the University's Departmental Research Ethics Committee (DREC). A participant information sheet informed participants of the nature of the study and their rights as a participant including details on the withdrawal of data if they wished to do so. Tick box options were provided with the consent form with validation options set disallowing participants to continue if they did not provide consent. A debrief form was displayed at the end of the study.

Data analysis

Correlations were run between SRC history, mental health, quality of life, cognitive flexibility, and physical pain, isolated and when controlling for physical pain. A MANOVA and MANCOVA investigated the difference in cognitive flexibility, depression, anxiety, and quality of life between those that have and have not sustained SRC; one with and one without controlling the effect of physical pain. It was predicted that those that have sustained SRC would produce lower accuracy scores and higher reaction times scores on the cognitive flexibility task, elevated

depressive and anxiety symptoms and lower scores of quality of life, as well as physical pain covarying significantly.

Results

Correlations

Correlation analysis was run on multiple variables; SRC history, depression, anxiety, cognitive flexibility, quality of life and physical pain, with and without physical pain as a covariate. Results are described in Table 2.

MANOVA/MANCOVA

Results from the MANOVA (without controlling for physical pain) and the MANCOVA where physical pain was added as a covariate are displayed in Table 3. Descriptive statistics are presented below in Table 4 to aid the interpretation of the findings from Table 3.

Post-Hoc tests

Following this, the assumption of homogeneity of regression slopes was tested, as this is vital when using MANCOVA analysis. The interaction term between SRC history and physical pain had a significant effect on TMT(B) accuracy ($p = .032$) violating the assumption of homogeneity of regression slopes and therefore results for this variable should be inferred with caution. However, this assumption has been adhered to for all other variables ($p > .05$), indicating that we can accept what has been found in the main analysis for these factors.

Discussion

The present study aimed to identify the role that SRC and physical pain have in conjuncture on mental health, cognitive ability, and quality of life. Initially, our analysis revealed that SRC negatively impacted cognitive flexibility performance, with lesser accuracy recorded on both aspects of a TMT. When accounting for physical pain, we found the same results regarding SRC and cognitive flexibility, but also that physical pain was associated with depression, state-trait anxiety, and quality of life. Therefore, despite literature that links both SRC and physical pain with poor mental health,^{3,4,14,16} cognitive impairment,^{5,6} and reduced quality of life,^{7,8,32,33} a more nuanced explanation would be that SRC is responsible for cognitive impairment and physical pain contributes to poor mental health and reduced quality of life.

SRC and impaired cognitive flexibility have been linked before,^{24,25} however, the role that physical pain has on cognitive flexibility is unclear. This is because studies assessing chronic pain in humans^{26,27} and animals²⁸ are

commonplace. Many athletes and non-athletes⁴⁵ live daily with acute physical pain, however, the effect of this on cognitive flexibility, to our knowledge, has not yet been examined. This study addresses this shortfall and provides evidence that acute physical pain has no detrimental impact to cognitive flexibility, which has positive practical implications given the many negative impacts of physical pain. From a sporting perspective, Vestberg et al.⁴⁶ have shown that professional and semi-professional football players had significantly better performances in switching tasks like TMT. Similar findings have been reported in rugby,⁴⁷ basketball⁴⁸ and tennis.⁴⁹ This signifies the importance of task switching ability and cognitive flexibility for optimal sporting performance. Therefore, it is reassuring that physical pain is not associated with impairment to cognitive flexibility but concerning that SRC history is.

Although it is thought cognitive impairment subsides shortly after concussion,⁵⁰ and some studies suggest cognitive impairments resolve within a week,^{51,52} this study indicates that the effects can last up to four years following concussion. Our sample consisted of athletes that had sustained concussion between 1 and 48 months prior to their participation with a large variance as highlighted by the standard deviation. Therefore, cognitive flexibility may be impaired following a concussion that never returns to baseline (however, pre-morbid assessments were not conducted in this study). Cognitive flexibility is important for various reasons within⁵³ and outside of sport,^{21,54} and therefore this study highlights the importance of avoiding SRC.

Moreover the prevention of physical pain is paramount. Where SRC history is associated with poorer cognitive flexibility, physical pain appears to contribute to poorer mental health (depression and anxiety) and reduced quality of life. These findings support previous literature that have found physical pain to be associated with poor mental health^{14,16} and reduced quality of life.^{29,30} However, this study brings into question studies that conclude concussion is responsible for negative changes to mental health and quality of life, that do not account for physical pain.^{3,4,7,8} As concussion and physical pain often co-exist,^{9,10} it could be that previous studies have misattributed these negative consequences to concussion, where in fact physical pain may be the responsible factor. Therefore, from the present study, it appears that SRC is associated with cognitive impairment and physical pain is responsible for poor mental health and reduced quality of life.

Knowledge application

This main findings from this study are that SRC is linked with poorer cognitive flexibility, even long after the event, and acute physical pain has a negative impact on mental health and quality of life. These findings are

Table 2. Correlations between SRC history and mental health, cognitive flexibility, and quality of life with and without controlling for physical pain.

Control Variables	SRC	Depression	State-anxiety	Trait-anxiety	Quality of Life	TMT(A) Accuracy	TMT(A) RT	TMT(B) Accuracy	TMT(B) RT	TMT Accuracy Difference	TMT RT Difference	Physical Pain
None												
SRC	—	—	—	—	—	—	—	—	—	—	—	—
Depression	-.097	—	—	—	—	—	—	—	—	—	—	—
State-anxiety	-.055	.762**	—	—	—	—	—	—	—	—	—	—
Trait-anxiety	.008	.789**	.787**	—	—	—	—	—	—	—	—	—
Quality of Life	.102	-.777**	-.701**	-.750**	—	—	—	—	—	—	—	—
TMT(A)	.230*	.066	.132	.089	-.166	—	—	—	—	—	—	—
Accuracy												
TMT(A) RT	-.059	.075	-.008	.047	-.033	-.043	—	—	—	—	—	—
TMT(B)	.321**	-.122	-.111	.010	.121	.419**	.021	—	—	—	—	—
Accuracy												
TMT(B) RT	.125	.185	.090	.206	.256*	.022	.436**	-.122	—	—	—	—
TMT	-.049	-.148	-.212	-.091	.201	-.824**	.060	.170	-.100	—	—	—
Accuracy Difference												
TMT RT	-.161	-.067	-.079	-.112	.167	-.063	.706**	.118	-.329**	.142	—	—
Difference												
Physical Pain												
SRC	-.022	.301**	.277*	.236*	-.328**	-.040	-.062	-.040	.185	.019	-.210	—
Depression	—	—	—	—	—	—	—	—	—	—	—	—
State-anxiety	-.095	—	—	—	—	—	—	—	—	—	—	—
Trait-anxiety	-.051	.741**	—	—	—	—	—	—	—	—	—	—
Quality of Life	.014	.774**	.772**	—	—	—	—	—	—	—	—	—
TMT(A)	.101	-.752**	-.673**	-.733**	—	—	—	—	—	—	—	—
TMT(B)	.229*	.082	.149	.102	-.137	—	—	—	—	—	—	—
Accuracy												
TMT(A) RT	-.061	.098	.010	.064	-.057	-.046	—	—	—	—	—	—
TMT(B)	.320**	-.116	-.104	.020	.114	.418**	.019	—	—	—	—	—
Accuracy												
TMT(B) RT	.132	.138	.041	.170	-.211	.030	.456**	-.117	—	—	—	—
TMT	-.049	-.161	-.226*	-.098	.219*	-.824**	.061	.171	-.105	—	—	—
Accuracy Difference												
TMT RT	-.169	-.004	-.022	-.066	.106	-.073	.711**	.112	-.302**	.149	—	—
Difference												
Physical Pain	—	—	—	—	—	—	—	—	—	—	—	—

Note: * $-p < .05$ ** $-p < .01$.

Table 3. MANCOVA depicting the influence of SRC history on cognitive flexibility, depression, anxiety, and quality of life with pain as a covariate.

	Df	η_p^2	F	p
MANOVA				
SRC				
TMT (A) Accuracy (%)	1, 81	.053	4.524	.036*
TMT (A) Reaction Time (s)	1, 81	.004	0.286	.594
TMT (B) Accuracy (%)	1, 81	.103	9.296	.003**
TMT (B) Reaction Time (s)	1, 81	.016	1.295	.258
TMT Accuracy Difference	1, 81	.002	0.198	.658
TMT Reaction Time Difference	1, 81	.026	2.154	.146
Depression	1, 81	.009	0.774	.381
State-Anxiety	1, 81	.003	0.243	.623
Trait Anxiety	1, 81	.000	0.005	.947
Quality of Life	1, 81	.010	0.857	.357
MANCOVA				
SRC				
TMT (A) Accuracy (%)	1, 80	.053	4.441	.038*
TMT (A) Reaction Time (s)	1, 80	.004	0.297	.587
TMT (B) Accuracy (%)	1, 80	.103	9.146	.003**
TMT (B) Reaction Time (s)	1, 80	.017	1.415	.238
TMT Accuracy Difference	1, 80	.002	0.192	.662
TMT Reaction Time Difference	1, 80	.029	2.363	.128
Depression	1, 80	.009	0.730	.395
State-Anxiety	1, 80	.003	0.205	.652
Trait Anxiety	1, 80	.000	0.015	.902
Quality of Life	1, 80	.010	0.819	.368
Physical Pain				
TMT (A) Accuracy (%)	1, 80	.001	0.106	.746
TMT (A) Reaction Time (s)	1, 80	.004	0.319	.574
TMT (B) Accuracy (%)	1, 80	.001	0.098	.755
TMT (B) Reaction Time (s)	1, 80	.036	2.962	.089
TMT Accuracy Difference	1, 80	.000	0.025	.875
TMT Reaction Time Difference	1, 80	.047	3.927	.051
Depression	1, 80	.090	7.904	.006**
State-Anxiety	1, 80	.076	6.613	.012*
Trait Anxiety	1, 80	.056	4.745	.032*
Quality of Life	1, 80	.107	9.607	.003**

* $p < .05$, ** $p < .01$.

Note: significant results are highlighted in **bold**.

concerning given prior evidence that physical activity appears to promote cognitive flexibility performance.^{55,56} From this study, it seems physical activity is not enough to offset the effects that SRC has on cognitive ability. Given that age between concussion groups were not significantly different, it is unlikely that age was a confound in this sample whereby changes in cognitive flexibility could be attributed. Additionally, as both groups are young, it is alarming that concussion has had a negative long-lasting influence on cognitive ability. Based on these findings, sports coaches, especially in contact sports where concussion is prevalent,⁵⁷ should incorporate cognitive flexibility tasks within training sessions to combat cognitive flexibility changes. There is evidence that cognitive training can slow down cognitive decline,⁵⁸ and as

Table 4. Descriptive statistics of cognitive flexibility, depression, anxiety, and quality of life by SRC history.

	SRC		Non-SRC	
	M	SD	M	SD
TMT (A) Accuracy (%)	94.51	12.22	98.70	2.46
TMT (A) Reaction Time (s)	45.90	28.54	43.30	11.72
TMT (B) Accuracy (%)	95.14	6.53	98.50	2.51
TMT (B) Reaction Time (s)	49.21	13.64	53.33	19.04
TMT Accuracy Difference	0.63	11.25	-0.20	3.62
TMT Reaction Time Difference	33.12	24.49	10.03	15.96
Depressive Symptoms	18.60	11.33	16.60	9.23
State-Anxiety Symptoms	40.37	13.01	39.00	12.28
Trait Anxiety Symptoms	45.05	12.92	45.25	12.03
Quality of Life Scores	23.65	5.93	24.83	5.59
Physical Pain	2.77	2.06	2.68	2.18

Note: significant results are highlighted in **bold**.

there are benefits to efficient cognitive flexibility within⁵³ and outside of sport,^{21,54} this is highly beneficial for athletes and sports teams.

Additionally, this study links physical pain with reduced quality of life, with the reasons for improving quality of life obvious. Providing athletes with pain questionnaires to complete regularly may benefit athletes and their teams. Physical pain has been associated with depression,¹⁴ and depression has in-turn been linked with lower quality of life.⁵⁹ Therefore, pain questionnaires may indicate those at risk of reduced quality of life. There is also evidence that female athletes will experience depression at lower levels of physical pain than male athletes,¹⁶ and therefore coaches should consider this. Therefore, regular anonymous pain questionnaires would aid the identification of struggling athletes and in-turn supporting those that require it.

Limitations

Although the present study has a relatively low sample size it was not underpowered as highlighted by the sample size calculations. Therefore, the risk of type II error was low. There is, however, a risk of self-selection bias due to the voluntary nature of our recruitment. As the study was advertised as exploring the impact that SRC has on mental health, cognitive ability, and quality of life, post-concussed athletes that have noticed deterioration of these three factors may be over-represented in this sample. This could explain the high prevalence of depression in the sample regardless of concussion history, as well as impaired cognition and lower quality of life in the post-concussed sample. Additionally, data collection commenced largely during COVID-19 restrictions, and as these restrictions were found to be detrimental to mental health,⁶⁰ this could have influenced results regarding mental health and quality of life. However, this is unlikely to have any

impact regarding SRC history, as all would have been subjected to these restrictions. Although this confound was unavoidable, future research should continue to investigate the impact that SRC has on mental health and quality of life.

Another issue was the difference in timeframe between SRC history and physical pain. SRC history was recorded as anyone that had previously sustained SRC in the past four years. By contrast, physical pain scores were recorded as to the level of pain experienced in the week preceding the study. Therefore, this disparity in timeframe may be reflected in the present findings and could explain why physical pain is associated with mental health and quality of life. However, while research tends to suggest cognitive impairment lasts for around a week,^{51,52} this study suggests that these effects persist over a longer period, and future studies should investigate the mid to long-term effects of SRC on cognitive ability.

Importantly, it must be noted that participants completed questionnaires and the cognitive task just once and were then separated into concussion groups based on their reporting of their SRC history. Due to this, we cannot know the cognitive abilities of the concussed group prior to their concussion. It is possible that this group would have always performed less accurately on the cognitive flexibility task, regardless of their concussion, and that we are falsely attributing this to the concussive event. However, our approach is typical within concussion literature due to practical issues associated with pre-morbid assessments. Therefore, the results still hold value, despite this limitation.

That said, the usual approach within concussion literature is to match participants for potential confounds so that the impact of SRC is evident. However, in this study, there was a significant difference between the biological sex of participants and their status as an athlete. As there are significant differences between these variables, they could both act as confounds when assessing the negative impacts of SRC. In this study, there were more males than females that had sustained SRC, and therefore the results here are perhaps more applicable to males. Moreover, athlete status is unlikely to effect the outcomes we were interested in as SRC, by nature, is only possible in athletes. In fact, some athletes could withdraw from sport due to sustaining SRC and experiencing its' negative consequences (i.e., there were 5 participants that were non-athletes that has sustained SRC) and would therefore be classed as a non-athlete in this study. Therefore, although this limitation should be considered when inferring the results, it is perhaps better to evade this misclassification of athlete status in the analysis.

Conclusion & future directions

The present study attempted to investigate the impact of SRC and physical pain on mental health, cognitive

ability, and quality of life. Despite what is often reported,^{51,52} SRC appears to have a long-lasting negative impact on cognitive ability. The SRC group performed significantly worse on TMT (A) and (B) than the non-SRC group with and without controlling for physical pain, indicating that SRC is responsible for this reduced accuracy. It is noteworthy that this effect was twice as strong for TMT(B) than (A). Additionally, physical pain was significantly associated with poor mental health and reduced quality of life, supporting previous literature that deems this to be the case.^{29,30} The effect on quality of life was stronger than any effect on mental health measures, and just under twice as strong as the effect on trait-anxiety. Given what we know about the link between physical pain and depression,^{14,16} and depression and quality of life,⁵⁹ it is sensible to treat them similarly. Sports teams could utilise pain questionnaires to identify vulnerable athletes that may require emotional support.


Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Daniel Walker  <https://orcid.org/0000-0002-9369-6953>

References

1. Chambers R, Gabbett TJ, Cole MH, et al. The use of wearable microensors to quantify sport-specific movements. *Sports Med* 2015; 45: 1065–1081.
2. Póvoas SC, Ascensão AA, Magalhães J, et al. Physiological demands of elite team handball with special reference to playing position. *J Strength Cond Res* 2014; 28: 430–442.
3. Hoyle C. Mental Health Outcomes of Elite Athletes Following Sport-Related Concussion. <https://doi.org/10.7939/r3-ctcq-9f80>. 2020.
4. Mrazik M. Mental health outcomes of elite athletes following sport-related concussion: a prospective study. *Arch Clin Neuropsychol* 2021; 36: 661.
5. Gonzalez AC, Kim M, Keser Z, et al. Diffusion tensor imaging correlates of concussion related cognitive impairment. *Front Neurol* 2021; 12: 1–12. <https://doi.org/10.3389/fneur.2021.639179>
6. Kodali S and Fisque MAMM. Cognitive impairment induced by sports-related concussion in adolescents: a review. *Int J Child Adolesc Health* 2021; 14: 117–126.
7. Doroszkiewicz C, Gold D, Green R, et al. Anxiety, depression, and quality of life: a long-term follow-up study of patients with persisting concussion symptoms. *J Neurotrauma* 2021; 38: 493–505.

8. Walton SR, Broshek DK, Kranz S, et al. Mood, psychological, and behavioral factors of health-related quality of life throughout recovery from sport concussion. *J Head Trauma Rehabil* 2021; 36: 128–136.
9. Mollaveva T, Cassidy JD, Shapiro CM, et al. Concussion/mild traumatic brain injury-related chronic pain in males and females: a diagnostic modelling study. *Medicine (Baltimore)* 2017; 96. <https://doi.org/10.1097/MD.00000000000005917>
10. Provance AJ, Howell DR, Potter MN, Wilson PE, D'Lauro AM and Wilson JC. Presence of neck or shoulder pain following sport-related concussion negatively influences recovery. *Journal of child neurology* 2020; 35: 456–462.
11. WHO. *Depression*. 2023, February 23. <https://www.who.int/news-room/fact-sheets/detail/depression>.
12. Rosenthal JA, Foraker RE, Collins CL, et al. National high school athlete concussion rates from 2005–2006 to 2011–2012. *Am J Sports Med* 2014; 42: 1710–1715.
13. Wolanin A, Hong E, Marks D, et al. Prevalence of clinically elevated depressive symptoms in college athletes and differences by gender and sport. *Br J Sports Med* 2016; 50: 167–171.
14. Walker D and Marchant D. The impact of injury and physical pain on depressive symptoms in student athletes. *Sport Exerc Psychol Rev* 2020; 16: 60–71.
15. Yang J, Peek-Asa C, Corlette JD, et al. Prevalence of and risk factors associated with symptoms of depression in competitive collegiate student athletes. *Clin J Sport Med* 2007; 17: 481–487.
16. Walker D and McKay C. Interactive effects of sex and pain on elevated depressive symptoms in university student athletes. *Int J Sport Exerc Psychol* 2022; 1–12. <https://doi.org/10.1080/1612197X.2022.2057571>.
17. Yang J, Peek-Asa C, Covassin T, et al. Post-concussion symptoms of depression and anxiety in division I collegiate athletes. *Dev Neuropsychol* 2015; 40: 18–23.
18. Elbinoune I, Amine B, Shyen S, et al. Chronic neck pain and anxiety-depression: prevalence and associated risk factors. *Pan Afr Med J* 2016; 24. <https://doi.org/10.11604/pamj.2016.24.89.8831>
19. Gureje O. Comorbidity of pain and anxiety disorders. *Current psychiatry reports* 2008; 10: 318–322.
20. Walding MF. Pain, anxiety and powerlessness. *J Adv Nurs* 1991; 16: 388–397.
21. Diamond A. Executive functions. *Annu Rev Psychol* 2013; 64: 135–168.
22. Aslan S. Examination of cognitive flexibility levels of young individual and team sport athletes. *J Educ Train Stud* 2018; 6: 149–154.
23. Harmon KG, Clugston JR, Dec K, et al. American Medical society for sports medicine position statement on concussion in sport. *Br J Sports Med* 2019; 53: 213–225.
24. Hume PA, Theadom A, Lewis GN, et al. A comparison of cognitive function in former rugby union players compared with former non-contact-sport players and the impact of concussion history. *Sports Med* 2017; 47: 1209–1220.
25. Wilmoth K, Mau K, Guzowski N, et al. Direct comparison of multidimensional clinical assessment tools: sensitivity to concussion in student athletes. *Arch Clin Neuropsychol* 2019; 34: 737–737.
26. Hageman MG, Briet JP, Oosterhoff TC, et al. The correlation of cognitive flexibility with pain intensity and magnitude of disability in upper extremity illness. *J Hand Microsurg* 2014; 6: 59–64.
27. Moriarty O, Gorman CL, McGowan F, et al. Impaired recognition memory and cognitive flexibility in the rat L5–L6 spinal nerve ligation model of neuropathic pain. *Scand J Pain* 2016; 10: 61–73.
28. Cowen SL, Phelps CE, Navratilova E, et al. Chronic pain impairs cognitive flexibility and engages novel learning strategies in rats. *Pain* 2018; 159: 1403.
29. Gard A, Lehto N, Engström Å, et al. Quality of life of ice hockey players after retirement due to concussions. *Concussion* 2020; 5: CNC78.
30. Niv D and Kreitler S. Pain and quality of life. *Pain Pract* 2001; 1: 150–161.
31. Ponsford J, Nguyen S, Downing M, et al. Factors associated with persistent post-concussion symptoms following mild traumatic brain injury in adults. *J Rehabil Med* 2019; 51: 32–39.
32. Samadi A, Salehian R, Kiani D, et al. Effectiveness of duloxetine on severity of pain and quality of life in chronic low back pain in patients who had posterior spinal fixation. *J Orthop Trauma Rehabil* 2021; 1–6. <https://doi.org/10.1177/2210491720983333>
33. Vaegter HB, Christoffersen LO, Enggaard TP, et al. Socio demographics, pain characteristics, quality of life and treatment values before and after specialized interdisciplinary pain treatment: results from the Danish clinical pain registry (PainData). *J Pain Res* 2021; 14: 1215.
34. Voormolen DC, Polinder S, Von Steinbuechel N, et al. The association between post-concussion symptoms and health-related quality of life in patients with mild traumatic brain injury. *Injury* 2019; 50: 1068–1074.
35. Downie WW, Leatham PA, Rhind VM, et al. Studies with pain rating scales. *Ann Rheum Dis* 1978; 37: 378–381.
36. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas* 1977; 1: 385–401.
37. Husaini BA, Neff JA, Harrington JB, et al. Depression in rural communities: validating the CES–D scale. *J Community Psychol* 1980; 8: 20–27.
38. Spielberger CD, Gorsuch RL and Lushene RE. *The state trait anxiety inventory manual*. Palo Alto, Cal.: Consulting Psychologists, 1970.
39. Ware JE Jr, Kosinski M and Keller SD. A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996; 34(3): 220–233. <https://doi.org/10.1097/00005650-199603000-00003>
40. Reitan RM. The relation of the trail making test to organic brain damage. *J Consult Psychol* 1955; 19: 393.
41. Peirce JW, Gray JR, Simpson S, et al. PsychoPy2: experiments in behavior made easy. *Behav Res Methods* 2019; 51: 195–203. <https://doi.org/10.3758/s13428-018-01193-y>
42. Arnett JA and Labovitz SS. Effect of physical layout in performance of the trail making test *Psychol Assess* 1995; 7: 220–221.
43. Sharma B. A focus on reliability in developmental research through Cronbach's alpha among medical, dental and para-medical professionals. *Asian Pac J Health Sci* 2016; 3: 271–278.

44. Qualtrics. *Qualtrics*. <https://www.qualtrics.com>. 2005.
45. Ferreira AS, de Oliveira Silva D, Del Priore LB, et al. Differences in pain and function between adolescent athletes and physically active non-athletes with patellofemoral pain. *Phys Ther Sport* 2018; 33: 70–75.
46. Vestberg T, Gustafson R, Maurex L, et al. Executive functions predict the success of top soccer players. *PLoS One* 2012; 7: 1–5.
47. Faubert J. Professional athletes have extraordinary skills for rapidly learning complex and neutral dynamic visual scenes. *Sci Rep* 2013; 3: 1154. <https://doi.org/10.1038/srep01154>
48. Cortis C, Tessitore A, Lupo C, et al. Inter-limb coordination, strength, jump, and sprint performances following a youth men's basketball game. *J Strength Cond Res* 2011; 25: 135–142.
49. Shim J, Carlton LG, Chow JW, et al. The use of anticipatory visual cues by highly skilled tennis players. *J Mot Behav* 2005; 37: 164–175.
50. Lifshitz J, Witgen BM and Grady MS. Acute cognitive impairment after lateral fluid percussion brain injury recovers by 1 month: evaluation by conditioned fear response. *Behav Brain Res* 2007; 177: 347–357.
51. Bleiberg J, Cernich AN, Cameron K, et al. Duration of cognitive impairment after sports concussion. *Neurosurgery* 2004; 54: 1073–1080.
52. Delaney JS, Lacroix VJ, Leclerc S, et al. Concussions among university football and soccer players. *Clin J Sport Med* 2002; 12: 331–338.
53. Mujika I, Halson S, Burke LM, et al. An integrated, multifactorial approach to periodization for optimal performance in individual and team sports. *Int J Sports Physiol Perform* 2018; 13: 538–561.
54. Roy MH and Dugal SS. Developing trust: the importance of cognitive flexibility and co-operative contexts. *Manage Decis* 1998; 561–567. <https://doi.org/10.1108/00251749810239441>
55. Dupuy O, Billaut F, Raymond F, et al. Effect of acute intermittent exercise on cognitive flexibility: the role of exercise intensity. *J Cognit Enhancement* 2018; 2: 146–156.
56. Ludyga S, Gerber M, Mücke M, et al. The acute effects of aerobic exercise on cognitive flexibility and task-related heart rate variability in children with ADHD and healthy controls. *J Atten Disord* 2020; 24: 693–703.
57. Bakhos LL, Lockhart GR, Myers R, et al. Emergency department visits for concussion in young child athletes. *Pediatrics* 2010; 126: 550–556.
58. Hussein F, Damirchi A and Babaei P. Effect of brain training on cognitive performance in elderly women diagnosed with mild cognitive impairment. *Casp J Neurol Sci* 2016; 2: 25–31.
59. Diaz R, Miller EK, Kraus E, et al. Impact of adaptive sports participation on quality of life. *Sports Med Arthrosc* 2019; 27: 73–82.
60. Banks J, Fancourt D and Xu X. Mental health and the COVID-19 pandemic. *World happiness report 2021*, 24. 2021.