



This is a peer-reviewed, final published version of the following document, © 2025 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). and is licensed under Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0 license:

Johnson, Richard M., Pandey, Bindhi Wasini, Chand, Kesar, Davies, Ceri L ORCID logo ORCID: <https://orcid.org/0000-0003-3484-2866>, Edwards, Debra, Edwards, Esther, Jeffers, James, King, Kieran, Kuniyal, Jagdish Chandra, Mishra, Himanshu, Phillips, Victoria, Roy, Nikhil, Seviour, Jessica, Sharma, Dev Dutt, Sharma, Pushpanjali, Singh, Harkanchan and Singh, Ram Babu (2025) 'HiFlo-DAT': A flood hazard event-disaster database for the Kullu District, Himachal Pradesh, Indian Himalaya. *International Journal of Disaster Risk Reduction*, 120. art: 105336. doi:10.1016/j.ijdr.2025.105336

Official URL: <https://doi.org/10.1016/j.ijdr.2025.105336>

DOI: <http://dx.doi.org/10.1016/j.ijdr.2025.105336>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/14883>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.


PLEASE SCROLL DOWN FOR TEXT.



ELSEVIER

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

International Journal of Disaster Risk Reduction

journal homepage: www.elsevier.com/locate/ijdr

‘HiFlo-DAT’: A flood hazard event-disaster database for the Kullu District, Himachal Pradesh, Indian Himalaya

Richard M. Johnson^{a,b,*}, Bindhy Wasini Pandey^d, Kesar Chand^e, Ceri L. Davies^c, Debra Edwards^b, Esther Edwards^b, James Jeffers^b, Kieran King^b, Jagdish Chandra Kuniyal^f, Himanshu Mishra^d, Victoria Phillips^b, Nikhil Roy^d, Jessica Seviour^b, Dev Dutt Sharma^g, Pushpanjali Sharma^g, Harkanchan Singh^{h,2}, Ram Babu Singh^{d,1}

^a Institute of Science and Environment, University of Cumbria, Ambleside, LA22 9BB, UK

^b Climate, Hazards and Environmental Change Research Cluster, School of Sciences, Bath Spa University, Bath, BA2 9BN, UK

^c Geography, University of Gloucestershire, The Park, Cheltenham, GL50 2RH, UK

^d Department of Geography, and Centre for Himalayan Studies, University of Delhi, Delhi, 110007, India

^e G.B. Pant National Institute of Himalayan Environment, Himachal Regional Centre, Mohal-Kullu, 175126, Himachal Pradesh, India

^f College of Forestry, Veer Chandra Singh Garhwal University of Horticulture and Forestry, Ranichauri, 249199, Uttarakhand, India

^g Department of Geography, Himachal Pradesh University, Shimla, 171005, Himachal Pradesh, India

^h Kullu District Disaster Management Authority (DDMA), Kullu, 175101, Himachal Pradesh, India

ARTICLE INFO

Keywords:

Flood
Hazard
Disaster
Database
India
Himalaya

ABSTRACT

‘HiFlo-DAT’ (Himalayan Flood Database) contributes to the disaster risk reduction (DRR) agenda of developing methodologies for the assembly, analysis, and application of disaggregated/sub-national disaster loss data; here for mountain floods in the Kullu District, Himachal Pradesh, India. The HiFlo-DAT architecture is aligned to international best practice/local needs. It uses English-language documents, principally newspapers and government reports (1835–2020), and comprises 128 flood events, at 59 locations, over 175 years (1846–2020). This open-access database brings a substantial improvement over existing compilations. Subject to the fidelity of historical event recording, analyses highlight temporal/process patterns inclusive of flood-rich periods (1890–1900s; 1990s-present: 68 % of events), increasing flood occurrence towards the present, the prevalence of rainfall causation (55 %), and the dominance of summer monsoon flooding (June–September: 87 %). Spatially, of the 59 locations recording floods, 76 % record a single event, 24 % have two or more events, and four tributaries record 8–14 events. Key flood impact receptors were roads (55 floods), bridges (54 floods and 94 impacts) and vulnerable labourer-migrant communities (70 % fatalities and 83 % affected) notably associated with construction projects in remote/exposed locations. Key opportunities for policy and practice

* Corresponding author. Institute of Science and Environment, University of Cumbria, Ambleside, LA22 9BB, UK.

E-mail addresses: richard.johnson@cumbria.ac.uk (R.M. Johnson), bwpdsegeo@gmail.com (B.W. Pandey), kesar82@yahoo.com (K. Chand), cdavies30@glos.ac.uk (C.L. Davies), debra.edwards@hotmail.co.uk (D. Edwards), edwards.esther21@gmail.com (E. Edwards), j.jeffers@bathspa.ac.uk (J. Jeffers), kieranleeking@btinternet.com (K. King), jckuniyal@gmail.com (J.C. Kuniyal), himanshumishra136@gmail.com (H. Mishra), vp141300@gmail.com (V. Phillips), nikhilroy.geo@gmail.com (N. Roy), jessicaseviour@live.co.uk (J. Seviour), ddshpu@gmail.com (D.D. Sharma), aditi.sharma799@gmail.com (P. Sharma), harkanchan15dm2@gmail.com (H. Singh).

¹ Author deceased.

² Presently: Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ), Shimla, Himachal Pradesh, India.

<https://doi.org/10.1016/j.ijdr.2025.105336>

Received 21 December 2024; Received in revised form 20 February 2025; Accepted 22 February 2025

Available online 25 February 2025

2212-4209/© 2025 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

development include transference of the HiFlo-DAT methodology across the wider Indian Himalayan Region and trans-boundary basins; multi-disciplinary approaches to corroborate and extend documentary-based databases; improved access to public archive materials; routine integration of historical flood data into DRR/climate change adaptation management planning and infrastructure development design; and deeper multi-agency partnership to record contemporary flood impacts to provide effective data for current/future DRR.

1. Introduction

In considering flood hazard/risk in mountain catchments it is important to contextualise the global significance and diversity of mountains. Gardner [1], Walz et al. [2], Hock et al. [3], Sharma et al. [4], and Adler et al. [5] provide synthesis, where the capital of mountains includes their biodiversity, cultural heritage, and ecosystem services (e.g., energy, water, food), for global human populations. Mountains can however be remote, form disputed trans-boundary settings, may have populations with higher levels of poverty, and have hazard-prone environmental systems. Rapidly changing environment-society conditions in mountains are increasing disaster exposure, vulnerability and risk; driven by integrated stressors including climate variability, habitat degradation, conflict, globalisation, infrastructure development, tourism, urbanisation and population change. Amplifying concerns and contributing to an urgent call for action, are complex/contested future climate change trajectories in the Indian Himalayan Region. These signal increased temperatures, typically increasing precipitation but with regional diversity, likely increasing summer monsoon precipitation which generates many floods, and likely decreasing low flow and more extreme high flow events within river channels [6, 7,8,9]. We are reminded of these challenges by high-magnitude hazard events across the Indian Himalayan Region. For example, the September 2014 floods in the Kashmir Valley in north-west India [10] and further east in Uttarakhand, the Kedarnath disaster in June 2013 [11,12,13] and the Chamoli disaster in February 2021 [14,15,16]. More broadly Vaidya et al. [17], using CRED EM-DAT (global disaster data; 1980–2015), establish that Hindu Kush Himalaya nations account for 21 % of all major disaster events, in which India recorded 438 events (second to China). Furthermore, floods are a prominent hazard type both globally [18,19] and in the Himalayan region [19,20]. Taking a 5-year snapshot of CRED EM-DAT data (2016–2020) for India, shows recorded floods accounted for 44 % of hazard events ($n = 36$), 68 % of deaths ($n = 6549$), 53 % of affected persons ($n = 53.9$ million) and 57 % of damage costs.

The UNDRR ‘Sendai Framework for Disaster Risk Reduction 2015–2030’ (SFDRR) is the key policy driving DRR efforts, of which India is a signatory. Regional policy groups bring further focus; India is a member of the ‘Asia-Pacific Ministerial Conference on Disaster Risk Reduction’ (APMCDRR) who adopted the Asia-Pacific Action Plan 2021–2024 in December 2021, and 2024–2027 in October 2024 [21,22]. Building on the precursor Asia Regional Plan [23] these plans target the development of methodologies for the compilation of disaster loss data to help reduce future disaster impacts; aligning with SFDRR Priority 1 ‘*understanding disaster risk*’, inclusive of ‘*the collection, analysis, management and use of relevant data and practical information and ensure its dissemination ...*’ (clause 24a). India has operationalised this and other DRR priorities via National, State and District disaster management authorities. A small number of States have existing multi-hazard atlases of historical events (e.g., Gujarat, Himachal Pradesh, and Uttarakhand), ‘Memoranda of Loss and Damages’ of contemporary events (e.g., 2013–2024 in Himachal Pradesh), and ‘Post Disaster Needs Assessments’ (e.g., 2023 Monsoon floods/landslides in Himachal Pradesh; [24,25]). In parallel, national efforts [26] to extend understanding of flood risk include a national ‘Flood Affected Area Atlas of India’ [27], based on satellite imagery (1998–2022), but has limited information for Himachal Pradesh. Alongside this the India Meteorological Department (IMD) maintains an annual record of ‘Disastrous Weather Events’ (1967-present). This details heavy rainfall/flood events, at state and district levels, but entries are typically over generalised. More broadly, large bi-lateral initiatives like the ‘Indian Himalayas Climate Adaptation Programme’ (IHCAP) in Himachal Pradesh, not only assembled historical flood data, but also recommended ‘*the dissemination of information and training related to hazard zones, elements-at-risk and best practices*’ [28]. Thereby empowering local agencies and communities to reflect on their hazard/risk perceptions and to foster greater societal engagement for risk informed development and disaster preparedness/resilience near river channels.

Accordingly, standing observations likely persist, that Indian disaster data recording and analysis requires more coordination [29]; governance structures are not capitalising on the depth of local knowledge [30]; that data holdings remain ‘incomplete’ and ‘incomprehensive’ [31] and are not regularly maintained/updated; and there is a tendency for insufficient data disaggregation and missing data (i.e., gaps) in existing databases resulting in skewed appreciation of disaster risk and thus inefficient/ineffective DRR [11, 32]. Similarly, an independent assessment by ADPC-UNDRR [33], p20 reports: ‘*despite these initiatives and stakeholders involved, risk and disaster information in India has remained fractured across various agencies, ministries and administrative levels, with little cross-compatibility and harmonization which limits comprehensive analysis of risks in the country ...*’. Encouragingly a change to legislation, i.e., The Disaster Management (Amendment) Bill December 12, 2024, may support greater action, as this mandates the creation of national and state level disaster databases [34].

These debates withstanding, global hazard/disaster databases such as ADRC (Asian Disaster Reduction Centre, Japan), EM-DAT (Emergency Events Database: Centre for Research on the Epidemiology of Disasters, Belgium), NatCatSERVICE (Munich Re, Germany) and Reliefweb (UN Office for the Coordination of Humanitarian Affairs) are all established portals. Other inventories, such as the Dartmouth Flood Observatory: Global Active Archive of Large Flood Events (University of Colorado, USA; Brackenridge, no date) and IFNet (Infrastructure Development Institute, Japan) also operate internationally but with a focus on flood events. These all have restrictive criteria for database inclusion, typically focussing on the largest/most devastating events, thus filtering out smaller flood

events of local significance ([32,35–37]) and assume the trigger process type is clearly identifiable. Whilst these global compilations can assist macro-level policy/practice, they can be a poor fit to a country's needs. Instead, national/sub-national stakeholders require more comprehensive, localised and hazard process specific databases to effectively implement national policies for development planning and anticipative risk management ([30,32,33,35]).

Given these Indian flood disaster challenges and policy targets to enhance data quality, the objectives of 'HiFlo-DAT' (Himalayan Flood Database) are to: (1) use documentary archives to generate a new, locally focussed and open-access database of historical floods in the Kullu District (Himachal Pradesh, India), enabling a revised appraisal of past flood characteristics; (2) outline a transferable methodology for flood databases, applicable to the wider Indian Himalayan Region (>530,000 km², extending 2500 km across 13 States/Union Territories/Districts); and to (3) offer recommendations for policy-practice development.

2. Study region and existing flood databases

2.1. Kullu and Manali Tehsils, Kullu District, Himachal Pradesh

Situated in northern India, the Kullu District (Fig. 1) has a resident population of c. 440,000 [38], swelled by transient economic migrants, tourists, and pilgrims (>1 million influx in 2004; [39]) which exacerbate exposure, vulnerability and risk to hazards [40]. The succession of governance (pre-colonial, British colonial, and post-Independence) in this region is significant, as this has conditioned the language, quantity and current location of historical materials which may detail past floods. Accordingly, HiFlo-DAT uses globally dispersed English-language records, incorporating the period of British colonial administration (1846–1947: 'Kulu sub-division of Kangra District'; where 'Kulu' is the pre-independence spelling of both the area and principal settlement), post-1947 Indian independence under Punjab administration (1947–1966) and Himachal administration (1966 onwards).

HiFlo-DAT's spatial extent is the area of the Manali and Kullu Tehsil's (Fig. 1) in the northern part of the Kullu District (3561 km², c. 0.67 % of the Indian Himalayan Region; [42]). Selecting these Tehsil's yields many English-language documentary records during the colonial to post-independence periods, perhaps with greater abundance than most of the Indian Himalaya. This reflects: (1) being focal points of former British colonial administration/meteorological monitoring in Kulu and Naggar [43,44] (2) reporting interest by former media, e.g. the Civil and Military Gazette (newspaper) had a correspondent based in Kulu in the late 19th Century; (3) alignment with long-standing/more intensively settled locations, key trade/transport routes and economic activity in the District [44, 45]; and (4) a high frequency of hazard events as a consequence of local seismic, hydro-meteorological (i.e., Indian summer monsoon, winter snowfall, snowmelt) and active sediment transfer cascades ([40,46,47]). Moreover, these areas are rapidly developing,




Fig. 1. 'HiFlo-DAT' spatial location: Manali and Kullu Tehsils in the Kullu District, Himachal Pradesh, India. (DEM dataset: [41]; Kullu District: c. 31° 58' N; 77° 06' E; 5503 km²; 1089 to >6500 m AMSL).

changing the hazard-risk equation, for example hydro-electric power schemes [48], road widening and tunnelling [49,50], proposed Kullu-Manali airport runway extension [51], planned railways [52], and tourism growth [53].

2.2. Kullu District hazard-disaster profile

Accounts of hazard-disaster events, in the Kullu District incorporate research [40,46,54], government/NGO policy-practice [55–57, 58], and community-based knowledge [59,60]; commonly including earthquakes, floods, slope failures, snow avalanches, fires, crop infestations and epidemics. Existing databases of flood events for the Manali and Kullu Tehsils (Table 1), showcase the paucity of publicly available data, reporting 1–15 flood events since 1994. This may in part reflect known losses of documentary records during past hazard events. For example, the May 21, 1894 Phojal Nalla flood destroyed the record offices in Dawara and Dobhi [61]; more recently the September 4, 1995 Beas River floods washed away records in Patlikuhal. HiFlo-DAT responds, being a new open-access compilation of historical floods, empowering more effective flood risk assessments, disaster risk reduction, and development in the Kullu District [62,63].

3. 'HIFLO-DAT' method

3.1. An overview

Developing a robust database for the Kullu District, which is potentially transferable across the Indian Himalayan Region necessitates extended methodological articulation. Our approach was highly consultative, grounded in best practice and project managed, empowering a bi-lateral team to work with globally dispersed materials. Fig. 2 details the key phases: HiFlo-DAT design, data selection, data capture, data review, data entry, database analysis, and dissemination.

3.2. HiFlo-DAT design and structure

A workshop in Delhi (June 2018) comprising Indian State and District government, DRR NGOs and researchers, distilled user aspirations for the database design: a simple and intuitive format; a comprehensive set of data fields inclusive of temporal and spatial attributes; use of commonly accessible file types; and online hosting. This consultation was nested in appraisal of existing global hazard/disaster databases. Using evaluation of research outputs via Scopus® (13 keyword combinations, 5518 hits), alongside inspection of existing flood and/or disaster databases, i.e., ADCRC; CRED EM-DAT; Dartmouth Flood Observatory; and PAGES (Past Global Changes) flood working group. These were filtered to $n = 64$ sources and systematically reviewed detailing database structures/fields/hierarchies, data entry and validation protocols, analytical foci, and communication approaches. Whilst representing global knowledge, these sources are dominated by European databases reflecting a concentration of practice. Specifically focussed on Mediterranean and/or mountain catchments in Portugal ([65]: 'DISASTER'), Spain ([66]: 'INUNGAMA'; [67]: 'PRESS-GAMA'; [68]: 'PREDIFLOOD'), France ([69]: 'BDHI'), Germany ([70]: 'HANG'), Italy ([71]: 'APAT'; [72]: 'PEOPLE'), Switzerland ([73]: 'WSL'), Greece [74], and multi-nationally across the north-west Mediterranean ([37,75]: 'HYMEX').

For accessibility HiFlo-DAT is shared as an English language MS Excel © spreadsheet for textual data and Google Earth © mapping for locational data. The HiFlo-DAT architecture (Fig. 3) includes 'groupings' of like information and sub-divisions into 'categories'; these vary slightly between the 'unmerged sheet' (i.e., data entry of individual sources) and 'merged sheet' (i.e., flood event synthesis from one or more sources). Groups ($n = 10–11$) detail database management, source citation, event timing (when), location (where), causation (what), environmental and societal impacts (what and who), and the event management lifecycle. Herein 'categories' ($n = 80–95$) provide detailed quantitative and qualitative data according to entry rules.

3.3. Documentary source selections and capture approaches

HiFlo-DAT uses English-language documents from public and private collections in India, the UK and the USA. Since 2013 the research team have systematically evaluated materials, but most intensively 2018–2020, at the Prime Ministers' Museum and Library (PMML, formerly the Nehru Memorial Museum and Library [NMML], Delhi) and the British Library (London).

Table 2 details key repositories, their accessibility and the material types collected; including newspapers, reports, published literature and a wider array of materials inclusive of grey-literature (e.g., exploration and mountaineering accounts). These collections were selected and accessed with the aid of catalogues (where existing) alongside the expert knowledge of curators and research partners. These provide capacity to reconnect society with forgotten flood events, despite constraints on comprehensiveness associated with editorial style and decision making on inclusion for publication [76].

Table 3 details the overlapping sources used in HiFlo-DAT: dominated by national and regional newspapers and government reports for 1835–2020 (185 full years). Amongst digitally searchable newspapers (i.e., The Indian Express, The Times of India, and The Tribune) keyword combinations ($n = 22–92$) with spelling variants of hazard processes and location/person names permitted filtration at the point of collection, resulting in $n = 662$ files. These span 1838–2020 (i.e., 182 full years), with all month coverage, and continuous regional insight via 'The Tribune' from 1881.

In contrast, analogue sources (i.e., microfilm and hardcopy) were targeted reflecting time-cost constraint; largely limited to summer monsoon months (i.e., June–September). These correspond with the dominant rainfall-generated flood season in South Asia [77,78], including mountain states, such as Ladakh [79], Himachal Pradesh [46,54] and Uttarakhand [80]. Unfiltered materials ($n =$

Table 1
Key existing hazard/disaster databases incorporating historical floods in the Kullu District, Himachal Pradesh.

| Database | Author | Spatial Extent | Temporal Range | Number of Entries for Kullu District | Number of Entries for Manali/Kullu Tehsils (Years) | Public Availability and Link | Baseline Datasets | Comment |
|--|--|--------------------|----------------------------|---|--|--|--|---|
| District Disaster Management Plan (2017) | Kullu District Disaster Management Authority (DDMA) | Regional: District | 1988–2003 | 9, ‘prominent flash floods’ | 5 (1995–2003) | YES: https://hpkkullu.nic.in/documents-2/ | ? | NA |
| HVRA (‘Hazard Vulnerability and Risk Atlas’) | Himachal Pradesh State Disaster Management Authority (HPSDMA) | Regional: State | Unknown | c. 10 | c. 10 | PARTIALLY: http://www.hpsdma.hp.gov.in/Home_Disaster.aspx | Household surveys | Public version has redacted flood location data. However, screenshots of the full database maps in HPSDMA presentations show historical flood locations (no metadata) |
| IHCAP (Indian Himalayas Climate Adaptation Programme) | Indio-Swiss consortium | Regional: State | 1950–2014 | 44 ‘significant flood’ events for the State, and most in the Kullu District | ? | NO: previously available, but the website is now redundant | Scientific publications, technical reports, existing databases (DFO, EM-DAT) and media sources | IHCAP reports provide synopsis accounts, but more detailed data are not in the public domain |
| IMD (‘Disastrous Weather Events’, which populate the ‘Climate Hazards & Vulnerability Atlas’) | India Meteorological Department | National | 1967–2020 (excepting 1977) | c. 25 affirmative flood process accounts ^{#1} | c. 15 (1994–2019) | YES: https://imdpune.gov.in/library/publication.html | IMD reports and wider media information | Data are typically over-generalised/aggregated, making determination of process and location specifics challenging, which leads to count exclusions |
| ADRC | Asian Disaster Reduction Centre, Japan | Global | 1998-present ^{#2} | 1 | 1 (2003) | YES: https://www.adrc.asia/latest_disaster.php | Media, governmental/NGO reports and remote sensing sources | NA |
| DFO: ‘Global Active Archive of Large Flood Events’ (Dartmouth Flood Observatory) | University of Colorado, USA | Global | 1985-present ^{#2} | 7 | 3 (1994–2003) | YES on request: http://floodobservatory.colorado.edu/index.html | Media, governmental, instrumental, and remote sensing sources | Data on spatial location can be generalised, which prevents detailed local assessments |
| EM-DAT (‘Emergency Events Database’) | Centre for Research on the Epidemiology of Disasters (CRED), Belgium | Global | 1900-present ^{#2} | 5 | 3 (2003–2012) | YES: https://emdat.be/ | UN/NGO/insurance reports, research publications and media | Data on spatial location can be generalised, which prevents detailed local assessments |

^{#1} = IMD [64] reports 51 flood events in the Kullu District in the period 1969–2019; ^{#2} = July 2020 end point of database review here.




Fig. 2. Synopsis of the 'HiFlo-DAT' production journey.

>162,823 files) were captured using photography (.jpeg) and scanning (.pdf), enabling off-site review. Core were 'The Civil and Military Gazette', a regional newspaper (Lahore, Pakistan) alongside one of its precursor publications 'The Moffsilité'. These bring semi-continuous coverage 1845–1963 (104 of 119 years), although following partition (August 1947) information about India substantially declines.

3.4. Documentary source review approaches

Extracting relevant information from source materials was a substantial undertaking. The workflow is detailed in Supplementary Materials 1; key steps being: (1) team distributed review to identify potentially relevant information; (2) moderation of team selected materials resulting in a final selection. For consistency team data review was guided by a protocol note and contemporary map. A record sheet detailed whether an account had application (i.e., a negative or interim positive hit). Hereafter rigorous moderation confirmed that all materials were reviewed; followed by a check on whether accounts detail overbank flood processes and generate




Fig. 3. 'HiFlo-DAT' database architecture.

Table 2

Key documentary archive organisations in India, the UK and the USA underpinning 'HiFlo-DAT'.

| Country | Key Organisation | | | | | Accessibility Conditions | | | | | Material Type Collected | | | | | | |
|---------|---|---------------------|-----------------------------|---|--------------------------|--------------------------|---------|----------------|----------------|-----------------|-------------------------|--------------------|--------------------------|-----------------------------|-----------------------|---------|-------|
| | Name | Location | Type | Digital/ Online Catalogue (At Time of Access) | Access Year (s) | Open Access | Request | Paid Permit | Outputs Pay | Outputs Free | Newspaper | Report/ Records | Numerical Data Series | Book/ Article/ Thesis | Personal Documents | Webpage | Other |
| India | DDMA, Kullu District | Kullu | Government Authority | No | 2015, 2018 | ✓ | ✓ | | | ✓ | | | | | | ✓ | |
| | HPSDMA, Himachal Pradesh | Shimla | Government Authority | No | 2015, 2018 | ✓ | ✓ | | | ✓ | | | | | | ✓ | ✓ |
| | Himachal Pradesh State Archive | Shimla | Archive | Paper Offline | 2013–2015 | | ✓ | | | ✓ | | | | ✓ | ✓ | | |
| | Himachal Pradesh University | Shimla | University Library | No | 2013 | | ✓ | | | ✓ | | | | ✓ | | | |
| | ICAR Indian Agricultural Research Institute | Katra | Government Organisation | No | 2014, 2017 | | ✓ | | ✓ | | | | | ✓ | | | |
| | Indian Institute of Advanced Studies | Shimla | Research Library | Yes | 2015 | | ✓ | | ✓ | | | | | ✓ | | | |
| | Kullu Library | Kullu | Library | No | 2015 | | ✓ | | | ✓ | | | | ✓ | | | |
| | National Archives of India | Delhi | Archive | Yes | 2016 | | ✓ | ✓ | ✓ | | | | ✓ | | ✓ | | ✓ |
| | Prime Ministers' Museum & Library | Delhi | Archive | Yes | 2017–2018 | | ✓ | ✓ | ✓ | | ✓ | | | | | | |
| | Punjab State Archive | Chandigarh | Archive | Paper Offline | 2014–2016 | | ✓ | | | ✓ | | | ✓ | ✓ | ✓ | ✓ | |
| | Ratan Tata Library | Delhi | University Library | Yes | 2016 | | ✓ | | | ✓ | | | | ✓ | | | |
| | The Times of India | Online | Publisher Website | Yes | 2020 | ✓ | | | | ✓ | ✓ | | | | | | |
| | The Tribune | Chandigarh + Online | Publisher Archive + Website | Yes | 2015–2016, 2020 (Online) | ✓ | ✓ | | | ✓ | ✓ | | | ✓ | | | |
| | The Indian Express | Panchkula | Publisher Archive | Yes | 2017 | | ✓ | | ✓ | | ✓ | | | | | | |
| UK | British Library | London | Archive | Yes | 2016–2020, 2022 | ✓ | ✓ | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | | ✓ | |
| | Pagoda Tree Press | Bath | Archive | No | 2015, 2017 | | ✓ | | ✓ | ✓ | | | ✓ | | | ✓ | |
| | Penelope Chetwode Collection | Brighton | Family Archive | No | 2017 | | ✓ | | | ✓ | | | ✓ | ✓ | | ✓ | |
| USA | RGS-IBG | London | Archive | Yes | 2013 | | ✓ | | ✓ | | | | ✓ | | | ✓ | |
| | American Alpine Club | Golden, CO | Library | Yes | 2016 | | ✓ | | | ✓ | | | ✓ | | | | |
| | PAHAR | Denver, CO | Personal Collection | No | 2016 | | ✓ | | | ✓ | | | | | | ✓ | |

Table 3

Key continuous documentary data sources selectively used for 'HiFlo-DAT'.

| Source Series Details | | | | Timespan Reviewed | | | | Data Output Characteristics | | |
|--|--|-------------------|-----------------------------|-------------------|--------------------|------------|------------------------------|-----------------------------|-------------------------------|-----------------------------------|
| Publication Name | Provider | Focus | Newspaper (N) or Report (R) | Start Year | End Year | Year Count | Months | File Format | Number of Files ^{#6} | Filtered or Unfiltered at Capture |
| The Indian Express (Delhi Edition) | The Indian Express | National | N | 1954 | 2017 (April) | 64 | ALL | .JPEG | 36 | Filtered (Common Keywords) |
| The Indian Express (Chandigarh Edition) | Express | | | 1977 | 2010 | 34 | | | | |
| The Times of India | BL (British Library) Online | National | N | 1838 | 2005 | 168 | ALL | .PDF | 90 | |
| | | | | 2018 | 2020 (July) | 3 | | | 10 | |
| The Tribune | The Tribune Online | Regional | N | 1881 | 2016 | 136 | ALL | .PDF | 513 | |
| | | | | 2016 | 2020 (July) | 5 | | | 13 | |
| Civil and Military Gazette (CMG) ^{#1} | BL | Regional | N | 1876 | 1914 | 39 | June–Sept. (where available) | .PDF | 39,947 | Unfiltered |
| | | | | 1947 | 1949 ^{#5} | 3 | (1894 Jan.–Oct.) | | | |
| | PMML (Prime Ministers' Museum & Library) | | | 1915 | 1938 | 24 | June–Sept. (where available) | .JPEG | 65,461 | |
| | | | | 1947 | 1949 ^{#5} | 3 | | | | |
| | | | | 1956 | 1963 | 8 | | | | |
| Englishman | PMML | National | N | 1894 | 1894 | 1 | May–Sept. | .JPEG | 2109 | |
| India Administration Report (IAR) ^{#2} | BL | Regional Sections | R | 1855 | 1870 | 16 | Annual Publication | .PDF | 339 | |
| Mofussilite | BL | Regional | N | 1845 | 1845 | 1 | June–Sept. (where available) | .JPEG | 904 | |
| | | | | 1847 | 1875 | 29 | | .PDF | 6052 | |
| Report on the Administration of the Punjab Territories (RAPT) ^{#3} / Punjab Administration Report (PAR) | BL | Regional | R | 1849 | 1855 | 7 | Annual Publication | .PDF | 629 | |
| | | | | 1868 | 1883 | 16 | | .PDF | 1755 | |
| | | | | 1884 | 1934 | 51 | | .JPEG | 4865 | |
| The Friend of India/Statesman (FOI) ^{#1,4} | BL | National | N | 1835 | 1882 | 48 | June–Sept. | .PDF | 9507 | |
| | PMML | | | 1883 | 1889 | 7 | | .JPEG | (31,255) | |
| | | | | 1915 | 1927 | 13 | | | | |

#1 During the project BL holdings were microfilm only, since 2020 searchable digital copies are being uploaded to the British Newspaper Archive. CMG: <https://www.britishnewspaperarchive.co.uk/titles/civil-military-gazette-lahore> (In January 2024 = 1876–1951, 1954–1963).

FOI: <https://www.britishnewspaperarchive.co.uk/titles/friend-of-india-and-statesman> (In January 2024 = 1852–1883).

#2 IAR: A precursor national publication to PAR with Punjab regional sections.

#3 RAPT (1849–1855): A precursor to both the IAR (1855–1870) and PAR (1868–1934), showing changing reporting regimes.

#4 FOI: To prioritise, HiFlo-DAT only reviewed holdings sourced from the BL (1835–1882). PMML holdings (n = 31,255 files, and after 1927) remain un-reviewed.

#5 Repeated entry as different monthly holdings at PMML and BL for the Independence/Partition period, so best assembled from both sources.

#6 A file count of n = 1 varies between part of, single and double pages.

impacts in-area (i.e., Manali and Kullu Tehsils). Decisions were guided by local stakeholder consultation and cross-referral to modern and historical documents/maps. All accepted interim positive accounts were given a unique # source number (# 1–267); these were iteratively reduced to a final n = 220 # sources to populate the HiFlo-DAT database (section 3.5).

3.5. 'HiFlo-DAT' database population (unmerged and merged), validation and open access

As detailed in Fig. 3 and section 3.2, HiFlo-DAT has two Microsoft Excel © worksheets, i.e., unmerged and merged (n = 29,047 and 10,139 cells respectively). Like Lang et al. [69] this architecture facilitates initial data entry followed by flood event synthesis. This is necessary given an individual # source may provide details of multiple floods in time and space, whilst also recognising that knowledge of a discrete flood (F-event) is often a synthesis of one or more # sources. The workflow is detailed in Supplementary Materials 1 (steps 3–6) and 2.

Quantitative and qualitative data entry to the unmerged worksheet is strictly controlled, complying with cell entry explanations, does not over interpret the # sources, and uses third party information (e.g., maps, imagery, and websites) to best validate historical

information. Supplementary data includes historical rainfall records (see section 3.6), and carefully selected latitude and longitude coordinates using Google Earth. This generates $n = 308$ entries from $n = 220$ # sources.

The merged worksheet is generated by sorting # sources into flood event groups, first by date and second by location. Where a flood event ranges from valley-scale/coupled drainage network footprints to more localised occurrences. These groups are then synthesised into single rows of data constituting each F-event entry ($n = 128$). The sometimes fragmentary and contested nature of grouped # source data requires judgements to be made and data ranges to be specified; for consistency Supplementary Materials 2 details the rules





Fig. 4. Temporal analyses of floods in the Kullu District 1846–2020, (A) annual; (B) decadal; (C) monthly; and (D) seasonal.

(A)



(B)




Fig. 5. Spatial analyses of floods in the Kullu District, (A) flood event locations; and (B) frequency of floods by location.

adopted. All steps in this process are cross-checked to assure data accuracy. Fulfilling user group aspirations (section 3.2) the database outputs are open access via BathSPAdata (Resource 1).

3.6. Historical instrumental rainfall data series

Historical daily rainfall data (rain-day format) are integrated into HiFlo-DAT bringing better understanding of floods. These are for Naggar Farm (c. 1660 m AMSL; Lat. 32.115647°, Long. 77.160752°; 1891–1950 and 1962–May 2017) and Kullu (exact location unknown; 1891–1950). They offer a new and carefully constructed synthesis of ‘Daily Rainfall of India’ reports (1891–1950) held at the British Library (IOR/V/18/62–120), and ICAR-IARI-Katrain data 1962 to present-as partly used (1962–2009) by Jangra and Singh [81]. In using these data, it is recognised that individual floods may not be rainfall generated, and the recorded rainfall may not be spatially representative.

4. HiFlo-DAT results and discussion

Showcasing the contribution of HiFlo-DAT (merged) to knowledge of historical flooding in the Kullu District, key characteristics are evaluated, including temporal and spatial signatures, and environmental, societal and human impacts. Whilst these HiFlo-DAT results advance understanding of historical floods, it is important to caveat their fidelity. As they are drawn from ‘reported’ documentary data, it is appropriate to recognise they will not be an absolute record of historical flood occurrence and future flood risk across the region. Indeed, it may be the case that the spatial distribution of historical floods is in part informed by population density and infrastructure presence, which conditioned past awareness and societal interest.

4.1. Temporal characteristics

Over 175-years (1846–2020) using 127 of 128 events with an affirmative year, 57 years recorded one or more floods (32.6 % of all years), whilst the longest period of no-floods is 26 years (1912–1937). Corresponding mean annual frequencies are 0.73 events (all years) and 2.22 events (in flood only years). Herein, two years record 9 events (1894 and 2018); two years record 5 events (1846 and 1899); four years record 4 events (1898, 1901, 1994, 1998); and nine years record 3 events (1875, 1889, 1900, 1905, 1938, 1995, 2003, 2011, and 2015). Fig. 4(A) depicts the annual frequency, cumulative frequency and 10-year running mean of floods. Here attention is drawn to flood-rich periods, e.g. 1890–1900s and 1990s-present (i.e., steeper cumulative curve) in contrast to flood-poor periods (i.e., flatter cumulative curve). Employing a 1-tailed (positive) classic Mann-Kendall test with continuity correction, using XLSTAT, indicates a statistically significant trend in the time series. Being able to reject the null hypothesis (computed p-value less than α), in favour of the alternative hypothesis, signifies an increasing number of annual floods over 175 years ($K \tau = 0.205$; significance level $[\alpha] = 0.05$; p-value = 0.00024).

Fig. 4(B) shows 17 of 19 decades registering floods (all except the 1850s and 1920s), at mean decadal frequencies of 6.68 events (all decades) and 7.00 events (1840–2010; complete decades). The record shows five decades with higher frequencies (>10 events), each accounting for >10 % of the population. Again, the two flood-rich multi-decadal spans are evident, i.e. 1890-1909 (35 events, 27.6 %) and 1990–2019 (51 events, 40.2 %). The Mann-Kendall 1-tailed (positive) test returns no statistically significant trend in the series of event counts per decade ($K \tau = 0.127$; significance level = 0.05; p-value = 0.240).

Fig. 4(C) shows monthly frequencies and percentages of flood events; the importance of the summer monsoon season in generating floods is evident with June to September accounting for 111 events or 87.4 % of the population. Extending this, Fig. 4(D) uses 30-year periods according to India Meteorological Department (IMD) defined seasons. Here monsoon season dominance ranges to 50–100 % of events in each period but typically exceeds 80 %. Whilst it appears that the monsoon season accounts for an increasing proportion of flood events since the 1930s, the Mann-Kendall 1-tailed (positive) test returns no statistically significant trend of annual flood event frequency in the monsoon season 1869–2020 ($K \tau = 0.103$; significance level = 0.05; p-value = 0.051).


4.2. Spatial characteristics

Fig. 5(A) and Resource 2 show the location of all $n = 128$ floods. These incorporate all positional accuracy categories (see Resource 1), where S = specific location or sub-catchment, G = generalised as an indicative location or an extensive impact area, and U = unspecified regional location. On Fig. 5(A) there are $n = 59$ plotted points, reflecting multiple flood occurrences at some locations. Overall, these demonstrate a broad spread of historical floods along the length of the Kullu Valley, specifically the Beas River corridor and its tributaries. The Parbati River sub-catchment has fewer recorded flood event locations ($n = 9$), but this does not automatically translate into a lower future flood risk.

Fig. 5(B), using $n = 128$ floods, reveals the frequency distribution using graduated circles. The two purple circles are surrogate locations incorporating ‘U’ category positional accuracy data, and therefore potentially overstate the frequency of flood events at these locations but remain representative of wider impacts in the catchment for these individual events. The largest circle ($n = 22$, U = 15, G = 7) is the right bank of the Akhara Bridge over the Beas River (Lat. 31.962323°; Long. 77.115693°). The smaller circle ($n = 6$; U = 3, G = 3) is the mouth of the Parbati River (Lat. 31.898510°; Long. 77.148252°). These withstanding, overall, 76.3 % ($n = 45$ of 59) of ‘locations’ have a single recorded flood occurrence, the remainder have multiple recorded floods, with 10.2 % ($n = 6$ of 59) having two floods, and 13.6 % ($n = 8$ of 59) having three or more floods.

Refining this evaluation, Fig. 6(A) is a rank analysis, using a restricted dataset, i.e., 110 floods with affirmative locational data (i.e.,

(A)



(B)




Fig. 6. Spatial analyses of floods in the Kullu District, (A) rank analysis of flood spatial recurrence; (B) spatial pattern of floods over the entire time.

S and G categories). This shows the number of times each location was affected (y-axis) against the rank (x-axis, ranks 1–59), where the starting rank (1) is for the highest number of events. Here $n = 14$ locations with two or more recorded events account for 59.1 % of the total flood ‘events’ ($n = 65$ of 110), and those with a single recorded event are 40.9 % ($n = 45$ of 110). Overall, they are approximately power law distributed ($y = 10.522x^{-0.655}$; $R^2 = 0.93$). The location most afflicted by flooding is the Beas River ($n = 28$ of 110, 25.5 %), albeit this is spatially extensive, with events being recorded at multiple locations. At a more granular level, four Beas River right-bank tributaries are prevalent: Phojal Nalla (13 events, 11.8 % associated with a single dominant point location; but 14 events when considering multiple named points), Mohal Khad (10 events, 9.1 %), Bajaura Khad (9 events, 8.2 %, but 10 events at multiple named points), and Sarvari Nalla (5 events, 4.5 %; but 8 events at multiple named points). The fourth ranked (7 events, at the Akhara Bridge location in Fig. 5 B) is one of multiple locations associated with Beas River.

Fig. 6 (B), using 110 floods, reveals the reported distance density pattern for the entire 1846–2020 period, with 10 km (geodesic distance) concentric bands from a node at the Akhara Bridge on the right bank of the Beas River (Lat. 31.962323°; Long. 77.115693°). This shows a dominance of flood events within 20 km of Kullu and decreasing outwards. Specifically, 0–10 km (41 events, or 37.3 % of the total), 10–20 km (40, 36.4 %), 20–30 km (14, 12.7 %), 30–40 km (10, 9.1 %), 40–50 km (5, 4.5 %). However, spatio-temporal signatures are more complex when segregated into 30-year time slices from 1840 (excepting 2020- a single year). Herein, Fig. 7, based on 109 floods (as F12 has inexact timing) reveals clustering of floods within 20 km for 1840–1869 and 1900–1929; in contrast floods are consistently more widespread for 1870–1899 and 1990–2019, within 40 km and 50 km, respectively, corresponding to flood rich periods. The remaining periods 1930–1959, 1960–1989, and 2020 have some of the lower event frequencies and more variable patterns. A further analysis of the geodesic distances from the node to each flood position (excluding 2020, so $n = 108$), demonstrate that minimum distances are always < 1 km, and maxima fluctuate between 13.5 and 43.4 km. More revealing are average distances which in the aggregated period 1840–1959 range 8.3 ± 4.8 to 10.9 ± 9.2 km, compared to the aggregated period 1960–2019 ranging 15.1 ± 14.2 to 19.9 ± 13.3 km. This generalised outward expansion may reflect better reporting, growth of societal activities into more peripheral locations, and increasing event frequencies. Despite this, no floods are recorded in the headwaters of the Parbati river catchment (i.e., >50 km distance from Akhara Bridge).




Fig. 7. Spatial pattern of floods over 30-year time slices (1840–2020).

4.3. Hydro-meteorological and geomorphic characteristics

Interconnections between meteorological, hydrological and geomorphological processes are important in understanding floods. Herein, processes recorded as responsible for each flood ($n = 128$) were: 14.8 % river floods (long-duration rainfall), 39.8 % flash floods (short-duration rainfall), 39.8 % unspecified floods; with the remainder being landslides (3.1 %), snowmelt (0.8 %) or unknown (1.6 %). Withstanding the technical accuracy of source reporting, this suggests a dominance of rainfall causation. However, the likely importance of slope instability-channel coupling is poorly appreciated. For example, HiFlo-DAT captures named secondary processes, and these signal $n = 12$ landslides, $n = 1$ debris-flow, and $n = 3$ Landslide Lake Outburst Floods (LLOF) resulting from avalanche, landslide and earthquake derived barriers. Whilst seemingly rare, LLOFs in 1875 (F9), 1894 (F20) and 1905 (F52) generated large and impactful floods. As LLOF incidence is likely to increase with climate change and anthropogenic development in the region (e.g., [54, 62]), future DRR needs to better accommodate LLOF risk, which NDMA [82] acknowledges as a current gap.

Considering rainfall, reported quantitative data are sparse, including intensity ($n = 1$, 25 mm h^{-1} at Bhuntar airport, 2001), depth ($n = 5$: 63–127 mm, 1995–2018) and duration ($n = 20$: 0.5–61 h, 1888–2018). Wider qualitative narratives ($n = 92$ events) are typically too opaque for quantitative database entry, but they do indicate intensity is dominated by ‘heavy/torrential rainfall’ ($n = 49$ events) in contrast to ‘light rain’ ($n = 1$) and ‘showers/intermittent rain’ ($n = 3$). Duration expressions, include ‘steady-continuous/incessant/sustained’ rainfall ($n = 21$), and indicate many floods are associated with multiple episode rainfall ($n = 13$). Considering event type phrasing, ‘Thunderstorm/Cloudburst’ are dominant ($n = 38$ events) in contrast to ‘rain’ ($n = 7$), ‘monsoon’ ($n = 2$) and ‘snowmelt’ ($n = 2$). Changing vernacular is apparent, reporting ‘Thunderstorms’ 1889–1939, and ‘Cloudburst’ 1994–2020 (excepting 1899 and 1902). This language has wider implications, as many DRR agencies (e.g. [56,57]) often classify cloudbursts as an independent hazard process type, rather than an integrated meteorological mechanism which may result in channelised flood flows. This process-decoupling may hinder effective DRR in complex cascading systems.

Furthermore, independent rainfall records for Naggar Farm and Kullu (see section 3.6), offer some corroboration of event timing, rainfall event locations and antecedent conditions. For example, source narratives for the May 21, 1894 Phojal Nalla LLOF (F20) detail preceding heavy winter snowfalls, rain on the 15th May impacting the snow cover, and flood transferred snow deposits downstream. In corroboration, the Naggar Farm data (c. 10 km east), usefully detail: (1) 18/4/94 to 11/5/94 1 mm of rainfall, and thus likely sunnier days resulting in snowpack metamorphosis; (2) 12–16/5/94 59.4 mm of rainfall, which would have enhanced snow melt or brought new snow layers at elevation; and (3) 17–21/5/94 0 mm of rainfall. This record likely supports the absence of rainfall flood causation, and instead a snow avalanche/landslide flood mechanism.

Considering flood flow characteristics, alike rainfall quantitative data are sparse. Discharge records ($n = 3$: 2482–44,855 $\text{m}^3 \text{s}^{-1}$), are reconstructed (Phojal Nalla and Seri Nalla/Beas River, F20 and F11) and measured (Beas River, F86). Wider narrative expressions are abundant, detailing: location (main river [$n = 27$], and tributary [$n = 20$]), timing ($n = 18$), magnitude (velocity [$n = 2$], relative size [$n = 23$], stage [$n = 14$]) and hydrograph changes (rising and falling stage [$n = 20$], and flood waves [$n = 4$]). Whilst accounts of sediment load/turbidity are rarer ($n = 6$).

Table 4
Frequency and magnitude of flood event impacts on environmental and societal assets.

| Asset Group | Asset Type Damaged | Flood Events Recording Impact to the Asset Type (Frequency) | Sum Frequency of Impacts for Asset Group | Sum Magnitude of Impacts to the Asset Type |
|--|--|---|--|--|
| Buildings & Property | Domestic property | 22 | 52 | $n = 445$ |
| | Shops/Stalls/Kiosks | 12 | | $n = 159$ |
| | Other Business/Hotels/Industrial | 8 | | $n = 80$ |
| | Communal property | 9 | | $n = 52$ |
| | Religious/Cultural | 1 | | $n = 2$ |
| Transport Infrastructure | Bridge | 54 | 118 | $n = 94$ |
| | Road | 55 | | 47.3 km |
| | Track (unsurfaced) | 2 | | 2 reports |
| | Pedestrian pathway | 7 | | 7 reports |
| Vehicles | Damaged | 4 | 10 | $n = 16$ |
| | Washed away | 6 | | $n = 13$ |
| Water & Slope Infrastructure | Watermills | 5 | 30 | $n = 58$ |
| | Irrigation channels (Kuhls) | 2 | | 2 reports |
| | Protective engineering (channel & slope) | 8 | | 8 reports |
| | HEP/Dams | 4 | | 4 reports |
| | Water supply | 11 | | 11 reports |
| Power & Communication Utilities | Electrical power | 9 | 19 | 9 reports |
| | Telecommunications | 5 | | 5 reports |
| Plants, Trees & Animals | Postal | 5 | 55 | 5 reports |
| | Forestry land | 14 | | 14 reports |
| | Agricultural/Horticulture land | 28 | | 102.6 ha |
| | Orchards (trees) | 5 | | $n = 758$ |
| | Livestock | 8 | $n = 430$ | |

Reported geomorphological impacts include channel incision ($n = 8$), channel aggradation ($n = 18$), avulsion ($n = 21$), bank erosion/failure ($n = 15$), overbank deposition ($n = 21$), tree damage/woody debris ($n = 32$), and slope instability ($n = 19$). Extracting quantities from narrative accounts ($n = 55$), reveal over bank deposits ranged from fines to boulders up to 6 m deep, but inundation areas are rarely reported; in channel deposits are boulder sized up to 40 m deep; lateral channel shifts up to 500 m; and tree damages up to the thousands in a single event. These are likely a snapshot, as no source was a dedicated geomorphological appraisal.

4.4. Environmental and societal asset impact characteristics

Table 4 synthesises impacts to environment and society assets; whilst likely subject to source biases and incomplete capture/quantification, it provides insight into historical flood losses. Frequency counts express the number of flood events recording the impact type, and magnitude of impact is more diverse according to the asset type, comprising counts of individual assets, counts of reports (the same as frequency), and distance/area. Data reveal a concentration on transport infrastructure, in particular bridges (54 floods, 94 impacts) and roads (55 floods, 47.3 km of damage/burial/interruption). Though also noteworthy are the impacts to domestic properties (22 floods impacting 445 buildings) and agriculture and horticultural land (28 floods, 102.6 ha).

Further evaluating bridge impacts reveals the number of bridges damaged/lost in these floods range 1–12 (average 1.74 bridges). Temporally, over 19 decades, 12 record impacts and 7 do not; 4 decades account for 67 % of all reported bridge impacts: 1890s (20.4 %), 1900s (18.5 %), 1950s (13.0 %) and 2010s (14.8 %). Spatially, qualitative accounts detail 81 impacts (of the 94 total), at 29 locations, of which nearly half have a single bridge impact, and the remainder experienced repeated bridge impacts. Most accounts (65 %) detail bridge losses, opposed to bridge damage (35 %). These are identified as road bridges (35 %), foot bridges (20 %) or unspecified (45 %). The top 5 locations of recurring impacts are Dobhi ($n = 12$, 1894–2018, largely Phojal Nalla tributary, with episodes 1894, 1952–1957 and 2018); Kullu ($n = 8$, 1846–1995, largely Sarvari Nalla tributary); Bajaura ($n = 8$, 1899–1903, Bajaura Khad tributary); Mohal ($n = 7$, 1887–1902, Mohal Khad tributary); and Akhara in Kullu ($n = 6$, 1894–2019, Beas River [Main River]). These data reveal that Kullu, the largest settlement, is most afflicted. Further, tributary rivers are dominant, exhibiting short episodic impact periods (i.e., Bajaura and Mohal) and recurring impacts (i.e., Dobhi, Kullu).

Considering road impacts, these are recorded 1846–2020, in which 13 decades record impacts and 6 do not. Herein 5 decades each record ≥ 10 % of occurrences (i.e., 1890s, 1950s, 1990s, 2000s, 2010s). Moreso, 74.5 % of occurrences are since 1950, and 1990–2019 accounts for 50.9 %. This skew perhaps reflects more recent expansion of road networks and motorised transport. Spatially, 36.4 % of events have local impacts, 47.3 % are undefined, and extensive impacts are 16.3 %, including in 1894, 1995 and 2018. Narratives are thin, for example, 8 detail the length of impacted road (0.15–30 km), 6 give the duration of disruption (3 h–4 days), and a single account outlines the recovery cost (F86, September 1995). Exploring the nature of impacts, terminology is opaque but indicates: 21 events caused damage/disruption; 19 breached/inundated the road; 12 caused loss by destroying/washing away the road; and 17 impacted prevented/disrupted traffic flows.

Establishing reporting rationales for environmental and societal impacts is challenging, but they may reflect societal interest at the time of reporting. For example, Sah and Mazari [83], Berkes et al. [84] and Johnson et al. [60] remark that the traditional mountain economy of the Kullu District focusses attention on productive land which provides subsistence and income via market trade, access to which is governed by roads and bridges. This focus on transport infrastructure continues, as it underpins the growing tourist economy. For example, in the July–August 2023 Himachal Pradesh flood-landslide disaster, particular concern surrounded impacts on transport infrastructure. Closing 1300 roads and washing away c. 40 bridges [85], with estimated recovery costs of Rs 2458 Crore (c. £235 million in 2024), being 27 % of the total recovery cost [24].

4.5. Human impact characteristics

Selected quantitative and qualitative data highlight the harmful impact of floods on humans. The former being: (1) 'Total Fatalities' (i.e., sum of fatalities and the missing), hereafter 'fatalities'; and (2) 'Total Affected' (i.e., sum of the homeless and injured), hereafter 'affected'. HiFlo-DAT records 253 fatalities, across 24 events, where deaths per fatal event (range 1–50, average 10.5) occur in two periods: 1894–1905 (6 events and 78 fatalities) and 1994–2018 (18 events and 175 fatalities). Furthermore, 1322 people were affected across 15 events (range 1–500, average 88.1 people per event), occurring in two time periods: 1889 (1 event and 40 affected) and 1994–2012 (14 events and 1282 affected). These data show losses across all three centuries, but with a strong skew from the 1990s, although the 1894 Phojal Nalla flood recorded the largest fatalities ($n = c. 50$).

Qualitative accounts reveal vulnerability and exposure dimensions. Regarding vulnerability an important recipient population (i.e., who) are those associated with the keywords: migrant, labourer, herder, trader, worker and muleteer. 69.6 % ($n = 176$) of all fatalities are associated with event narratives with one or more of these words (8 events; flood event numbers [see Resource 1] = 20, 35, 81, 96, 97, 99, 108, 113). With the same filters (5 events; flood event numbers = 97, 99, 101, 108, 111) are associated with 1090 or 82.5 % of all affected persons. These are important findings, evidencing demographic vulnerability in a mountain setting; such contributions are considered pivotal by the UNDRR [32] in overcoming aggregated data obstacles.

Regarding exposure (i.e., where and what), the activities of the recipient population notably include those engaged in construction work typically in remote tributary locations (keywords: cable laying, construction, HEP, road avalanche shed), and importantly many were encamped in labourer colonies near to rivers at the time of the flood events (keywords: sleep, camp, community). 48.2 % ($n = 122$) of all fatalities are associated with event narratives with one or more of these words (7 events; flood event numbers = 35, 81, 97, 99, 100, 108, 113). With the same filters (4 events; flood event numbers = 97, 99, 100, 108) are associated with 191 or 14.4 % of all affected persons. Furthermore, many fatal losses occurred during the night hours (23:25 to 03:00 IST) with no or limited warning.

These analyses in the Kullu District highlight the recent vulnerability and exposure of migrants/labourers engaged in large-scale construction activities. Unfortunately, these circumstances recur across the Indian Himalaya, for example, high-profile events in Uttarakhand include the February 2021 Dhauliganga flood at Raini HEP and Tapovan HEP (with many fatalities), and November 2023 Silkyara Bend–Barkot tunnel collapse (with no fatalities, but large rescue costs). To reduce risks, further investigation is warranted, including consideration of what DRR benefits may be derived from more effective development/construction planning, design and management.

5. Wider discussion: applications of ‘HIFLO-DAT’

5.1. Enhanced knowledge for informed decision-making in the Kullu District

HiFlo-DAT detailing 128 floods, at 59 locations, over 175 years (1846–2020) in the Manali and Kullu Tehsils (Kullu District) reconnects society to past knowledge of flood occurrence and impacts. This is a major contribution, substantially improving upon existing compilations of past floods (Table 1: 1–15 entries from c. 1994) in terms of event frequency, timespan, and depth of information. Furthermore, analyses of HiFlo-DAT highlight increasing flood occurrence towards the present; dominance of rainfall causation in the monsoon season; high-magnitude LLOF events; hotspot tributaries subject to repeated floods (subject to limitations of documentary reporting patterns); and key impact receptors, namely roads, bridges and labourer-migrant communities associated with construction projects. These data and analyses are only of value if they inspire and inform action [32] strengthening local government and NGO decision making. Future tangible gains may include improved project selection and design for both disaster risk reduction/climate change adaptation (i.e., mitigation of hazard exposure and vulnerability, and resilience generation) and infrastructure development schemes (e.g., hydro-electric power, road widening and tunnelling, planned railways and airport runway upgrading, and settlement expansion). For example, in the absence of HiFlo-DAT, in May 2024 the HP State Executive Committee [86] approved substantial funding for flood protection works (Rs. 1761.57 Crore, c. £161.1 million in 2024). These 11 projects focus on the Beas River corridor, adjoining an at-risk national highway and key settlements; however, HiFlo-DAT brings alternative focus on tributary flood hazard/disaster losses. This perspective supports a growing call for informed decision making in the Kullu District, in respect to hazard and climate change challenges. For example, Allen et al. [63] detail how risk assessments using dendrogeomorphic and Glacial Lake Outburst Flood (GLOF) modelling data serve as ‘input for the prioritisation and design of adaptation actions’. Similarly, Chand et al. [87] articulate how flood vulnerability assessments strengthen resource allocation decisions. Whilst Gupta et al. [47] underline the need for hydro-meteorological process-cascade understanding for evidence-based policy making.

5.2. Transference of the methodology to the wider Himalaya and across trans-boundary basins

Whilst HiFlo-DAT is of direct value to the Kullu District, it can also help address fragmentary disaster loss data across the Indian Himalayan Region [29]. Specifically, HiFlo-DAT brings a grounded (i.e., best-practice and locally aligned) and robust methodology (see section 3) for the selection, capture, review, synthesis and analysis of documentary flood data. Wider application is however influenced by the quantity and quality of historical documentary sources. Accordingly, database assembly in more remote locations, with less established documentary histories than in the Kullu District case, may benefit from greater incorporation of grey-literature and local knowledge sources.

In offering this perspective, we are mindful of the time, cost and endeavour required to access and process documentary data sources, especially for dispersed and hardcopy collections. Therefore, as a pilot-scale test of transferability, in May–June 2023, we took the HiFlo-DAT methodology to the high-altitude Dhauliganga catchment upstream of Joshimath (Chamoli District, Uttarakhand). This location was selected to bring multi-event historical context to the large-scale February 2021 disaster ([14,15]) which may otherwise dominate local disaster risk reduction narratives, locations, and policy. Like the Kullu District approach, we explored documentary archives; this time prioritising digital resources for accessibility, particularly via the British Library, supplemented by review of hardcopy materials held by the Chamoli District Government in Gopeshwar. In addition, drawing on expertise [59,60] we hosted workshops in mountain villages (i.e., Raini, Pagarsu, Kosa and Malari) to explore local knowledge of past hazard-disaster events. The interim headline is one of success in deploying the HiFlo-DAT methodology to a different location, where local knowledge was of elevated importance given fewer accounts emerged from documentary records in this very remote location. This result should bring confidence that with appropriate team expertise and project resources, the transferability of the HiFlo-DAT methodology is both beneficial and achievable.

Of further merit would be trans-boundary implementation of HiFlo-DAT, given hazard process cascades may transcend national boundaries. Existing national [88] and international policy positions (e.g., [21,22,89,90]) already make the case for such trans-boundary hazard-disaster collaboration. Such may be particularly valuable for higher-magnitude LLOF/GLOF floods, which are likely to increase in frequency associated with climate change driven glacial retreat and lake formation, permafrost degradation and slope instability. Indeed, NDMA [82] recommend the need for a ‘systematic database of GLOFs and LLOFs’ as a foundation for more effective future management. For example, in response to the trans-boundary (China [Tibet]–India), 2000 and 2005 River Sutlej LLOF disasters [91], early warning systems [82] were implemented. It would however be interesting to re-evaluate these interventions with the benefit of new long-term historical flood knowledge.

5.3. Reflections on HiFlo-DAT data quality and further validation opportunities

Within HiFlo-DAT the technical specificity, depth and breadth of information afforded by documentary sources are variable. This likely reflects differential access to flood locations, lagged communication, audience/societal interest, reporting/editorial biases, author roles/responsibility/positionality, language, available technologies and technical expertise [76,92,93]. Whilst more recent accounts tend to offer greater detail, there are notable exceptions. For example, floods at Phojal Nalla (F20, 1894) and Bajaura Khad (F35, 1899) are well articulated. Conversely, the September 2018 Phojal Nalla flood (F124), may seem to offer a reasonable level of detail; however, our site inspections in October 2018, could significantly extend detail of geomorphological and societal impacts. This may raise questions about the factors influencing the inclusion and calibre of data entries in HiFlo-DAT. Both are conditioned by the original assembly of information, the subsequent accessibility of information, and how it is processed. As detailed in section 3, a robust regime was implemented in the construction of HiFlo-DAT, including filtering out accounts which did not clearly articulate evidence of flood processes or were opaque (e.g., elevated flow discharge), cross-referencing sources/existing databases and exploring third-party data such as Google Earth. Whilst HiFlo-DAT offers a significant step forward, it is bound by these data quality and quantity contexts. It is therefore valuable to highlight opportunities which may extend and test these and other historical flood databases. Systematic developments may include comparison to: (1) existing dendrochronological flood data in the Kullu District ([62,94]); (2) new multi-disciplinary (hydro-meteorological, bio-geomorphic and sedimentological) flood reconstructions using landscape evidence of past flood impacts [95,96]; (3) local knowledge oral accounts, for example the 1894 Phojal Nalla flood is recounted in song ([60]; <https://vimeo.com/285841577> [12 min. 06 sec.]); (4) village documents such as Panchayat record books [60,97]; (5) local language newspaper archives and video media; (6) untapped English language newspapers, increasingly accessible via digitisation; (7) diaspora family records not in official collections; (8) a wider array of Government of India data, such as a deeper dive into data underpinning the IMD 'Climate Hazards and Vulnerability Atlas' (Table 1; [64]) and the Central Water Commission (CWC) 'Flood Damage Statistics 1953–2020' [98]; and (9) modelling outputs.

5.4. Developing future DRR policy and practice

The introduction details current DRR policy-practice, including the importance of methodologies for the compilation of disaggregated disaster loss data at sub-national scales. Contributing to this agenda, we offer reflections related to: (1) specificity of historical databases; (2) data specificity and partnerships using technology for recording contemporary event impacts; and (3) access to archive data in service of global challenges.

UNDRR [32] in articulating the importance of data disaggregation, do so particularly regarding socio-economic metrics and generating data for contemporary events. We would extend this further, as HiFlo-DAT demonstrates the value of assembling disaggregated architectures across all metrics for past flood events. Having more detailed historical data will support the review and updating of existing key documents (e.g., Disaster Management Plans) and databases such as state-level HVRA collections [26].

The September 2018 Phojal Nalla flood (F124) revealed an untapped opportunity in developing more detailed records of contemporary events. We were able to compare the depth of our knowledge from live-time social media and field reconnaissance a month afterwards, versus that conveyed by the media and official reports. This demonstrated the residual capacity of technical knowledge/collaboration across sectors and disciplines. Whilst also highlighting the role digital technology (e.g. social media imagery/videos, mobile apps, and emerging AI capabilities) could contribute to capturing and evaluating flood processes and impacts. Such technologies are a growing enabler for decentralised/local community engagement and ownership of hazard and disaster knowledge at a local level [32,59,82,88,99].

Ease of access to archives is variable (Tables 2 and 3), and at times the experience is inefficient, restricted, and bound by paywalls; this only acts to inhibit disaster risk reduction and sustainable development progress. Whilst growing digitisation and open access collections may reduce the time and costs for data mining of historical data [100], many collections remain outside this scenario. Accordingly, in serving societal interest, there is a pressing need (e.g., [101]) for better inter-agency connectivity to facilitate improved access to public archive collections. Furthermore, efforts would be accelerated by long-term research funding supporting the systematic review of archive materials aligned to global challenges.

6. Conclusions

Disaster risk challenges are prominent in mountain environments and Asia more generally, where floods are a significant hazard/disaster loss process. International disaster risk reduction efforts, including the UNDRR Sendai Framework, highlight prevailing gaps in disaster loss databases, especially at sub-national scales. Accordingly, developing methodologies for the systematic assembly and analysis of such disaggregated data is a pressing need to reduce future disaster losses in an era of climate change. 'HiFlo-DAT' (Himalayan Flood Database) is an important contribution to this agenda, initially focussing on the Kullu District, Himachal Pradesh, India. It delivers a sizeable improvement in local knowledge, detailing 128 historical flood events, at 59 locations, over 175 years (1846–2020) reconnecting society/governing organisations/NGOs to their past documentary knowledge of historical flood occurrence and impacts. Further synopsis and visualisation is given by the accompanying HiFlo-DAT Film (<https://vimeo.com/1037424595>). Going forward our discussions highlight new opportunities, in overview:

- (1) Engage responsible agencies to embrace historical flood data to better inform their disaster risk reduction/climate change adaptation strategies, and to assist sustainable design of infrastructure projects in mountain environments.

- (2) The wider Indian Himalaya and trans-boundary basins would benefit from systematic approaches to assembling and sharing historical flood data, alike HiFlo-DAT.
- (3) Documentary based databases should be compared, corroborated and extended, where possible, using an array of sources, including local language and local community data, Government of India collections, diaspora accounts, and dendrochronological/hydro-geomorphic reconstruction. As an array of data are required to inform more comprehensive future flood risk assessments.
- (4) Recording details of future events would benefit from multi-agency partnerships to deliver richer information aligned to database needs and should consider how to effectively capture digital data.
- (5) Improving historical archive access, via new inter-agency cooperation would enhance strategic disaster risk reduction and sustainable development needs.

CRedit authorship contribution statement

Richard M. Johnson: Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **Bindhy Wasini Pandey:** Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Funding acquisition, Data curation, Conceptualization. **Kesar Chand:** Resources, Project administration, Methodology, Formal analysis. **Ceri L. Davies:** Writing – review & editing, Visualization, Software, Formal analysis. **Debra Edwards:** Investigation, Formal analysis, Data curation. **Esther Edwards:** Supervision, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **James Jeffers:** Methodology, Investigation, Formal analysis, Data curation. **Kieran King:** Methodology, Investigation, Formal analysis, Data curation. **Jagdish Chandra Kuniyal:** Writing – review & editing, Resources, Project administration, Methodology, Investigation, Funding acquisition, Conceptualization. **Himanshu Mishra:** Writing – review & editing, Resources, Project administration, Methodology, Investigation, Data curation. **Victoria Phillips:** Methodology, Investigation, Formal analysis, Data curation. **Nikhil Roy:** Methodology, Investigation, Data curation. **Jessica Seviour:** Methodology, Investigation, Formal analysis, Data curation. **Dev Dutt Sharma:** Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Funding acquisition, Data curation, Conceptualization. **Pushpanjali Sharma:** Methodology, Investigation, Formal analysis, Data curation. **Harkanchan Singh:** Methodology. **Ram Babu Singh:** Methodology, Funding acquisition, Conceptualization.

Data availability

Resource 1 ‘HiFlo-DAT’ database source and flood event spreadsheets (.xlsx): <https://doi.org/10.17870/bathspa.28053218>.

Resource 2 ‘HiFlo-DAT’ event locations (.kmz): <https://doi.org/10.17870/bathspa.28053254>.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

Sincere thanks to all archive organisations in India, the UK and the USA who have provided access to documentary source materials, key here are: The British Library (London, UK), The Prime Ministers’ Museum and Library (Delhi, India), The Tribune [Newspaper] (Chandigarh, India), and ICAR-IARI-Katrain (Kullu District, India). We also express thanks to our graduates and post-graduates for their sustained endeavours in this research, and to reviewers for their expertise which assisted development of this article. This work was supported by UGC-UKIERI (IND/CONT/G/20–21/A2 [G/17–18/19]) 2018–2021, and by Bath Spa University, the University of Delhi and the University of Cumbria. In kind contributions (i.e., time and expertise) were provided by project workshop delegates, including government agencies (e.g., Himachal Pradesh State Disaster Management Authority, Kullu District Disaster Management Authority), NGOs (e.g., DOERS, Geohazards Society India, IUCN India Office); creative industry (Kraft Films-Bengaluru); and Dr Vishwa Chandel (Panjab University, GIS data). The data selections, compilations and views expressed are those of the authors and are not conditioned by the research funders. Map lines delineate study areas and do not necessarily depict accepted national boundaries. We also pay our respects to our dear friend, colleague and team member, Prof. R B Singh (University of Delhi), for his insightful contributions to the project, prior to his passing in 2021.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijdr.2025.105336>.

Data availability

Research data are shared by an institutional data repository. The DOI links are given in the article.

References

- [1] J.S. Gardner, Risk complexity and governance in mountain environments, in: Fra, U. Paleo (Eds.), *Risk Governance*, Springer, Dordrecht, 2015, pp. 349–371, <https://doi.org/10.1007/978-94-017-9328-5>.
- [2] A. Walz, A. Grêt-Regamey, S. Lavorel, Social valuation of ecosystem services in mountain regions, *Reg. Environ. Change* 16 (2016) 1985–1987, <https://doi.org/10.1007/s10113-016-1028-x>.
- [3] R. Hock, G. Rasul, C. Adler, B. Cáceres, S. Gruber, Y. Hirabayashi, M. Jackson, A. Käb, S. Kang, S. Kutuzov, A.I. Milner, U. Molau, S. Morin, B. Orlove, H. Steltzer, High Mountain areas, in: H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (Eds.), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, 2019. https://www.ipcc.ch/site/assets/uploads/sites/3/2022/03/04_SROCC_Ch02_FINAL.pdf.
- [4] E. Sharma, D. Molden, A. Rahman, Y.R. Khatiwada, L. Zhang, S.P. Singh, T. Yao, P. Wester, Introduction to the Hindu Kush Himalaya assessment, in: P. Wester, A. Mishra, A. Mukherji, A.B. Shrestha (Eds.), *The Hindu Kush Himalaya Assessment: Mountains, Climate Change, Sustainability and People*, Springer, Nature Switzerland AG, Cham, 2019, pp. 1–16. <https://link.springer.com/book/10.1007/978-3-319-92288-1>.
- [5] C. Adler, P. Wester, I. Bhatt, G. Huggel, G.E. Insarov, M.D. Morecroft, V. Muccione, A. Prakash, Cross-chapter paper 5: mountains, in: H.-O. Pörtner, D. C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Lösckke, V. Möller, A. Okem, B. Rama (Eds.), *Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2022. https://www.ipcc.ch/report/ar6/wg2/downloads/report/IPCC_AR6_WGII_CrossChapterPaper5.pdf.
- [6] C. Mathison, A. Wiltshire, A.P. Dimiri, P. Falloon, D. Jacob, P. Kumar, E. Moors, J. Ridley, C. Siderius, M. Stoffel, T. Yasunari, Regional projections of North Indian climate for adaptation studies. *Science of The Total Environment* (2013) 468–469, <https://doi.org/10.1016/j.scitotenv.2012.04.066>. Supplement, S4–S17.
- [7] A. Dixit, S. Sahany, S.K. Mishra, Modeling the climate change impact on hydroclimate fluxes over the Beas basin using a high-resolution glacier-atmosphere-hydrology coupled setup, *J. Hydrol.* 627 (A) (2023) 130219, <https://doi.org/10.1016/j.jhydrol.2023.130219>.
- [8] R. Krishnan, A.B. Shrestha, G. Ren, Unravelling climate change in the Hindu Kush Himalaya: rapid warming in the mountains and increasing extremes, in: P. Wester, A. Mishra, A. Mukherji, A.B. Shrestha (Eds.), *The Hindu Kush Himalaya Assessment: Mountains, Climate Change, Sustainability and People*, Springer, Nature Switzerland AG, Cham, 2019, pp. 57–97. https://link.springer.com/chapter/10.1007/978-3-319-92288-1_3.
- [9] K.K. Rao, S.K. Patwardhan, A. Kulkarni, K. Kamala, S.S. Sabade, K.K. Kumar, Projected changes in mean and extreme precipitation indices over India using PRECIS, *Global Planet. Change* 113 (2014) 77–90, <https://doi.org/10.1016/j.gloplacha.2013.12.006>.
- [10] J.A. Ballesteros-Cánovas, T. Koul, A. Bashir, J.M. Bodoque del Pozo, S. Allen, S. Guillet, I. Rashid, S.H. Alamgir, M. Shah, M. Sultan Bhat, A. Alam, M. Stoffel, Recent flood hazards in Kashmir put into context with millennium-long historical and tree-ring records, *Sci. Total Environ.* 722 (2020) 137875, <https://doi.org/10.1016/j.scitotenv.2020.137875>.
- [11] UNDRR, Scoping study on compound, cascading and systemic risks in Asia-Pacific, United Nations Office for Disaster Risk Reduction - Regional Office for Asia and Pacific (2022) 60. <https://www.undrr.org/publication/scoping-study-compound-cascading-and-systemic-risks-asia-pacific>.
- [12] S.K. Allen, P. Rastner, M. Arora, C. Huggel, M. Stoffel, Lake outburst and debris flow disaster at Kedarnath, June 2013: hydrometeorological triggering and topographic predisposition, *Landslides* 13 (2016) 1479–1491, <https://doi.org/10.1007/s10346-015-0584-3>.
- [13] D.P. Dobhal, A.K. Gupta, M. Mehta, D.D. Khandelwal, Kedarnath disaster: facts and plausible causes, *Curr. Sci.* 105 (2) (2013) 171–174. <https://www.currentscience.ac.in/Volumes/105/02/0171.pdf>.
- [14] D.H. Shugar, M. Jacquemart, D. Shean, et al., (50 more authors), A massive rock and ice avalanche caused the 2021 disaster at Chamoli, Indian Himalaya, *Science* 373 (6552) (2021) 300–306. <https://www.science.org/lookup/doi/10.1126/science.abh4455>.
- [15] NDMA, Study of causes & impacts of the Uttarakhand disaster on 7th February 2021, in: Raunthi Gadhera (Ed.), *Rishiganga & Dhauliganga Valley: Measures to Reduce Disaster Risks*. National Disaster Management Authority, New Delhi, May 2022, 2022. https://ndma.gov.in/sites/default/files/PDF/Reports/Detailed_report_UK_Disaster.pdf.
- [16] A.B. Shrestha, J. Steiner, S. Nepal, S.B. Maharjan, M. Jackson, G. Rasul, B. Bajracharya, Understanding the Chamoli flood: Cause, process, impacts, and context of rapid infrastructure development, in: <https://www.icimod.org/article/understanding-the-chamoli-flood-cause-process-impacts-and-context-of-rapid-infrastructure-development/>, 2021.
- [17] R.A. Vaidya, M.S. Shrestha, N. Nasab, D.R. Gurung, N. Kozo, N.S. Pradhan, R.J. Wasson, A.B. Shrestha, C.G. Gurung, A. Bajracharya, Disaster risk reduction and building resilience in the Hindu Kush Himalaya, in: P. Wester, A. Mishra, A. Mukherji, A.B. Shrestha (Eds.), *The Hindu Kush Himalaya Assessment: Mountains, Climate Change, Sustainability and People*, Springer, Nature Switzerland AG, Cham, 2019, pp. 389–419. <https://link.springer.com/book/10.1007/978-3-319-92288-1>.
- [18] CRED-UNISDR, Poverty & Death: Disaster Mortality 1996–2015, UNISDR (now UNDRR), 2016. Geneva, <https://www.undrr.org/publication/poverty-death-disaster-mortality-1996-2015>.
- [19] CRED-UNDRR, The human cost of disasters: an overview of the last 20 years (2000-2019), Geneva, <https://www.undrr.org/publication/human-cost-disasters-overview-last-20-years-2000-2019>, 2020.
- [20] M.S. Shrestha, W.E. Grabs, V.R. Khadgi, Establishment of a regional flood information system in the Hindu Kush Himalayas: challenges and opportunities, *Int. J. Water Resour. Dev.* 31 (2) (2015) 238–252, <https://doi.org/10.1080/07900627.2015.1023891>.
- [21] UNDRR, Asia-pacific action plan 2021-2024 for the implementation of the Sendai framework for disaster risk reduction 2015-2030, United Nations Office for Disaster Risk Reduction - Regional Office for Asia and Pacific (2021) 17. <https://www.undrr.org/publication/asia-pacific-action-plan-2021-2024-implementation-sendai-framework-disaster-risk>.
- [22] UNDRR, APMCDRR 2024 Co-chairs' statement, United Nations Office for Disaster Risk Reduction - Regional Office for Asia and Pacific (2024) 6. <https://www.undrr.org/publication/apmcdrr-2024-co-chairs-statement>.
- [23] AMCDRR, Action plan 2018-2020 of the Asia regional plan for implementation of the Sendai framework for disaster risk reduction 2015-2030. https://www.preventionweb.net/files/56219_actionplan20182020final.pdf, 2018.
- [24] HPSDMA, Report on post disaster needs assessment, Himachal Pradesh monsoon - 2023, floods, cloudbursts and landslides. Himachal Pradesh state disaster management authority, Shimla (2023) 202. <https://hpsdma.nic.in/admnis/admin/showing.aspx?ID=3570>.
- [25] HPSDMA, Revised memorandum of damages (as of 15-09-2023) due to flash floods, floods, cloudbursts and landslides during monsoon season – 2023. Government of Himachal Pradesh Revenue Department (Disaster Management Cell) (2023) 191. <https://hpsdma.nic.in/admnis/admin/showing.aspx?ID=3573>.
- [26] NDMA, Proceedings of National Webinar with States/Union Territories for conducting a Hazard Vulnerability and Risk Assessment (HVRA) study. National Disaster Management Authority (2021) 35. New Delhi, June 2021, <https://www.ndma.gov.in/sites/default/files/PDF/Technical%20Documents/NDMA-Proceedings-of-Webinar-on-HVRA.pdf>.
- [27] NRSC, Flood Affected Area Atlas of India (1998-2022), Satellite Based Approach, NRSC/ISRO/Department of Space, 2023. March 2023, <https://ndma.gov.in/flood-hazard-atlases>.
- [28] IHCAP, Science brief: flood risk and early warning systems, *IHCAP* (2017) 16.
- [29] P. Rautela, Lack of scientific recordkeeping of disaster incidences: a big hurdle in disaster risk reduction in India, *Int. J. Disaster Risk Reduct.* 15 (2016) 73–79, <https://doi.org/10.1016/j.ijdrr.2015.12.005>.

- [30] A. Ogra, A. Donovan, G. Adamson, K.R. Viswanathan, M. Budimir, Exploring the gap between policy and action in Disaster Risk Reduction: A case study from India. *International Journal of Disaster Risk Reduction* 63 (2021) 102428, <https://doi.org/10.1016/j.ijdr.2021.102428>.
- [31] HPSDMA, *Activity Report 2011-2015*, Government of Himachal Pradesh, 2016.
- [32] UNDRR, Closing climate and disaster data gaps: new challenges, new thinking, united nations office for disaster risk reduction. <https://www.undrr.org/publication/closing-climate-and-disaster-data-gaps-new-challenges-new-thinking>, 2023.
- [33] ADPC-UNDRR, Disaster risk reduction in India, Asian Disaster Preparedness Center and UN Office for Disaster Risk Reduction (2020) 36. <https://www.preventionweb.net/publication/disaster-risk-reduction-india-status-report-2020-0>.
- [34] NDMA, DM (amendment) Bill. 12 December 2024. <https://x.com/ndmaindia/status/1867253828875346276>, 2024.
- [35] P. Adhikari, Y. Hong, K.R. Douglas, D.B. Kirschbaum, J. Gourley, R. Adler, G.R. Brakenridge, A digitized global flood inventory (1998-2008): compilation and preliminary results, *Nat. Hazards* 55 (2010) 405–422, <https://doi.org/10.1007/s11069-010-9537-2>.
- [36] L. Aceto, A.A. Pasqua, O. Petrucci, Effects of damaging hydrogeological events on people throughout 15 years in a Mediterranean region, *Advances in Geosciences* 44 (2017) 67–77, <https://doi.org/10.5194/adgeo-44-67-2017>.
- [37] M.C. Llasat, M. Llasat-Botija, O. Petrucci, A.A. Pasqua, J. Rossello, F. Vinet, L. Boissier, Floods in the north-western Mediterranean region: presentation of the HYMEX database and comparison with pre-existing global databases. *La Houille Blanche* 99 (1) (2013) 5–9, <https://doi.org/10.1051/lhb/2013001>.
- [38] *Census of India, Himachal Pradesh (Series 03, Part 12 A) District Census Handbook, Kullu, Village and Town Directory*, Directorate of Census Operations-Himachal Pradesh, 2011.
- [39] S. Singh, Destination development dilemma-case of Manali in Himachal Himalaya, *Tour. Manag.* 29 (2008) 1152–1156, <https://doi.org/10.1016/j.tourman.2008.02.018>.
- [40] J.S. Gardner, Natural Hazards Risk in the Kullu District, Himachal Pradesh, India. *Geographical Review* 92 (2) (2002) 282–306, <https://doi.org/10.1111/j.1931-0846.2002.tb00008.x>.
- [41] D.D.A.C. Asf, ALOS PALSAR Radiometric Terrain Corrected high res, includes material ©JAXA/METI 2007. <https://doi.org/10.5067/Z97HFCNKR6VA>, 2007.
- [42] Village Info. <https://villageinfo.in/himachal-pradesh/kullu.html>, 2021.
- [43] Punjab Government, Punjab District Gazetteers, Kangra District, Parts 2-4 Kulu and Saraj, Lahul and Spiti, Punjab Government, Lahore, 1918.
- [44] P. Chetwode, *Kulu: the End of the Habitable World*, John Murray, London, 1972.
- [45] A.F.P. Harcourt, *The Himalayan Districts of Kooloo, Lahoul and Spiti*, Vivek Publishing House, Delhi, 1871 (1972 republication).
- [46] M.P. Sah, R.K. Mazari, An overview of the geoenvironmental status of the Kullu Valley, Himachal Pradesh, India, *J. Mt. Sci.* 4 (1) (2007) 3–23, <https://doi.org/10.1007/S11629-007-0003-X>.
- [47] V. Gupta, B. Syed, A. Pathania, S. Raaj, A. Nanda, S. Awasthi, D.P. Shukla, Hydrometeorological analysis of July-2023 floods in Himachal Pradesh, India. *Natural Hazards* 120 (2024) 7549–7574, <https://doi.org/10.1007/s11069-024-06520-5>.
- [48] A. Diduck, R.M. Johnson, E. Edwards, A.J. Sinclair, J. Gardner, K. Patel, Small hydro and environmental justice: lessons from the Kullu District of Himachal Pradesh, in: A. Diduck, K. Patel, A.K. Malik (Eds.), *Advancing Environmental Justice for Marginalized Communities in India: Progress, Challenges and Opportunities*, Routledge, Abingdon, 2021. ISBN 9780367692810.
- [49] PWD, Public works department, information on national highways in Himachal Pradesh. <http://hppwd.hp.gov.in/information-on-national-highways-in-himachal-pradesh>, 2021.
- [50] P. Chauhan, Rohtang tunnel opens on Saturday, *Tribune* (2020), 2 October 2020, <https://www.tribuneindia.com/news/schools/rohtang-tunnel-opens-on-saturday-150012>.
- [51] A. Vashisht, Bhunter airport runway awaits extension, the *Tribune*, 22 August 2021, <https://www.tribuneindia.com/news/himachal/bhunter-airport-runway-awaits-extension-300430>, 2021.
- [52] *Tribune*, Aerial survey begins for Bilaspur-Leh railway line, The *Tribune* 19 September 2020, <https://www.tribuneindia.com/news/himachal/aerial-survey-begins-for-bilaspur-leh-railway-line-143057>, 2020.
- [53] J. Sood, P. Lynch, C. Anastasiado, Community non-participation in homestays in Kullu, Himachal Pradesh, India, *Tour. Manag.* 60 (2017) 332–347, <https://doi.org/10.1016/j.tourman.2016.12.007>.
- [54] IHCAP, *Climate Vulnerability, Hazards and Risk. An Integrated Pilot Study in Kullu District, Himachal Pradesh*. Synthesis Report, Indian Himalayas Climate Adaptation Programme (IHCAP), India, 2016.
- [55] P.G. Sarptodar, *Study on Early Warning Systems for Floods and Flash Floods: of the Most Vulnerable Communities of Beas River Basin in Selected 5 Villages of Kullu District, Kullu DDMA and UNDP*, 2016.
- [56] DDMA, *Kullu District disaster management plan*. District disaster management authority, Kullu (2017) 266. <https://hpkullu.nic.in/documents-2/>.
- [57] HPSDMA, Himachal Pradesh State Disaster Management Plan. Himachal Pradesh State Disaster Management Authority, Shimla (2017) 262. <https://www.hpsdma.nic.in//admnis/admin/showimg.aspx?ID=76>.
- [58] Pragyaa, Hazard, risk, vulnerability and capacity analysis (HRVCA) of selected 10 villages of Manali subdivision, in: District Kullu, Himachal Pradesh. Kullu DDMA and UNDP, 2016.
- [59] BSU and GBPNIHE, Accelerating change: engaging local communities in disaster risk reduction in the Indian Himalayan region, Science Policy Brief, Bath (UK) and Almora (India), June 2020 (2020), <https://doi.org/10.17870/bathspa.c.5036990.v1>.
- [60] R.M. Johnson, E. Edwards, J.S. Gardner, A.P. Diduck, Community vulnerability and resilience in disaster risk reduction: an example from Phoajal Nalla, Himachal Pradesh, India. *Regional Environmental Change* 18 (7) (2018) 2073–2087, <https://doi.org/10.1007/s10113-018-1326-6>.
- [61] *Civil and Military Gazette, The Kulu flood*. The *Civil and Military Gazette (Lahore)* 17 (5395) (1894) 4.
- [62] J.A. Ballesteros-Cánovas, D. Trappmann, M. Shekhar, A. Bhattacharyya, M. Stoffel, Regional flood-frequency reconstruction for Kullu district, western Indian Himalayas, *J. Hydrol.* 546 (2017) 140–149, <https://doi.org/10.1016/j.jhydrol.2016.12.059>.
- [63] S.K. Allen, J. Ballesteros-Cánovas, S.S. Randhawa, A.K. Singha, C. Huggel, M. Stoffel, Translating the concept of climate risk into an assessment framework to inform adaptation planning: insights from a pilot study of flood risk in Himachal Pradesh, Northern India, *Environ. Sci. Pol.* 87 (2018) 1–10, <https://doi.org/10.1016/j.envsci.2018.05.013>. September 2018.
- [64] IMD, *Climate Hazards and Vulnerability Atlas of India*, State: Himachal Pradesh. Office of Climate Research and Services, India Meteorological Department, 2022. Pune, <https://imd pune.gov.in/reports.php>.
- [65] M. Leal, C. Ramos, S. Pereira, Different types of flooding lead to different human and material damages: the case of the Lisbon Metropolitan Area. *Natural Hazards* 91 (2018) 735–758, <https://doi.org/10.1007/s11069-017-3153-3>.
- [66] M. Barnolas, M.C. Llasat, A flood geodatabase and its climatological applications: the case of Catalonia for the last century, *Nat. Hazards Earth Syst. Sci.* 7 (2) (2007) 271–281, <https://doi.org/10.5194/nhess-7-271-2007>.
- [67] M.C. Llasat, M. Llasat-Botija, L. López, A press database on natural risks and its application in the study of floods in Northeastern Spain. *Natural Hazards Earth System Sciences* 9 (2009) 2049–2061, <https://doi.org/10.5194/nhess-9-2049-2009>.
- [68] M. Barriandos, J.L. Ruiz-Bellet, J. Tuset, J. Mazón, J.C. Balasch, D. Pino, J.L. Ayala, The "Prediflood" database of historical floods in Catalonia (NE Iberian Peninsula) AD 1035–2013, and its potential applications in flood analysis, *Hydrol. Earth Syst. Sci.* 18 (2014) 4807–4823, <https://doi.org/10.5194/hess-18-4807-2014>.
- [69] M. Lang, D. Coeur, A. Audouard, M. Villanova-Oliver, J.P. Pene, BDHI: a French national database on historical floods, in: 3rd European Conference on Flood Risk Management (FLOODrisk 2016), Lyon, France, 2016, <https://doi.org/10.1051/e3sconf/20160704010>.
- [70] C. Copien, C. Frank, M. Becht, Natural hazards in the Bavarian Alps: a historical approach to risk assessment, *Nat. Hazards* 45 (2008) 173–181, <https://doi.org/10.1007/s11069-007-9166-6>.
- [71] B. Lasteria, M.R. Simonetti, M. Casaioli, S. Mariani, G. Monacelli, Socio-economic impacts of major floods in Italy from 1951 to 2003. *Advances in Geosciences* 7 (2006) 223–229, <https://doi.org/10.5194/adgeo-7-223-2006>.

- [72] O. Petrucci, P. Salvati, L. Aceto, C. Bianchi, A.A. Pasqua, M. Rossi, F. Guzzetti, The Vulnerability of People to Damaging Hydrogeological Events in the Calabria Region (Southern Italy). *International Journal of Environmental Research and Public Health* 15 (1) (2018) 48, <https://doi.org/10.3390/ijerph15010048>.
- [73] N. Hilker, A. Badoux, C. Hegg, The Swiss flood and landslide damage database 1972–2007. *Natural Hazards Earth System Sciences* 9 (2009) 913–925, <https://doi.org/10.5194/nhess-9-913-2009>.
- [74] M. Diakakis, G. Deligiannakis, Flood fatalities in Greece: 1970–2010. *Journal of Flood Risk Management* 10 (2017) 115–123, <https://doi.org/10.1111/jfr3.12166>.
- [75] M.C. Llasat, M. Llasat-Botija, O. Petrucci, A.A. Pasqua, J. Rosselló, F. Vinet, L. Boissier, Towards a database on societal impact of Mediterranean floods within the framework of the HYMEX project. *Natural Hazards Earth System Sciences* 13 (2013) 1337–1350, <https://doi.org/10.5194/nhess-13-1337-2013>.
- [76] J.M. Jeffers, Environmental knowledge and human experience: using a historical analysis of flooding in Ireland to challenge contemporary risk narratives and develop creative policy alternatives, *Environ. Hazards* 13 (3) (2014) 229–247, <https://doi.org/10.1080/17477891.2014.902800>.
- [77] O. Singh, M. Kumar, Flood events, fatalities and damages in India from 1978 to 2006, *Nat. Hazards* 69 (2013) 1815–1834, <https://doi.org/10.1007/s11069-013-0781-0>.
- [78] V.S. Kale, Is flooding in South Asia getting worse and more frequent? *Singapore Journal of Tropical Geography* 35 (2014) 161–178, <https://doi.org/10.1111/sjtg.12060>.
- [79] S.C. Bhan, A.K. Devrani, V. Sinha, An analysis of monthly rainfall and the meteorological conditions associated with cloudburst over the dry region of Leh (Ladakh), India, *Mausam* 66 (1) (2015) 107–122, <https://doi.org/10.54302/mausam.v66i1.371>.
- [80] V. Gupta, D.P. Dobhal, S.C. Vaideswaran, August 2012 cloudburst and subsequent flash flood in the Asi Ganga, a tributary of the Bhagirathi river, Garhwal Himalaya, India, *Curr. Sci.* 105 (2) (2013) 249–253, <https://www.currentscience.ac.in/Volumes/105/02/0249.pdf>.
- [81] S. Jangra, M. Singh, Analysis of rainfall and temperatures for climatic trend in Kullu Valley, *Mausam* 62 (1) (2011) 77–84, <https://doi.org/10.54302/mausam.v62i1.207>.
- [82] NDMA, National disaster management authority guidelines, management of Glacial Lake Outburst floods (GLOFs). National Disaster Management Authority, Government of India, New Delhi, 2020. <https://www.ndma.gov.in/sites/default/files/PDF/Guidelines/Guidelines-on-Management-of-GLOFs.pdf>.
- [83] M.P. Sah, R.K. Mazari, Anthropogenically accelerated mass movement, Kulu Valley, Himachal Pradesh, India, *Geomorphology* 26 (1998) 123–138, [https://doi.org/10.1016/S0169-555X\(98\)00054-3](https://doi.org/10.1016/S0169-555X(98)00054-3).
- [84] F. Berkes, J.S. Gardner, A.J. Sinclair, Comparative aspects of mountain land resources management and sustainability: case studies from India and Canada, *Int. J. Sustain. Dev. World Ecol.* 7 (2000) 375–390, <https://doi.org/10.1080/13504500009470056>.
- [85] E. Sana, A. Kumar, E. Robson, R. Prasanna, U. Kala, D.G. Toll, Preliminary assessment of series of landslides and related damage by heavy rainfall in Himachal Pradesh, India, during July 2023, *Landslides* 21 (2024) 919–931, <https://doi.org/10.1007/s10346-023-02209-1>.
- [86] HPSDMA, in: Proceedings of the 23rd Meeting of the HP State Executive Committee (SEC) Constituted under the Disaster Management Act (2005), 2024. <https://hpsdma.nic.in/WriteReadData/LINKS/DF93e78c7d-83ac-4d53-a6c8-11c880c1b4ff.pdf>.
- [87] K. Chand, A.K. Jamwal, G. Meraj, T. Thakur, M. Farooq, P. Kumar, S.K. Singh, S. Kanga, J. Debnath, Integrating geoenvironmental and socioenvironmental analyses for flood vulnerability assessment in the Kullu Valley, Himachal Pradesh, India, *Int. J. Disaster Risk Reduct.* 108 (2024) 104494, <https://doi.org/10.1016/j.ijdr.2024.104494>.
- [88] NDMA, National disaster management plan 2019. National Disaster Management Authority, Government of India, New Delhi, 2019. <https://ndma.gov.in/sites/default/files/PDF/ndmp-2019.pdf>.
- [89] UNDRR, Words into Action guidelines: implementation guide for addressing water-related disasters and transboundary cooperation. <https://www.undrr.org/publication/words-action-guidelines-implementation-guide-addressing-water-related-disasters-and>, 2018.
- [90] UNDRR and WMO, Global status of multi-hazard early warning systems. <https://www.undrr.org/reports/global-status-MHEWS-2023>, 2023.
- [91] V. Gupta, M.P. Sah, Impact of the Trans-Himalayan Landslide Lake Outburst Flood (LLOF) in the Satluj catchment, Himachal Pradesh, India. *Natural Hazards* 45 (2008) 379–390, <https://doi.org/10.1007/s11069-007-9174-6>.
- [92] G.C.D. Adamson, Institutional and community adaptation from the archives: a study of drought in western India, 1790–1860, *Geoforum* 55 (2014) 110–119, <https://doi.org/10.1016/j.geoforum.2014.05.010>.
- [93] J.M. Jeffers, Particularizing adaptation to non-predominant hazards: a history of wildfires in County Donegal, Ireland from 1903 to 2019, *Int. J. Disaster Risk Reduct.* 58 (2021) 102211, <https://doi.org/10.1016/j.ijdr.2021.102211>.
- [94] S.K. Shah, A. Bhattacharyya, M. Shekhar, Reconstructing discharge of Beas River basin, Kullu valley, western Himalaya, based on tree-ring data, *Quat. Int.* (286) (2013) 138–147, <https://doi.org/10.1016/j.quaint.2012.09.029>.
- [95] V.S. Kale, Palaeoflood Hydrology in the Indian Context. *Journal Geological Society of India* 71 (2008) 56–66. <https://www.geosocindia.com/index.php/jgsi/article/view/80874>.
- [96] D.E.J. Hobley, H.D. Sinclair, S.M. Mudd, Reconstruction of a major storm event from its geomorphic signature: The Ladakh floods, 6 August 2010. *Geology* 40 (6) (2012) 483–486, <https://doi.org/10.1130/G32935.1>.
- [97] R.M. Johnson, E. Edwards, J.S. Gardner, B. Mohan, Village heritage and resilience in damaging floods and debris flows, Kullu valley, Indian Himalaya, in: I. Convery, G. Corsane, P. Davis (Eds.), *Displaced Heritage: Responses to Disaster, Trauma and Loss*, Boydell & Brewer, Woodbridge, 2014, pp. 207–224. <https://boydellandbrewer.com/9781783274307/displaced-heritage/>.
- [98] CWC, Flood damage Statistics (statewise and for the country as a whole) during 1953 to 2020. Central Water Commission, Government of India, 2022. <https://cwc.gov.in/sites/default/files/flood-damage-data-merged.pdf>.
- [99] D.E. Alexander, Social media in disaster risk reduction and Crisis management, *Sci. Eng. Ethics* 20 (2014) 717–733, <https://doi.org/10.1007/s11948-013-9502-z>.
- [100] A. Yzaguirre, M. Smit, R. Warren, Newspaper archives + text mining = rich sources of historical geo-spatial data, *IOP Conf. Ser. Earth Environ. Sci.* 34 (2016) 012043, <https://doi.org/10.1088/1755-1315/34/1/012043>.
- [101] R.M. Koll, How India Can Tackle its Climate Challenges, *Hindustan Times*, 2024. <https://www.hindustantimes.com/opinion/how-india-can-tackle-its-climate-challenges-101724342256374.html>.