



UNIVERSITY OF
GLOUCESTERSHIRE

This is a peer-reviewed, final published version of the following document and is licensed under Creative Commons: Attribution 2.0 license:

Lai, C Benjamin, Zhang, Ying, Rogers, Sally L and Mager, Dixie L (2009) Creation of the two isoforms of rodent NKG2D was driven by a B1 retrotransposon insertion. Nucleic Acids Research, 37 (9). pp. 3032-43. doi:10.1093/nar/gkp174

Official URL: <http://nar.oxfordjournals.org/content/37/9/3032.long>

DOI: <http://dx.doi.org/10.1093/nar/gkp174>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/2001>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

This is the final published version and is licensed under a [Creative Commons Attribution 2.0 License](#):

Lai, C Benjamin and Zhang, Ying and Rogers, Sally L and Mager, Dixie L (2009). Creation of the two isoforms of rodent NKG2D was driven by a B1 retrotransposon insertion. Nucleic acids research, 37 (9), 3032-3043.

Published in Nucleic acids research, and available online at:

<http://nar.oxfordjournals.org/content/37/9/3032.long>

We recommend you cite the published version.

The URL for the published version is <http://dx.doi.org/10.1093/nar/gkp174>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

Creation of the two isoforms of rodent NKG2D was driven by a B1 retrotransposon insertion

C. Benjamin Lai, Ying Zhang, Sally L. Rogers and Dixie L. Mager*

Terry Fox Laboratory, British Columbia Cancer Agency and Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada

Received September 17, 2008; Revised February 26, 2009; Accepted March 3, 2009

ABSTRACT

The mouse gene for the natural killer (NK) cell-activating receptor *Nkg2d* produces two protein isoforms, NKG2D-S and NKG2D-L, which differ by 13 amino acids at the N-terminus and have different signalling capabilities. These two isoforms are produced through differential splicing, but their regulation has not been investigated. In this study, we show that rat *Nkg2d* has the same splicing pattern as that of the mouse, and we mapped transcriptional start sites in both species. We found that the splice forms arise from alternative promoters and that the NKG2D-L promoter is derived from a rodent B1 retrotransposon that inserted before mouse–rat divergence. This B1 insertion is associated with loss of a nearby splice acceptor site that subsequently allowed creation of the short NKG2D isoform found in mouse but not human. Transient reporter assays indicate that the B1 element is a strong promoter with no inherent lymphoid tissue-specificity. We have also identified different binding sites for the ETS family member GABP within both the mouse and rat B1 elements that are necessary for high-promoter activity and for full *Nkg2d-L* expression. These findings demonstrate that a retroelement insertion has led to gene-regulatory change and functional diversification of rodent NKG2D.

INTRODUCTION

The activity of natural killer (NK) cells against potential target cells is regulated by a balance of inhibitory and activating receptors (1–3). One of the most well-studied activating receptors is NKG2D, which recognizes ligands often upregulated in response to stress, such as during virus infection or malignant transformation (4). A recent

study using NKG2D deficient mice has demonstrated that expression of this receptor inhibits some types of cancer in spontaneous tumour models (5), indicating a significant role for NKG2D in tumour surveillance. The importance of NKG2D is also indicated by the existence of specific viral evasion mechanisms directed against it (6).

In the mouse, NKG2D is expressed on NK cells, activated CD8⁺ T cells, NKT cells, $\gamma\delta$ T cells and activated macrophages (7). Alternative splicing produces two different isoforms of NKG2D in mouse, one of which contains 13 extra amino acids at its N terminus (NKG2D-L) compared to the shorter isoform (NKG2D-S) (8). NKG2D-S can bind and signal through the adapters DAP10 and DAP12, while NKG2D-L can only bind DAP10 (8). Due to differential downstream pathways, DAP12 is capable of initiating cytotoxic as well as cytokine release responses while DAP10 can only activate an NK cell to kill (9). CD8⁺ T cells express only DAP10, whereas NK cells express both DAP10 and DAP12 (8). Resting mouse NK cells express very little NKG2D-S, but expression of this isoform is induced upon NK cell activation, enabling signalling through NKG2D-L/DAP10, NKG2D-S/DAP10 and NKG2D-S/DAP12 (8). These expression patterns of DAP10/12 and of mouse NKG2D isoforms result in functional diversification of this receptor. In mouse NK cells, NKG2D is strongly activatory, stimulating NK cells to attack bound targets. However, in CD8⁺ T cells, NKG2D serves only a co-stimulatory function (10). The situation in human is slightly different. The single human NKG2D isoform resembles mouse NKG2D-L and also only binds DAP10 (11–13). Not surprisingly, NKG2D stimulation using monoclonal antibodies in human NK cells initiates a cytotoxic response but not cytokine release (13). As in the mouse, NKG2D/DAP10 only acts in a co-stimulatory manner in human CD8⁺ T cells (14).

While many studies have concentrated on the function of NKG2D, surprisingly little is known about its transcriptional regulation. Here we show that the spliced isoforms of mouse NKG2D arise from the use of

*To whom correspondence should be addressed. Tel: +1 604 675 8139; Fax: +1 604 877 0712; Email: dmager@bccrc.ca

alternative promoters. We focus on the promoter for mouse NKG2D-L and show it to be derived from a rodent-specific B1 retrotransposon with a functional binding site for GABP, a member of the ETS family. We propose that insertion of this retrotransposon, through unusual donation of a polymerase II promoter, led to the alternative splicing patterns maintained in both mouse and rat that, in turn, result in functionally distinct NKG2D isoforms. These results document an intriguing example of how retrotransposons can affect the evolution of host gene expression and, more specifically, lend insight into the transcriptional regulation of NKG2D.

MATERIALS AND METHODS

Cell culture

LNK (mouse NK cell line), EL4 (mouse T cell line) and RNK16 (rat NK cell line) were cultured in RPMI 1640 medium supplemented with 10% (v/v) fetal bovine serum, 100 U/ml penicillin, 100 U/ml streptomycin, 50 μ M 2-ME and 2 mM L-glutamine. LNK cells were further supplemented with 1000 U/ml IL-2 (Peprotech). NIH 3T3 (mouse embryonic fibroblast cell line) was cultured in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum.

Mice

C57BL/6 (B6) mice were purchased from The Jackson Laboratory and bred in our animal facility. The use of animals for this study was approved by the Animal Care Committee of the University of British Columbia, and animals were maintained in accordance with the guidelines of the Canadian Council on Animal Care.

Reverse-transcriptase polymerase chain (RT-PCR) reaction and sequencing

All RT-PCR was performed using Superscript III (Invitrogen) using random primers according to manufacturer's instructions. For Sprague-Dawley rat total RNA (Clontech), PCR was then carried out using gene-specific primers: rat exon 1a forward, 5'-GATTCACAAGAAACAGGACCTC; rat exon 1b forward, 5'-TGGCATGGGTCGTGATC; rat exon 4 reverse, 5'-CAACAAGGACTCGAACAACG. PCR products were cloned into pGEMT vector (Promega) according to manufacturer's instructions. Sequencing was performed using the T7 primer by McGill University and Genome Québec Innovation Centre Sequencing Platform. For LNK, RNK16, EL4 and NIH 3T3 cell lines, total RNA was collected using Rneasy minikit and QIAshredder homogenizer (QIAGEN). The RNA samples were treated with DNase I twice: during purification using the on-column protocol of the Rneasy minikit and after elution using DNase I from Invitrogen. PCR primers used were as follows: mouse exon 1a forward, 5'-GAAACAGGATCTCCCTTCTCT; mouse exon 1b forward, 5'-ACAACCTGGATCAGTTTCTGAA; mouse exon 6 reverse, 5'-GGTTCCAGGTTTCTCTTCATT; rat exon 1a forward, 5'-GATTCAC

AAGAAACAGGACCTC; rat exon 1b forward, 5'-TGGCATGGGTCGTGATC; rat exon 6 reverse, 5'-GGTTCAGGCTTTGTTCTCA; GAPDH forward, 5'-GTGGA GTCTACTGGTGTCTTC; GAPDH reverse, 5'-GTGGCAGTGATGGCATGGAC.

Sequence alignment

Nucleotides positioned from -75 to +75 from the splice acceptor site of human *NKG2D* exon 2 were used as the reference sequence in UCSC Genome browser (<http://genome.ucsc.edu>). From the conservation tract, the orthologous sequences of rhesus, mouse, rat, guinea pig, hedgehog, horse, cow and armadillo were obtained. These sequences were aligned using Multiple Alignment using Fast Fourier Transform and G-INS-i strategy (MAFFT, <http://align.bmr.kyushu-u.ac.jp/mafft/online/server/index.html>). The same was done for -50 to +50 of the ATG in human exon 3 and for aligning PB1D9 with B1Mur3.

5'RACE

C57BL/6 mouse spleens were homogenized. Total RNA was extracted from LNK cells and mouse splenocytes using Rneasy minikit and QIAshredder homogenizer (QIAGEN). Sprague-Dawley rat total RNA was purchased from Clontech. 5'RACE was performed using FirstChoice RLM-RACE kit (Ambion) according to manufacturer's instructions. The primers used were as follows: mouse exon 6 reverse flanking, 5'-GGTTCCAGGTTTCTCTTCATT; mouse exon 6 reverse nested, 5'-TTGTTCTGTGACATATCCAGTT; rat reverse exon 6 flanking, 5'-GGTTCCAGGCTTTGTTCTCA; rat reverse exon 4 nested, 5'-CAACAAGGACTCGAACAACG.

Generation of luciferase reporter constructs

Promoter fragments were generated by PCR using forward primers with KpnI site inserted at the 5' end and reverse primers with NheI site inserted at the 5' end. PCR products were digested with KpnI/NheI (NEB) and cloned into pLG4.10 firefly luciferase promoter vector (Promega). The only exception is the generation of a construct containing the mouse B1 and a 391-bp fragment at separate sites. For this construct, the mouse B1 was cloned into the multiple cloning region using KpnI/NheI while the 391 bp fragment was inserted into a BamHI site downstream of the luciferase reporter gene. The primers used are as follows: mouse B1 forward: 5'-AAGGTACCCGGAAGTGGTGTCACATA; mouse B1 reverse: 5'-AAGCTAGCTTTTTTAAGACAGTGTATC; mouse -391 of B1 forward: 5'-AAGGTACCCTGACATTTATTTATCTT; mouse -1 of B1 reverse: 5'-AAGCTAGCATCTGTCTTAATTTCTG; mouse 391-bp BamHI forward: 5'-AAGGATCCCTGACATTTATTTATCTT; mouse 391-bp BamHI reverse: 5'-AAGGATCCCAATCTGTCCTAATTTCTG; rat B1 forward: 5'-AAGGTACC AACCAGAAGCAGTGTACAC; rat B1 reverse: 5'-AAGCTAGCTTTCTAAAGACAGTTTGTCTCTCT; B1 3' 32-bp truncation reverse: 5'-AAGCTAGCGGCTACCTCAGATTTACAG; B1 3' 59-bp truncation reverse: 5'-AAGCTAGCTGCCTCTGCTTCCTGAATG. The PB1D9 fragment truncated 20 bp at the 5' end and 59 bp at the

3' end was cloned by annealing complementary oligonucleotides, digesting with KpnI/NheI, and ligating into pGL4.10. The oligonucleotide sequences were as follows: 5'-AAGGTACCTTTAATCTCAGCATTTCAGG AAGCAGAGGCAGCTAGCAA and 5'-TTGCTAGCT GCCTCTGCTTCCTGAATGCTGAGATTAAAGGTA CCTT. One microgram of each oligonucleotide was mixed in 20 µl of water and heated to 95°C and then allowed to cool to room temperature. Primers used to clone GABP and AP1 mutants were as follows: GABP mutant forward: 5'-AAGGTACCCGTTAGTGGTGTACATA (mutation is underlined); AP1 mutant forward: 5'-AAG GTACCCGGAAGTGGTGGCACATA; double mutant forward: 5'-AAGGTACCCGTTAGTGGTGGCACATA.

Cell transfection and luciferase assays

LNK, EL4 and NIH 3T3 cells were transfected using Lipofectamine 2000 (Invitrogen) according to manufacturer's instructions. Briefly, 2.5×10^6 LNK and EL4 cells and 6×10^5 NIH 3T3 cells were seeded into 500 µl of growth media in 24-well plates for 24 h at 37°C. For each well, transfection media was prepared: 2 µl of Lipofectamine 2000 reagent was mixed with 1 µg of DNA for LNK and EL4 while 0.5 µl of Lipofectamine 2000 reagent was mixed with 0.6 µg of DNA for NIH 3T3. Transfection media was brought up to 100 µl using unsupplemented media and incubated at 37°C for 20 min. Next, growth media in each well was exchanged with 150 µl of unsupplemented media and 100 µl of transfection media. Cells were incubated at 37°C for 6 h before being topped up with 250 µl of media supplemented with 20% fetal bovine serum or fetal calf serum and IL-2 where necessary. Cells were then incubated at 37°C for 16 h before assaying.

Luciferase activity was assayed using the Dual-Luciferase Reporter Assay System (Promega) according to the manufacturer's instructions. Firefly luciferase activity was normalized relative to Renilla luciferase activity for each transfection and calculated as fold increase over pGL4.10-BASIC (pGL4B).

Electrophoretic mobility shift assays (EMSAs)

Double stranded probes were generated by annealing the following oligomers to their complementary oligomers: mouse wild type: 5'-AGGACAGATTCGGAAGT GGTGTACATATCTTTA; mouse GABP mutant: 5'-A GGACAGATTCGTTAGTGGTGTACATATCTTTA; mouse AP1 mutant: 5'-AGGACAGATTCGGAAGTGG TGGCACATATCTTTA; mouse double mutant: 5'-A GGACAGATTCGTTAGTGGTGGCACATATCTTTA; rat GABP1 wild type: 5'-GAGCCTGAACCAGAAGC AGTGTACCTA; rat GABP1 mutant: 5'-GAGCCTG AACCATAGCAGTGTACCTA; rat GABP2 wild type: 5'-TGGTCTATACAGGAAGTTCTAAGATATC; rat GABP2 mutant: 5'-TGGTCTATACAGTTAGTT CTAAGATATC. Nuclear extraction, probe labelling and the gel-shift assay were performed as described before (15). Protein concentration was determined using the Qubit Quantitation Platform (Invitrogen).

GABPα antibody (H-2 X) and c-Jun antibody (G-4 X) were purchased from Santa Cruz Biotechnology.

siRNA and real-time RT-PCR

LNK cells (5×10^4 per well) were seeded in 96-well plates and grown overnight in 100 µl serum-free RPMI 1640 medium supplemented with 1000 U/ml IL-2. The next day, the growth media was removed, and cells were transfected with 1 µM Accell SMARTpool mouse GABPA siRNA or Accell Non-Targeting Pool siRNA using 100 µl Accell siRNA Delivery media (Dharmacon) supplemented with 1000 U/ml IL-2. An untreated sample was also included where the media was exchanged, but no siRNA was added. Seventy-two hours later, total RNA collection and RT-PCR were carried out as described above. Three wells were pooled per treatment. The resultant cDNA was used as template in real-time PCR using the 7500 Fast Real-Time PCR System (Applied Biosystems). Relative expression was determined using the $2^{-\Delta\Delta CT}$ method. Threshold cycles for *Gabp*, *Nkg2d-S* and *Nkg2d-L* were normalized to *Gapdh*. Primers used were GAPDH forward: 5'-GACTTCAAC AGCAACTCCCAC; GAPDH reverse: 5'-TCCACCAC CCTGTTGCTGTA; NKG2D-S forward: 5'-GAAGGC TTTGACTCACAAGAAACAGG; NKG2D-L forward: 5'-TGGCATTGATTCTGATCGAAAGTCT; NKG2D reverse: 5'-CTCCAGGTTGACTGGTAGTTAGTGC.

RESULTS

A rodent SINE element is linked to alternative exon 1 usage

To begin to dissect the transcriptional regulation of mouse NKG2D, we first examined the mouse *Nkg2d* sequence via the UCSC Genome browser (<http://genome.ucsc.edu>) and noticed a repetitive B1 element, termed PB1D9, inserted at the 3' end of intron 1. An orthologous SINE element is found in the rat (termed B1Mur3) but not in the human, *NKG2D* gene. The two B1 elements differ by a 5' 3-bp deletion in the mouse B1 and an internal 30-bp insertion in the rat B1, but are otherwise 84.4% identical (aligned using MAFFT, <http://align.bmr.kyushu-u.ac.jp/mafft/online/server/index.html>). Although annotated differently in mouse and rat by Repbase (<http://www.girinst.org/rebase/index.html>), these B1 elements represent the same ancestral insertion. To avoid confusion, we will refer to the mouse element as mB1 and the rat element as rB1. While there are two mouse NKG2D isoforms arising from transcripts that alternatively encode exon 1a or 1b, there is only one human NKG2D protein from transcripts that always include exon 2, which is homologous to mouse exon 1b (Figure 1A). The situation in rat is less clear. A rat mRNA initiating near the region homologous to mouse exon 1b has been previously described (16). However, a rat genomic sequence that is highly similar to mouse exon 1a is found upstream of this initiation exon (Figure 1A). To see if rat *Nkg2d* splicing patterns resemble those of the mouse, we undertook RT-PCR on Sprague-Dawley rat splenic total RNA. Primers were designed in rat sequences orthologous to mouse exons 1a and 4. Sequencing of the amplified

product shows that a rat transcript indeed includes the region orthologous to mouse exon 1a and splices out the previously reported exon (Figure 1B). Furthermore, the splice sites used are the same as that of the mouse. We herein designate the exon found by RT-PCR as rat exon 1a and the alternative exon as rat exon 1b. As was done with the mouse (8), we also designate the isoform arising from exon 1a as rNKG2D-S and the isoform arising from exon 1b as rNKG2D-L. These results suggest that the *Nkg2d* splicing pattern in rat is identical to that of the mouse and that both species harbour a SINE element at the 3' end of intron 1. In comparison, human does not have the SINE element and splicing of *NKG2D* always gives rise to one protein.

The SINE element has polymerase II promoter potential

To determine if the SINE element is contributing an alternative promoter element, we first looked for transcription start sites (TSS) for mouse and rat *NKG2D-L* transcripts using 5' rapid amplification of cDNA ends (5'RACE). Primers were designed in mouse exon 6 and rat exons 4 and 6. We found two start sites within the mB1 element using C57BL/6 splenocytes (Figure 2A). Two additional sites were found using LNK, a mouse NK cell line that expresses both *NKG2D* isoforms (Figure 2B). Surprisingly, while one was located in the SINE element, the other lies downstream of the ATG1 start codon. Theoretically, this transcript could still be translated into *NKG2D-S* using the ATG2 start codon in exon 2. Further experiments are needed to verify this possibility. With RNA from Sprague-Dawley rat spleen, we found a dispersed pattern of TSS (Figure 2C). Two start sites were within the rB1 element while seven were downstream of the SINE but upstream of ATG1.

The 5'RACE experiments suggest that the rodent SINE element contains a functional promoter. To examine this possibility, we tested the mouse and rat SINE elements and the upstream region in luciferase reporter assays. Various fragments including and excluding mB1 and rB1 were cloned into pGL4B. These constructs were transiently transfected into LNK cells and promoter activity was measured. As shown in Figure 2D, both mB1 and rB1 elements by themselves display promoter activity. A fragment 391-bp upstream of mB1 shows no appreciable activity. Paradoxically, when this upstream region combined with mB1 was tested, the promoter activity was abolished. However, if this fragment was separated from mB1 in the same construct, only a small decrease in promoter activity was observed compared to a construct with mB1 alone. These results demonstrate that transcription not only starts within the SINE elements, but also that they contain polymerase II promoters.

The mouse B1 element contributes a functional GABP-binding site

Transposable elements harbouring functional enhancer sites are not uncommon (17). Thus, we interrogated the mouse mB1 sequence for potential transcription factor-binding sites using TESS (<http://www.cbil.upenn.edu/cgi-bin/teess/teess>). Curated results containing only




Figure 1. *NKG2D* genomic organization and splicing pattern. (A) A comparison of *NKG2D* exon-intron organization and splice variants in human, mouse and rat. Human exon nomenclature was revised based on 5'RACE from another study (13). Exons are labelled with roman numerals. Translated regions are shown as thicker solid boxes. Open boxes indicate the B1 element present in mouse and rat. Rat exon 1a was previously uncharacterized, and hence the first intron is shown as dotted line. (B) RT-PCR detection of an *Nkg2d-S* transcript in rat. Total RNA from Sprague-Dawley rat was used for reverse transcriptase reactions. Primers used for PCR are marked by arrows. PCR product was sequenced and shown as spliced transcript in the diagram.

mouse-specific, hematopoietic or lymphocyte-related transcription factors are shown in Figure 3A. To determine which of these sites contribute functional regulatory elements, luciferase reporter assays were carried out using truncated fragments of mB1 in pGL4B, transiently transfected in LNK cells. Surprisingly, fragments that truncate the 3' end of mB1 display much higher activity than the entire 109-bp mB1 (Figures 2D and 3B). A 77-bp fragment that eliminates the predicted TCF1, CEBP β and YY1-binding sites at the 3' end shows 40-fold activity over basal. Of note, this piece removes an ATG at the +78 position. When the 77-bp segment was further 3' truncated to 50 bp, eliminating two more TCF1 and one GATA3 binding site, 97-fold activity over basal was observed. Luciferase activity only dropped to 3-fold over basal when 20 bp were deleted at the 5' end of the 50-bp segment. This construct deletes a GABP, an AP1 and a GATA3-binding site. These results suggest that the essential transcription factor-binding sites are within the first 20 bp of mB1.

To further narrow down the essential transcription factor-binding sites, we mutated the potential GABP and AP1-binding sites and repeated the reporter assays in LNK cells. We chose to mutate the 77-bp construct, because the ATG at position +78 may be interfering with luciferase expression. Promoter activity does not




Figure 2. Detection of polymerase II promoter in SINE elements. (A) RLM-5'RACE of mouse *Nkg2d*. Sequence shown corresponds to chr6:129 572 548–29 572 772 of the mouse genome from UCSC

decrease appreciably using an AP1 mutant construct (Figure 3B). On the other hand, the GABP mutant and double mutant only show 2- and 4-fold activity over basal, respectively. These results indicate that the GABP-binding site is essential to enhancing transcription from the mB1 polymerase II promoter.

GABP binds the identified mB1 enhancer site *in vitro*

The core GGAA nucleotides in the binding sites of ETS family members, of which GABP is a member, are highly conserved. To determine if GABP is indeed binding to this mB1 element, EMSA were performed. When a DNA oligomer spanning the 5' end of mB1 was incubated with LNK nuclear extracts, two complexes are clearly visible in EMSAs, labelled I and II (Figure 4). Complex I contains GABP, since it can be supershifted by anti-GABP α antibody. This complex is decreased or abolished using wild-type unlabelled oligomers in a dose-dependent manner (10-, 25-, 50- and 100-fold excess). Complex I is still clearly visible when unlabelled oligomers containing a mutated GABP site or with mutations in both the GABP and AP1 sites were used as competitors. The fact that these signals are lower compared to complex I without competitors is likely the result of non-specific competition at 100-fold excess. In agreement with the previous experiments, the oligomer did not seem to bind AP1. A competition assay using unlabelled oligomers with a mutated AP1 site did not produce any complexes. In addition, anti-AP1 antibody did not induce a supershift. The identity of complex II is currently unknown. It is possible that this complex contains GATA3, since the oligomer spans a GATA3-binding site. Furthermore, all mutant competitors were equally effective at abolishing this complex and neither anti-GABP nor anti-AP1 antibodies supershifted the complex. Taken together, these results suggest that GABP, but not AP1, is an important factor in transcription from mB1.

GABP binds a separate site in rat B1 *in vitro*

We wondered if GABP regulation of the B1 promoter is conserved between mouse and rat. An alignment of mB1

Genome Browser (mm9 assembly). Underlined sequence represents the mB1 element inserted in front of exon 1b. Single stars mark transcription start sites found in C57BL/6 mouse spleen. Stacked double stars mark transcription start sites found in LNK mouse cell line. ATG1 start codon for mNKG2D-L is in brackets. Splice donor site of exon 1b is denoted with an arrow. (B) Detection of *Nkg2d* splice forms in LNK, RNK16, EL4 and NIH 3T3 cell lines with RT-PCR. The mouse exon-intron structure of *Nkg2d* is shown with exons labelled in roman numerals. Primers used are shown as arrowheads. (C) 5'RACE of rat *Nkg2d*. Sequence shown corresponds to chr4:166 923 873–166 924 142 of the rat genome from UCSC Genome Browser (rn4 assembly). Single stars mark transcription start sites found in Sprague-Dawley rat spleen. Underlined sequence represents the rB1 element. ATG1 start codon for rNKG2D-L is in brackets. Splice donor site of exon 1b is denoted with an arrow. (D) Luciferase reporter assay of SINE elements. Fragments including the rodent SINEs and 391-bp upstream of the mouse SINE were cloned into pGL4B and assayed using a dual-luciferase system. Promoter activity was normalized to co-transfected Renilla luciferase activity and calculated as fold above pGL4B. Data indicate mean (\pm SD) of three independent experiments.






Figure 3. Identification of transcription factor binding sites that contribute to *Nkg2d-L* regulation. (A) TESS search for transcription factor binding sites in mB1. The sequence of the mB1 inserted in front of mouse *Nkg2d* exon 1b is shown. Potential transcription factor binding sites are indicated with brackets. Only mouse-specific, hematopoietic or lymphocyte-related factors are shown. (B) Luciferase reporter assay of truncated and mutated SINE elements. mB1 sequence was truncated at the 5' and 3' ends or mutated at specific transcription factor binding sites, cloned into pGL4B and assayed using a dual-luciferase system. Mutated sites are indicated with an X. Promoter activities were normalized to co-transfected Renilla luciferase activity and calculated as fold above pGL4B. Data indicate mean (\pm SD) of two independent experiments.

WT AGGACAGATTCGGAAGTGGTGTACATATCTTTA
 GABP mutant AGGACAGATTCGTTAGTGGTGTACATATCTTTA
 API mutant AGGACAGATTCGGAAGTGGTGGCACATATCTTTA
 Double mutant AGGACAGATTCGTTAGTGGTGGCACATATCTTTA

LNK nuclear extract	-	+	+	+	+	+	+	+	+	+	+
WT competitor	-	-	+	+	+	+	-	-	-	-	-
GABP mutant competitor	-	-	-	-	-	-	+	-	-	-	-
API mutant competitor	-	-	-	-	-	-	-	+	-	-	-
Double mutant competitor	-	-	-	-	-	-	-	-	+	-	-
Anti-GABPb	-	-	-	-	-	-	-	-	-	+	-
Anti-c-Jun	-	-	-	-	-	-	-	-	-	-	+

Complex I ▶
 Complex II ▶




Figure 4. GABP forms protein-DNA complex within mB1 in EMSA. Sequences of probes and competitors used are shown at the top. Mutations are underlined. As control, lane 1 contains only γ^{32} P-labelled WT probe. LNK nuclear extract was incubated with labelled WT probe at 4°C for 20 min (lane 2). For competition assays, unlabelled competitors were incubated with LNK nuclear extract at 4°C for 20 min before addition of labelled WT probe for another 20-min incubation. WT competitors were used at 10-, 25-, 50- and 100-fold molar excess (lanes 3–6). Competitors with GABP, API or both binding sites mutated were used at 100-fold molar excess (lanes 7–9). Supershift assays were carried out using 5 μg of anti-GABPα (lane 10) or anti-c-Jun (lane 11) antibodies.

with rB1 shows that rB1 has an AGAA sequence instead of the core GGAA sequence at the mB1 GABP-binding site (Figure 5A). However, a separate potential GABP-binding site is found in the internal insertion of rB1 using TESS (<http://www.cbil.upenn.edu/cgi-bin/teess/teess>). These sites in rB1 were labelled GABP1 and GABP2, respectively. To see if either of these sites bind rat GABP, we subjected them to EMSA. DNA oligomers containing GABP2, but not GABP1, formed DNA-protein interactions with RNK16 (an *Nkg2d* expressing rat NK cell line, Figure 2B) nuclear extract (Figure 5B). The identity of the protein in this sole complex, labelled III, was confirmed to be GABP in a supershift assay using an anti-GABPα antibody. Complex III can be abolished using wild-type unlabelled oligomers at 100-fold excess but not with oligomers where the GABP2 site is mutated. These results suggest that GABP also regulates transcription from the rB1 promoter but uses a different binding site compared to mB1.

GABP is required for full NKG2D-L expression *in vivo*

To show that GABP is involved in the transcription of the natural *Nkg2d* locus, we adopted an RNAi strategy. We examined the effects of transient siRNA knockdown




Figure 5. GABP forms protein–DNA complex within rB1 in EMSA. **(A)** DNA sequence alignment of mB1 and rB1. Identical bases are indicated by vertical lines, and absent bases are shown as double dashes. Potential GABP-binding sites are labelled with brackets. **(B)** Sequences of probes and competitors used are shown at the top. Mutations are underlined. As control, lane 1 contains only γ^{32} P-labelled GABP1 and GABP2 WT probes. RNK16 nuclear extract was incubated with labelled GABP1 and GABP2 WT probes at 4°C for 20 min (lanes 2 and 5). For competition assays, unlabelled competitors were incubated with LNK nuclear extract at 4°C for 20 min before addition of labelled WT probe for another 20-min incubation. WT competitors were used at 100-fold molar excess (lanes 3 and 6). Mutant competitors were also used at 100-fold molar excess (lanes 4 and 7). Supershift assays were carried out using 5 μ g of anti-GABP α (lanes 8 and 9) antibody.

of *Gabpa* in the LNK cell line. Transcript levels of *Gabpa*, *Nkg2d-S* and *Nkg2d-L* were assayed via real-time RT-PCR. As shown in Figure 6, *Gabpa* transcripts decreased 70% with *Gabpa* siRNA. The negative control siRNA did not decrease the expression of the three genes compared to untreated samples. *Gabpa* knockdown had no effect on *Nkg2d-S* levels but decreased the expression of *Nkg2d-L* by 24% compared to the negative control. This suggests that the mB1 associated *Nkg2d-S*, but not *Nkg2d-L*, is regulated at least in part by GABP.

Prevalence of potential GABP sites derived from B1 elements

Neither the mB1 nor the rB1 GABP-binding sites are present in the consensus sequences of 15 B1 subfamily members found in RepBase, suggesting that these sites were created through mutation after insertion. Nonetheless, we wondered how many B1 elements in the mouse genome contain such sites and if there is evidence of these elements contributing to a regulatory network as has been revealed recently for other repetitive elements (see ‘Discussion’ section). A bioinformatics scan of all potential GABP-binding motifs to see how many are located in B1 elements is impractical since the core binding sequence is short. We opted instead to compare the prevalence of GABP-site-containing B1 elements throughout

the genome and those near promoters. A survey of all B1 elements (252 832) in the mouse genome revealed that 4877 (1.93%) contain at least one GABP-binding site. For the elements (70 804) that lie in a 10 kb window centred around TSSs of all mouse RefSeq genes, we found 1240 (1.75%) to have at least one GABP-binding site. We then subjected these RefSeq genes to gene ontology (GO) analysis using BiNGO (18). A comparison of the genes with any B1 elements in the 10-kb window to genes with GABP motif-associated B1 elements showed no differences between these groups.

Mouse B1 promoter activity is not tissue specific

Since GABP is a ubiquitous transcription factor, it likely does not confer tissue specificity. To investigate this issue, we tested a 3’ truncated fragment of mB1 in pGL4B using the EL4 and NIH 3T3 cell lines. EL4 is a mouse T-lymphoblastoid cell line and NIH 3T3 is a mouse embryonic fibroblast cell line. Neither cell lines express NKG2D (Figure 2B). The promoter activity of this construct in EL4 is comparable to that of LNK cells (Figure 7). The activity in NIH 3T3 is ~3-fold higher than in LNK and EL4 cells. This data demonstrates that the promoter and enhancers within the truncated mB1 retrotransposon are not tissue specific.




Figure 6. Transient knockdown reveals requirement of GABP for full expression of *Nkg2d-S* but not *Nkg2d-L*. LNK cells were treated with GABP α or negative control Accell siRNA. Transcript levels of *Gabpa*, *Nkg2d-S* and *Nkg2d-L* were assayed by real-time RT-PCR and normalized to the levels of *Gapdh*. mRNA levels from untreated cells (incubated with carrier media but no siRNA) were set to one. Results are the mean (\pm SD) of three independent experiments. * $P = 0.03$ (Student's t -test).

Splice signals are conserved in mammals except rodents

We wondered if the splicing pattern seen in rodents and human correlated with divergence of splice signals. Alignment of nine mammalian species at the intron1–exon1b border shows the absence of splice acceptor signals in rodents: mouse, rat and guinea pig (Figure 8A). It is possible that the selection pressure to maintain this splice signal was alleviated after the inserted SINE became capable of initiating transcription of *Nkg2d-L*. An added effect is the emergence of *Nkg2d-S*, initiating at exon 1a and skipping exon 1b now that the splice acceptor is eliminated. Interestingly, out of the nine mammals, only mouse and rat have a SINE B1 element at this splice junction. Thus, the evolution of guinea pig *Nkg2d* is different from the mouse and rat subclade. Furthermore, we noted that guinea pig and cow were the only species out of the nine examined that have lost the ATG1 start codon (Figure 8B). Both retain the ATG2 start codon while hedgehog does not (Figure 8C). This data shows that the SINE is specific to a subclade of rodents, and the splice acceptor signal at the intron1–exon1b junction is not present in rodents.

DISCUSSION

In the present study, we show that the two splice forms of mouse and rat *Nkg2d* arise from usage of alternative promoters. The promoter for mouse and rat *Nkg2d-L* is a SINE retrotransposon with a functional GABP-binding site. Comparative genomics data suggest a complex evolutionary model. Since guinea pigs lack the orthologous SINE element, we believe the B1 inserted after the divergence of guinea pig from the mouse and rat subclade. The identified GABP-binding site in the mouse is not found




Figure 7. Detection of promoter activity in different cell lines. A truncated 77-bp mB1 sequence cloned into pGL4B was assayed using a dual-luciferase system in LNK, EL4 and NIH 3T3 cells. Promoter activity were normalized to co-transfected Renilla luciferase activity and calculated as fold above pGL4B. Data indicate mean (\pm SD) of at least two independent experiments.

in the consensus B1 sequence, likely emerging through mutation. This novel regulatory element may have allowed the SINE to mediate transcription of the long isoform of *Nkg2d* directly from exon 1b, lifting the selection pressure to maintain a splice acceptor signal at the intron1–exon1b junction. The mutation of this splice acceptor site forces the transcript starting from exon 1a to splice directly to exon 2, thus creating NKG2D-S. The mutation and its consequences on splicing we have just described may apply to both mouse and rat. The only difference between the two species is the position of the acquired GABP-binding site. Thus, this study presents an interesting example of convergent evolution to achieve the same means in animals that share similar niches.

The 13 amino-acid difference at the N-terminus of the two isoforms results in differential adapter affinities: NKG2D-L binds only DAP10 while NKG2D-S binds both DAP10 and DAP12 (8). Signal cascades of these two adapters are reviewed elsewhere (19). In essence, DAP12-coupled Syk and ZAP70 signalling are strongly stimulatory and initiate cytotoxic and cytokine release pathways. On the other hand, DAP10 signalling resembles co-stimulatory pathways of CD28 in T cells. While DAP10 can initiate NK killing response through Grb2/Vav1, it cannot trigger cytokine release. This is shown in the inability of anti-NKG2D treated, DAP12 knockout mouse NK cells to produce IFN γ (9). Malarkannan *et al.* (20) also showed that DAP12/Syk/ZAP70-dependent activation of PLC γ 2 leading to the Bcl10/NF κ B pathway is absolutely essential to mouse NK cytokine production. Diefenbach *et al.* reported that while the long form of NKG2D is always expressed,




Figure 8. Analysis of *NKG2D* splice signal and start codons. (A) DNA sequence alignment of nine mammalian species near the splice acceptor site of human exon 2 using the human sequence as reference. Identical bases to the human reference are shown as dashes. Absent bases are shown as double dashes. The 5' end of B1 sequences in mouse and rat are underlined. (B) DNA sequence alignment near the ATG1 start codon of human *NKG2D* using the human sequence as reference. (C) DNA sequence alignment near the ATG2 start codon of mouse *Nkg2d* using the human sequence as reference.

the short form is only expressed after NK priming with IL-2 or polyinosinic:polycytidylic acid (8). Thus, it is possible that the two differentially expressed, signalling divergent splice forms provide the mouse and rat with beneficial functional flexibility in a context-dependent manner. In our model, we also believe that the losses of ATG1 in guinea pig and in cow are separate events of convergent evolution. The *Nkg2d* genes in these two animals likely produce only the short protein form. Indeed, Fikiri *et al.* (21) has reported the characterization of a short *NKG2D* form in cow. Although the flexibility of signalling made possible to mouse and rat may not be present in guinea pigs and cows, *NKG2D* in the latter two can at least signal through DAP12 and may therefore be beneficial. Last, the loss of ATG1 in guinea pigs may also have lifted selection pressure for retention of the splice acceptor site, allowing it to mutate away.

Our results that repetitive elements aid in the regulation of an NK receptor gene resonate with findings that link an AluSx and a LINE L1M5 element to the regulation of a human NK inhibitory receptor: *KIR3DL1* (22). Of particular interest is the fact that this AluSx SINE element also contains a binding site for an ETS member, ELK1. Removal of this SINE element abolished *KIR3DL1* distal promoter activity. The enrichment of transposable elements within immune genes has been noted before as a possible mechanism of rapid evolution and functional diversification (23). Our findings fit this model. While an Alu element may have contributed to the evolution of *KIR3DL1* by supplying the gene with regulatory elements, a B1 has managed to both provide regulatory signals and aid in the formation of a second, functionally distinct isoform of *NKG2D* in mouse and rat.

Recent ChIP-sequencing studies have shown that some classes of repetitive elements bind specific transcription

factors and establish transcription regulatory networks on a genome-wide basis. Binding for the transcription factor p53 was shown to be enriched in LTR10 and MER61 elements of the endogenous retrovirus ERV1 family (24,25). Bourque *et al.* (25) further showed that ESR1, POU5F1-SOX2 and CTCF-binding sites are pervasive in MIR, ERVK and B2 repeats, respectively. Finally, Roman *et al.* (26) described a new subclass of B1 element that harbours functional AhR and Slug-binding sites. Thus some classes of transposable elements are postulated to disperse specific control elements and drive regulatory expansion. We therefore questioned if B1 elements in the mouse genome may also be involved in the regulatory network of GABP. Since only human GABP ChIP-sequencing data is available (27) and B1 is a rodent-specific SINE element, we took an *in silico* approach. Genomic distribution and GO analysis did not reveal signs of selective retention of B1 elements with GABP sites near promoters of any gene classes. Thus, there is presently no evidence for this transposon family acting as the foundation of GABP-regulatory networks. However, it is possible that other B1 elements do bind GABP *in vivo* and could regulate genes. A ChIP-sequencing experiment in mouse would be needed to address this possibility directly.

B1 elements evolved from the 7SL RNA gene, a small non-coding RNA species. As such, they are thought to be transcribed through an internal polymerase III promoter (28). The B1 element identified that regulates mouse and rat *Nkg2d* represents a novel source of polymerase II promoter signals. B2 elements, a close relative of B1 that evolved from tRNA, have been found to provide a polymerase II promoter to the mouse gene *lama3* (29). To our knowledge, this is the first instance that B1 elements are shown to also contain polymerase II promoters.

Presently, we cannot explain why promoter activity decreases when the mB1 element is combined with 391 bp of upstream region. At least two non-mutually exclusive explanations are possible. First, the 391-bp fragment may contain an uncharacterized repressor. A search using TESS locates multiple transcription factor-binding sites. It remains to be seen if any of these are functional and bind negative regulators. However, the promoter activity of a construct containing mB1 decreases only slightly if the 391-bp fragment was inserted downstream of the reporter gene. This suggests that any repressor present on the fragment would be position dependent. The other possibility is that of chromatin effects. The reporter assays were conducted using a transiently transfected plasmid. This experimental system does not take into account the three-dimensional conformations or the degree of histone packaging of the DNA fragment in the context of the *Nkg2d* gene. It is possible that the suppressive effect of the 391-bp segment is not seen in the endogenous context.

The mechanisms that underlie the increased promoter activity with 3' truncated mB1 are also unclear. Again, it is possible that the truncations removed repressor-binding sites. At least four sites found via TESS in the truncated regions bind transcription factors known to be both activating and repressing. Two sites can bind TCF1, a factor that is activating when bound to β -catenin but repressive when interacting with Groucho (30). YY1 is a repressor when not interacting with E1A (31), while E12/E47 has been shown to act as a repressor to E-cadherin (32). Whether any of these sites are functional remains to be shown. An additional possibility is the presence of an ATG sequence that is removed in all 3' truncation constructs. Others have noted the possibility that ATG start codons can compete with reporter gene start codon and decrease reporter signal (33).

The ETS member GABP was first implicated to regulate a range of house-keeping genes such as ribosomal protein genes and a cytochrome c oxidase gene (34–36). GABP has also been linked to critical S-phase genes and cell-cycle progression (37). Recently, a string of reports have shown that GABP plays an important role in immune functions. In the myeloid lineage, GABP is known to regulate the genes for integrin *CD18* as well as IgA Fc receptor (38,39). The effect of GABP in lymphocyte biology is even more prominent. GABP was recently shown to regulate Fc receptor γ using monocyte, basophile and T cell lines (40). In T cells, GABP bridges JNK/SAPK signalling and enhanced IL-2 expression when artificially stimulated (41). GABP is also known to enhance transcription of Fas, IL-16 and IL7 receptor α (42–44). Last, GABP deficiency leads to defective B cell development, presumably because GABP binds to regulatory regions of *Pax5* and *CD79a* (45). In this report, we link GABP directly to NK cells by showing its positive regulatory effects on mouse and rat *Nkg2d*.

ETS family members share highly similar binding sites. In a genome wide study, Hollenhorst *et al.* (46) reported widespread co-occupancy of GABP, ETS1 and ELF1 at promoters of human genes. We cannot rule out the possibility that the binding site we discovered bind these factors as well. GABP may compete with ETS1 or

ELF1 for binding while all three may act synergistically to enhance *Nkg2d-L* transcription. This would reflect the *CD18* example, where GABP and PU.1 compete for binding sites but cooperatively activate the gene (38). Alternatively, these ETS members may not be competing but rather forming a complex in mB1. Physical interaction between the three GABP, ETS or ELF1 has not been described, but GABP has been shown to bind two other ETS members, ER71 and ER81 (47).

While EMSA experiments showed robust binding of GABP to the B1 elements, knocking down *Gabpa* showed only a moderate decrease in *Nkg2d-L* transcript levels. This could be due to an incomplete knockdown of the transcription factor. We only achieved ~70% silencing. The remaining GABP may be enough to sustain the observed *Nkg2d-L* expression. Another possibility is that GABP augments but is not absolutely required for *Nkg2d-L* expression. Other transcription factors that bind the B1 promoter may need to be silenced to see a more dramatic decrease to *Nkg2d-L* mRNA levels. Finally, similar ETS family binding motifs may allow other ETS members to bind the B1 element and compensate for the loss of GABP.

In mouse, NKG2D is known to be expressed only on NK cells, activated CD8⁺ T cells and some NKT and $\gamma\delta$ T cells (48). The fact that the mB1 promoter is capable of inducing reporter gene expression in NK, T and embryonic fibroblast cell lines leaves the question of NKG2D tissue-specific expression open. This is perhaps not surprising since GABP is a factor expressed in a broad range of tissue (47). We can think of at least two ways that specificity of *Nkg2d-L* expression can be achieved. There could be additional enhancer sites that bind tissue-specific transcription factors we have not uncovered. Another possibility is epigenetic regulation through DNA methylation and histone modifications. Although the upstream regions of *Nkg2d* exons 1a and 1b are CpG poor, the expression of *Ly49a*, another NK receptor gene, was shown to correlate with methylation at a CpG poor promoter (49). The expression of other NK receptor genes within the NK complex, *Ly49g* and *Nkg2a*, have also been linked to CpG methylation or histone acetylation (50,51). In addition, Sautourlidis *et al.* (52) suggests in a recent report that euchromatic histone marks open up the KIR loci early in development but use DNA methylation to silence specific KIR members and establish clonal expression later on.

As more studies emerge, the differences between human and mouse NK cells are starting to be appreciated. While mouse NKG2D-L is the isoform that more closely mimics human NKG2D, there are notable discrepancies. NKG2D is expressed on human CD8⁺ T cells regardless of activation status in human, but only on activated mouse CD8⁺ T cells. The receptor is also found on $\gamma\delta$ T cells of intestinal intraepithelial origin in human but not mouse (48). Since human NKG2D does not harbour a SINE element at the same position as the mouse, the mode of regulation is unknown and likely different from mouse. Furthermore, nothing is known about the regulation of the mouse short isoform, which responds to IL-2 and polyinosinic:polycytidylic acid treatment (8).

Dissecting the genomic and epigenetic factors that control these processes have therapeutic potential since it may allow the manipulation of an important NK-activating receptor involved in cancer immunosurveillance, viral clearance and autoimmune diseases.

ACKNOWLEDGEMENTS

We thank Drs Irina Maksakova and Ping Xiang for valuable advice and Drs Arefeh Rouhi and Fumio Takei for advice and critical reading of the manuscript.

FUNDING

Canadian Institutes of Health Research [15082 and 10825] and the Cancer Research Society, BC Cancer Agency; University of British Columbia Graduate Fellowship (to C.B.L.); Alexander Graham Bell Graduate Scholarship, The Natural Sciences and Engineering Research Council of Canada (to Y.Z.). Funding for open access charge: Canadian Institutes of Health Research.

Conflict of interest statement. None declared.

REFERENCES

- Malarkannan, S. (2006) The balancing act: inhibitory Ly49 regulate NKG2D-mediated NK cell functions. *Semin. Immunol.*, **18**, 186–192.
- Lanier, L.L. (2008) Up on the tightrope: natural killer cell activation and inhibition. *Nat. Immunol.*, **9**, 495–502.
- Vivier, E., Nunes, J.A. and Vely, F. (2004) Natural killer cell signaling pathways. *Science*, **306**, 1517–1519.
- Eagle, R.A. and Trowsdale, J. (2007) Promiscuity and the single receptor: NKG2D. *Nat. Rev. Immunol.*, **7**, 737–744.
- Guerra, N., Tan, Y.X., Joncker, N.T., Choy, A., Gallardo, F., Xiong, N., Knoblaugh, S., Cado, D., Greenberg, N.M. and Raulet, D.H. (2008) NKG2D-deficient mice are defective in tumor surveillance in models of spontaneous malignancy. *Immunity*, **28**, 571–580.
- Jonjic, S., Babic, M., Polic, B. and Krmpotic, A. (2008) Immune evasion of natural killer cells by viruses. *Curr. Opin. Immunol.*, **20**, 30–38.
- Jamieson, A.M., Diefenbach, A., McMahon, C.W., Xiong, N., Carlyle, J.R. and Raulet, D.H. (2002) The role of the NKG2D immunoreceptor in immune cell activation and natural killing. *Immunity*, **17**, 19–29.
- Diefenbach, A., Tomasello, E., Lucas, M., Jamieson, A.M., Hsia, J.K., Vivier, E. and Raulet, D.H. (2002) Selective associations with signaling proteins determine stimulatory versus costimulatory activity of NKG2D. *Nat. Immunol.*, **3**, 1142–1149.
- Zompi, S., Hamerman, J.A., Ogasawara, K., Schweighoffer, E., Tybulewicz, V.L., Di Santo, J.P., Lanier, L.L. and Colucci, F. (2003) NKG2D triggers cytotoxicity in mouse NK cells lacking DAP12 or syk family kinases. *Nat. Immunol.*, **4**, 565–572.
- Markiewicz, M.A., Carayannopoulos, L.N., Naidenko, O.V., Matsui, K., Burack, W.R., Wise, E.L., Fremont, D.H., Allen, P.M., Yokoyama, W.M., Colonna, M. *et al.* (2005) Costimulation through NKG2D enhances murine CD8⁺ CTL function: similarities and differences between NKG2D and CD28 costimulation. *J. Immunol.*, **175**, 2825–2833.
- Wu, J., Cherwinski, H., Spies, T., Phillips, J.H. and Lanier, L.L. (2000) DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. *J. Exp. Med.*, **192**, 1059–1068.
- Billadeau, D.D., Upshaw, J.L., Schoon, R.A., Dick, C.J. and Leibson, P.J. (2003) NKG2D-DAP10 triggers human NK cell-mediated killing via a syk-independent regulatory pathway. *Nat. Immunol.*, **4**, 557–564.
- Andre, P., Castriconi, R., Espeli, M., Anfossi, N., Juarez, T., Hue, S., Conway, H., Romagne, F., Dondero, A., Nanni, M. *et al.* (2004) Comparative analysis of human NK cell activation induced by NKG2D and natural cytotoxicity receptors. *Eur. J. Immunol.*, **34**, 961–971.
- Groh, V., Rhinehart, R., Randolph-Habecker, J., Topp, M.S., Riddell, S.R. and Spies, T. (2001) Costimulation of CD8 $\alpha\beta$ T cells by NKG2D via engagement by MIC induced on virus-infected cells. *Nat. Immunol.*, **2**, 255–260.
- Maksakova, I.A. and Mager, D.L. (2005) Transcriptional regulation of early transposon elements, an active family of mouse long terminal repeat retrotransposons. *J. Virol.*, **79**, 13865–13874.
- Berg, S.F., Dissen, E., Westgaard, I.H. and Fossum, S. (1998) Molecular characterization of rat NKR-P2, a lectin-like receptor expressed by NK cells and resting T cells. *Int. Immunol.*, **10**, 379–385.
- Jordan, I.K., Rogozin, I.B., Glazko, G.V. and Koonin, E.V. (2003) Origin of a substantial fraction of human regulatory sequences from transposable elements. *Trends Genet.*, **19**, 68–72.
- Maere, S., Heymans, K. and Kuiper, M. (2005) BiNGO: a cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. *Bioinformatics*, **21**, 3448–3449.
- Tassi, I., Klesney-Tait, J. and Colonna, M. (2006) Dissecting natural killer cell activation pathways through analysis of genetic mutations in human and mouse. *Immunol. Rev.*, **214**, 92–105.
- Malarkannan, S., Regunathan, J., Chu, H., Kutlesa, S., Chen, Y., Zeng, H., Wen, R. and Wang, D. (2007) Bcl10 plays a divergent role in NK cell-mediated cytotoxicity and cytokine generation. *J. Immunol.*, **179**, 3752–3762.
- Fikri, Y., Nyabenda, J., Content, J. and Huygen, K. (2007) Cloning, sequencing, and cell surface expression pattern of bovine immunoreceptor NKG2D and adaptor molecules DAP10 and DAP12. *Immunogenetics*, **59**, 653–659.
- Stulberg, M.J., Wright, P.W., Dang, H., Hanson, R.J., Miller, J.S. and Anderson, S.K. (2007) Identification of distal KIR promoters and transcripts. *Genes Immun.*, **8**, 124–130.
- van de Lagemaat, L.N., Landry, J.R., Mager, D.L. and Medstrand, P. (2003) Transposable elements in mammals promote regulatory variation and diversification of genes with specialized functions. *Trends Genet.*, **19**, 530–536.
- Wang, T., Zeng, J., Lowe, C.B., Sellers, R.G., Salama, S.R., Yang, M., Burgess, S.M., Brachmann, R.K. and Haussler, D. (2007) Species-specific endogenous retroviruses shape the transcriptional network of the human tumor suppressor protein p53. *Proc. Natl Acad. Sci. USA*, **104**, 18613–18618.
- Bourque, G., Leong, B., Vega, V.B., Chen, X., Lee, Y.L., Srinivasan, K.G., Chew, J.L., Ruan, Y., Wei, C.L., Ng, H.H. *et al.* (2008) Evolution of the mammalian transcription factor binding repertoire via transposable elements. *Genome Res.*, **18**, 1752–1762.
- Roman, A.C., Benitez, D.A., Carvajal-Gonzalez, J.M. and Fernandez-Salguero, P.M. (2008) Genome-wide B1 retrotransposon binds the transcription factors dioxin receptor and slug and regulates gene expression in vivo. *Proc. Natl Acad. Sci. USA*, **105**, 1632–1637.
- Valouev, A., Johnson, D.S., Sundquist, A., Medina, C., Anton, E., Batzoglou, S., Myers, R.M. and Sidow, A. (2008) Genome-wide analysis of transcription factor binding sites based on ChIP-seq data. *Nat. Methods*, **5**, 829–834.
- Kriegs, J.O., Churakov, G., Jurka, J., Brosius, J. and Schmitz, J. (2007) Evolutionary history of 7SL RNA-derived SINES in supraprimates. *Trends Genet.*, **23**, 158–161.
- Ferrigno, O., Virolle, T., Djabari, Z., Ortonne, J.P., White, R.J. and Aberdam, D. (2001) Transposable B2 SINE elements can provide mobile RNA polymerase II promoters. *Nat. Genet.*, **28**, 77–81.
- Roose, J. and Clevers, H. (1999) TCF transcription factors: molecular switches in carcinogenesis. *Biochim. Biophys. Acta*, **1424**, M23–M37.
- Gordon, S., Akopyan, G., Garban, H. and Bonavida, B. (2006) Transcription factor YY1: structure, function, and therapeutic implications in cancer biology. *Oncogene*, **25**, 1125–1142.
- Perez-Moreno, M.A., Locascio, A., Rodrigo, I., Dhondt, G., Portillo, F., Nieto, M.A. and Cano, A. (2001) A new role for E12/E47

- in the repression of E-cadherin expression and epithelial-mesenchymal transitions. *J. Biol. Chem.*, **276**, 27424–27431.
33. Davies, G.E., Locke, S.M., Wright, P.W., Li, H., Hanson, R.J., Miller, J.S. and Anderson, S.K. (2007) Identification of bidirectional promoters in the human KIR genes. *Genes Immun.*, **8**, 245–253.
 34. Genuario, R.R., Kelley, D.E. and Perry, R.P. (1993) Comparative utilization of transcription factor GABP by the promoters of ribosomal protein genes rpl30 and rpl32. *Gene Expr.*, **3**, 279–288.
 35. Genuario, R.R. and Perry, R.P. (1996) The GA-binding protein can serve as both an activator and repressor of ribosomal protein gene transcription. *J. Biol. Chem.*, **271**, 4388–4395.
 36. Sucharov, C., Basu, A., Carter, R.S. and Avadhani, N.G. (1995) A novel transcriptional initiator activity of the GABP factor binding ets sequence repeat from the murine cytochrome c oxidase vb gene. *Gene Expr.*, **5**, 93–111.
 37. Yang, Z.F., Mott, S. and Rosmarin, A.G. (2007) The ets transcription factor GABP is required for cell-cycle progression. *Nat. Cell Biol.*, **9**, 339–346.
 38. Rosmarin, A.G., Caprio, D.G., Kirsch, D.G., Handa, H. and Simkevich, C.P. (1995) GABP and PU.1 compete for binding, yet cooperate to increase CD18 ($\beta 2$ leukocyte integrin) transcription. *J. Biol. Chem.*, **270**, 23627–23633.
 39. Shimokawa, T. and Ra, C. (2005) C/EBP α functionally and physically interacts with GABP to activate the human myeloid IgA fc receptor (fc α R, CD89) gene promoter. *Blood*, **106**, 2534–2542.
 40. Takahashi, K., Hayashi, N., Shimokawa, T., Umehara, N., Kaminogawa, S. and Ra, C. (2008) Cooperative regulation of fc receptor gamma-chain gene expression by multiple transcription factors, including Sp1, GABP, and elf-1. *J. Biol. Chem.*, **283**, 15134–15141.
 41. Hoffmeyer, A., Avots, A., Flory, E., Weber, C.K., Serfling, E. and Rapp, U.R. (1998) The GABP-responsive element of the interleukin-2 enhancer is regulated by JNK/SAPK-activating pathways in T lymphocytes. *J. Biol. Chem.*, **273**, 10112–10119.
 42. Li, X.R., Chong, A.S., Wu, J., Roebuck, K.A., Kumar, A., Parrillo, J.E., Rapp, U.R., Kimberly, R.P., Williams, J.W. and Xu, X. (1999) Transcriptional regulation of fas gene expression by GA-binding protein and AP-1 in T cell antigen receptor.CD3 complex-stimulated T cells. *J. Biol. Chem.*, **274**, 35203–35210.
 43. Bannert, N., Avots, A., Baier, M., Serfling, E. and Kurth, R. (1999) GA-binding protein factors, in concert with the coactivator CREB binding protein/p300, control the induction of the interleukin 16 promoter in T lymphocytes. *Proc. Natl Acad. Sci. USA*, **96**, 1541–1546.
 44. Xue, H.H., Bollenbacher, J., Rovella, V., Tripuraneni, R., Du, Y.B., Liu, C.Y., Williams, A., McCoy, J.P. and Leonard, W.J. (2004) GA binding protein regulates interleukin 7 receptor α -chain gene expression in T cells. *Nat. Immunol.*, **5**, 1036–1044.
 45. Xue, H.H., Bollenbacher-Reilly, J., Wu, Z., Spolski, R., Jing, X., Zhang, Y.C., McCoy, J.P. and Leonard, W.J. (2007) The transcription factor GABP is a critical regulator of B lymphocyte development. *Immunity*, **26**, 421–431.
 46. Hollenhorst, P.C., Shah, A.A., Hopkins, C. and Graves, B.J. (2007) Genome-wide analyses reveal properties of redundant and specific promoter occupancy within the ETS gene family. *Genes Dev.*, **21**, 1882–1894.
 47. Brown, T.A. and McKnight, S.L. (1992) Specificities of protein-protein and protein-DNA interaction of GABP α and two newly defined ets-related proteins. *Genes Dev.*, **6**, 2502–2512.
 48. Raulet, D.H. (2003) Roles of the NKG2D immunoreceptor and its ligands. *Nat. Rev. Immunol.*, **3**, 781–790.
 49. Rouhi, A., Gagnier, L., Takei, F. and Mager, D.L. (2006) Evidence for epigenetic maintenance of Ly49a monoallelic gene expression. *J. Immunol.*, **176**, 2991–2999.
 50. Rouhi, A., Brooks, C.G., Takei, F. and Mager, D.L. (2007) Plasticity of Ly49g expression is due to epigenetics. *Mol. Immunol.*, **44**, 821–826.
 51. Rogers, S.L., Rouhi, A., Takei, F. and Mager, D.L. (2006) A role for DNA hypomethylation and histone acetylation in maintaining allele-specific expression of mouse NKG2A in developing and mature NK cells. *J. Immunol.*, **177**, 414–421.
 52. Santourlidis, S., Graffmann, N., Christ, J. and Uhrberg, M. (2008) Lineage-specific transition of histone signatures in the killer cell ig-like receptor locus from hematopoietic progenitor to NK cells. *J. Immunol.*, **180**, 418–425.