



This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document, This is a non-final version of an article published in final form in Read, Paul and Oliver, J L and De Ste Croix, Mark B and Myer, G D and Lloyd, R S (2016) Reliability of the Tuck Jump Injury Risk Screening Assessment in Elite Male Youth Soccer Players. Journal of Strength and Conditioning Research, 30 (6). pp. 1510-1516. ISSN 1064-8011. and is licensed under Creative Commons: Attribution-Noncommercial 3.0 license:

**Read, Paul, Oliver, Jon L, De Ste Croix, Mark B ORCID
logoORCID: <https://orcid.org/0000-0001-9911-4355>, Myer,
Gregory D and Lloyd, Rhodri S (2016) Reliability of the Tuck
Jump Injury Risk Screening Assessment in Elite Male Youth
Soccer Players. Journal of Strength and Conditioning
Research, 30 (6). pp. 1510-1516.
doi:10.1519/JSC.0000000000001260**

Official URL:

http://journals.lww.com/nsca-jscr/Abstract/2016/06000/Reliability_of_the_Tuck_Jump_Injury_Risk_Screening.2.aspx

DOI: <http://dx.doi.org/10.1519/JSC.0000000000001260>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/3152>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

This is a peer-reviewed, post-print (final draft post-refereeing) version of the following published document:

Read, Paul and Oliver, J L and De Ste Croix, Mark B and Myer, G D and Lloyd, R S (2015). *Reliability Of The Tuck Jump Injury Risk Screening Assessment In Elite Male Youth Soccer Players*. Journal of Strength and Conditioning Research. ISSN 1064-8011

Published in Journal of Strength and Conditioning Research, and available online at:

<https://www.terkko.helsinki.fi/journal-of-strength...>

We recommend you cite the published (post-print) version.

The URL for the published version is <https://www.terkko.helsinki.fi/journal-of-strength...>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

AUTHOR QUERIES

DATE 12/28/2015

JOB NAME JSCR

ARTICLE JSCR-08-6219

QUERIES FOR AUTHORS Read et al

THIS QUERY FORM MUST BE RETURNED WITH ALL PROOFS FOR CORRECTIONS

- AU1) Please check the running head has been taken from pdf. Please check.
- AU2) Please provide middle initial for the author "Paul Read." Also note there is discrepancy between manuscript and pdf in the same author name.
- AU3) Please provide department/unit (if any) for affiliations "1, 3, and 6." Also provide city details for affiliations "2 and 3."
- AU4) Please note that the corresponding author details has been taken from pdf. Please check.
- AU5) Please note Current Position is not allowed as per style. Hence this has been deleted. Please check.
- AU6) Please provide the age ranges for all subjects mentioned in the Subjects section if you have not already. If any subjects were <18 years of age, please confirm that parental consent was obtained. Please also confirm, if you have not already, that informed consent documents were signed for all subjects.
- AU7) Please check whether the edits to the sentence "Further analysis was also completed ..." are correct.
- AU8) Please check the suggested in-text citation for Figure 1 and correct the location if necessary.
- AU9) Please note that the original sentence "Whereas, observed knee and thigh motion ..." has been modified to "however, observed knee and thigh motion ..." and run on with its previous sentence "Criterion deficits of knee valgus, feet ..." for clarity and continuity. Please check whether the edits made are correct and approve.
- AU10) Please note that the references have been renumbered both in text and in list so as to maintain alphabetical order in the reference list, as per the style. Please check.
- AU11) Please provide accessed date for reference 14. Also check the same reference.
- AU12) Please update reference 36.
- AU13) Please provide a suitable caption for Figure 1 in the article.

RELIABILITY OF THE TUCK JUMP INJURY RISK SCREENING ASSESSMENT IN ELITE MALE YOUTH SOCCER PLAYERS

PAUL READ,¹ JON L. OLIVER,² MARK B.A. DE STE CROIX,³ GREGORY D. MYER,^{4,5,6} AND

AU2 RHODRI S. LLOYD²

¹School of Sport, Health and Applied Science, St Mary's University, London, United Kingdom; ²Youth Physical Development Unit, School of Sport, Cardiff Metropolitan University, United Kingdom; ³Exercise and Sport Research Centre, School of Sport and Exercise, University of Gloucestershire, United Kingdom; ⁴Division of Sports Medicine, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio; ⁵Department of Pediatrics and Orthopaedic Surgery, College of Medicine, University of

AU3 Cincinnati, Cincinnati, Ohio; and ⁶The Micheli Center for Sports Injury Prevention, Boston, Massachusetts

ABSTRACT

Read, P, Oliver, JL, de Ste Croix, MBA, Myer, GD, and Lloyd, RS. Reliability of the tuck jump injury risk screening assessment in elite male youth soccer players. *J Strength Cond Res* XX(X): 000–000, 2015—Altered neuromuscular control has been suggested as a mechanism for injury in soccer players. Ligamentous injuries most often occur during dynamic movements, such as decelerations from jump-landing maneuvers where high-risk movement patterns are present. The assessment of kinematic variables during jump-landing tasks as part of a preparticipation screen is useful in the identification of injury risk. An example of a field-based screening tool is the repeated tuck jump assessment. The purpose of this study was to analyze the within-subject variation of the tuck jump screening assessment in elite male youth soccer players. Twenty-five pre-peak height velocity (PHV) and 25 post-PHV elite male youth soccer players from the academy of a professional English soccer club completed the assessment. A test-retest design was used to explore the within-subject intersession reliability. Technique was graded retrospectively against the 10-point criteria set out in the screening protocol using two-dimensional video cameras. The typical error range reported for tuck jump total score (0.90–1.01 in pre-PHV and post-PHV players respectively) was considered acceptable. When each criteria was analyzed individually, kappa coefficient determined that knee valgus was the only criterion to reach substantial agreement across the two test sessions for both groups. The results of this study suggest that although tuck jump total score may be reliably assessed in elite male youth soccer players, caution should be applied in solely interpreting the composite

score due to the high within-subject variation in a number of the individual criteria. Knee valgus may be reliably used to screen elite youth male soccer players for this plyometric technique error and for test-retest comparison. **AU5**

KEY WORDS injury screening, landing assessment, youth, soccer

INTRODUCTION

The sport of soccer imposes high physiological demand with aerobic endurance (24), speed, agility, strength, and power key determinants of elite performance (22,38). An inherent risk of injury is also present because of frequent repetitions of high-intensity movements that expose players to high joint and muscular forces (5). Incidence data in elite male youth soccer indicate that injuries occur mainly in the lower extremities (71–80%), are largely noncontact in nature, and are predominantly comprise ligament sprains at the ankle and knee (4,20,34). Specifically, knee injury incidence rates in elite male youth soccer have been reported as 0.71 injuries per player each year, equating to an average of 17 training days and 2 matches missed per season (25). Furthermore, 35% of these injuries were classified as severe (>28 days absent), and other data are available to confirm the knee as the most frequent site of major injury in elite male youth soccer, including the anterior cruciate ligament and medial collateral ligament (MCL) sprains (40).

Although limited evidence is available in male youths to confirm pertinent risk for injuries, altered neuromuscular control has been suggested as a mechanism in females and adults (1,33,36). Deficits in neuromuscular control can impose excessive stress to the passive ligamentous structures, exceeding their strength limit and increase the potential for compromised structural integrity and injury (21). Ligamentous injuries most often occur during dynamic movements, such as decelerations from jump-landing maneuvers where high risk movement patterns are present (17).

AU4 Address correspondence to Paul Read, paul.read@stmarys.ac.uk.

00(00)/1–7

Journal of Strength and Conditioning Research
© 2015 National Strength and Conditioning Association

The assessment of kinematic variables during jump-landing tasks as part of a pre-participation screen is useful in the identification of injury risk (26–28,30). A common diagnostic tool used within the available literature is a drop vertical jump (3,7,12,28,29,32). These studies have confirmed that this assessment provides both valid and reliable indicators of anterior cruciate ligament injury risk factors, including knee valgus and high knee abduction moments (12,28,29,32). Owing to the need for expensive laboratory equipment and labor-intensive testing procedures, recent investigations have explored more clinician-friendly alternatives using standard video cameras to analyze drop vertical jump tasks (28,32). However, these clinic-based methods still present limitations for youth soccer coaches working with large groups. There is a paucity of literature available in youth male populations, and when assessing large groups comprising a variety of age ranges and anthropometric profiles, it is inherently difficult to select an appropriate drop height that accounts for individual jumping and landing abilities. Additionally, during drop-landing tasks, participants begin in an elevated position; muscles that were previously at rest while standing on the box control impact forces at the point of ground contact eccentrically. In more functional soccer tasks (jumping, cutting, etc.), the involved musculature are required to perform propulsive motions (i.e., the initial jump or running action) before ground contact, and it could be inferred that drop jumping tasks may artificially induce feed-forward stabilization mechanisms. This could be considered a learned skill as preparatory muscle activation strategies are known to develop as a result of age and maturation (19). Furthermore, despite the aforementioned methods (28) providing a more time efficient and clinician-friendly approach than laboratory techniques, there is still a requirement for digitization of joint angles to determine knee valgus motion and maximal flexion angles which is time inefficient.

Repeated jumping tasks whereby athletes are required to respond to movement perturbations and forces may be more ecologically valid in sports such as soccer. An example of a field-based screening tool is the repeated tuck jump assessment (27). This method has been developed to identify neuromuscular deficits and identify lower extremity technique flaws during a plyometric exercise (27,31,35). It has been suggested that the presence of these neuromuscular deficits may be indicative of an increased risk of traumatic knee injury (26,30). Assessed using a 10-point rating scale, initial pilot studies have indicated moderate-to-strong interrater reliability (range = 0.72–0.97) (27). More recently, intra- and intertester reliability have been reported demonstrating excellent agreement (kappa measurement, $k = 0.88$) (11).

The tuck jump assessments were originally designed as a coach-friendly plyometric landing assessment and training tool which targets risk factors for ACL injury mechanisms (27,31,35). However, less information is available to confirm its reliability and validity as a diagnostic screening tool, especially in male youth soccer players who have a high incidence

of MCL injuries (25,34) where a valgus mechanism has been suggested as a high-risk movement pattern (9,16). It has been suggested that baseline levels and deficits can be compared following targeted training interventions to objectively track improvements in performance (31,35). However, no data is currently available that report within-subject variation using test-retest experimental design, especially in pediatric populations. Given the inherent risk of lower limb injury to male youth soccer players, especially around the time of peak height velocity (PHV) (39), and reported higher variability in jump performance in less mature individuals (10), it would seem pertinent to confirm the reliability of the tuck jump assessment with this particular population. The aim of this study is to determine the reliability of the tuck jump assessment tool in both pre- and post-PHV elite male youth soccer players.

METHODS

Experimental Approach to the Problem

This study used a test-retest design to explore the intersession reliability of the tuck jump assessment. Participants were required to attend the club training ground on 3 occasions, each separated by a period of 7 days. The first session was used to familiarize participants to the test, allowing both visual demonstrations and practice attempts until the principal researcher was satisfied that the technique was appropriate and consistent. Data collected in the second and third sessions were subsequently analyzed to determine within-subject variation. In each of the 2 experimental test sessions, a 10-minute standardized warm-up inclusive of a series of dynamic stretches was completed before commencement. Testing was completed at the same time on each day, and participants were asked to wear the same clothing and footwear and refrain from strenuous exercise for at least 48 hours before testing. Participants were also asked to eat according to their normal diet and avoid eating and drinking substances other than water 1 hour before each test session.

Subjects

Twenty five pre-PHV (age 11.93 ± 0.43 years; height 151.40 ± 4.84 cm; body mass 41.05 ± 5.62 kg; maturational offset -2.34 ± 0.41 years;) and 25 post-PHV (age 17.26 ± 0.69 years, stature 178.22 ± 5.47 ; body mass 72.27 ± 6.93 kg; maturity offset 2.91 ± 0.81 years) youth soccer players from the academy of a professional English Championship soccer club volunteered to take part in the study. The research design required players to complete a familiarization session and 2 experimental test sessions separated by a period of 1-week. None of the players reported injuries at the time of testing, and all were participating regularly in football training and competitions. Parental consent and participant assent were collected before the commencement of testing. In addition, subjects completed a physical activity readiness questionnaire to determine whether their health status was

appropriate and that there was no significant physical reason why they should not partake in the research project. Ethical approval was granted by the institutional ethics committee in accordance with the declaration of Helsinki. The study conforms to the Code of Ethics of the World Medical Association (approved by the ethics advisory board of Swansea University) and required players to provide informed consent before participation.

Procedures

During each recorded test session, tuck jumps were performed in place for a period of 10 seconds (27). Before the test, participants were required to place their feet 35 cm apart on two vertical strips of tape, which were connected by a horizontal line forming an H-shape (35). Participants began by performing a countermovement before jumping in a vertical direction as high as possible while simultaneously pulling their knees up toward their chest. Upon ground contact, participants were

encouraged to minimize ground contact time and use a toe to mid-foot rocking landing strategy. They were also instructed to land in the same footprint with each jump. Players were screened using a specified 10-point criteria as suggested previously (27,31,35). Deficiencies in plyometric technique were marked if it was not maintained for 10 seconds.

To increase assessment accuracy, 2-dimensional video cameras were used to capture the test, and each player's technique was graded retrospectively. Kinematic data were collected at 50 Hz using 2 high-definition video cameras (Samsung, New Jersey, USA) positioned in the frontal and sagittal planes at a height 0.70 m and at a distance of 5 m from the center of the capture area. To allow visible tracking of the knees, subjects were required to wear shorts with a line at approximately mid-thigh. When scoring each trial of the tuck jump assessment, each of the 10 criteria were analyzed, and if a deficit was present on two or more repetitions, it was




Figure 1. Reprinted, by permission, from G.D. Myer, K.R. Ford, and T.E. Hewett, 2008, "Tuck jump assessment for reducing anterior cruciate ligament injury risk," *Athletic Therapy Today* 13(5): 39–44. Human Kinetics, Inc. Adaptations are themselves works protected by copyright. So in order to publish this adaptation, authorization must be obtained both from the owner of the copyright in the original work and from the owner of copyright in the translation or adaptation.

TABLE 1. Descriptive statistics and reliability values for tuck jump (TJ) total score.*†

Test	Number of jumps		TJ total score		Change in mean	ICC	Typical error
	Test 1	Test 2	Test 1	Test 2			
TJ total score pre-PHV	14.4 ± 1.6	14.7 ± 1.48	6.88 ± 1.33	6.64 ± 1.50	-0.24 ± 1.27	0.53	0.90
TJ total score post-PHV	15.8 ± 0.8	14.4 ± 0.81†	5.44 ± 1.29	5.40 ± 1.83	0.17 ± 1.43	0.55	1.01

*ICC = intraclass correlation coefficient; PHV = peak height velocity.

†Significantly different from test 1 ($p \leq 0.05$).

marked as previously stated (27,31,35). Additionally, a composite score tallied from the recorded deficits was provided, and higher scores were indicative of reduced performance, 10 being the maximum score attainable, suggesting players should be targeted for further technique training (27). The same rater marked and recorded deficits to avoid the presence of interrater error. Further analysis was also completed to determine the frequency when each of the criteria was present and the total number of jumps in each session (Figure 1).

Statistical Analyses

Tests for normal distribution were completed on the recorded data, and means and standard deviations for tuck jump total score were calculated across the two testing sessions. Twenty-five participants per group were selected to provide a sufficiently small confidence interval to the reliability statistics. This provided a confidence interval factor of approximately 1.35 to apply random variation. Doubling the sample size would only reduce the precision of the confidence interval factor to 1.22 (13) and thus have limited effects on the outcomes or interpretations of reliability.

Reliability statistics included: change in mean, intraclass correlation coefficient (ICC) to determine rank order repeatability and typical error estimate. Reliability data were

computed through Microsoft Excel 2010 using a freely available spread sheet (14). To determine systematic bias between trials, paired samples *t*-tests were used with a *p* value ≤ 0.05 . The total number of jumps completed during each trial was calculated, and a paired samples *t*-test was used to check for a significant difference between test sessions. Furthermore, Pearson correlation coefficients were calculated to assess the relationship between tuck jump total score and the number of jumps performed. Paired samples *t*-tests and Pearson correlation coefficients were processed using SPSS (V.21, Chicago, IL, USA).

To determine the accuracy of observations across the 2 test sessions for each individual criterion displayed in the tuck jump assessment checklist, agreement was assessed using kappa coefficient with the following threshold classifications: >0.8 almost perfect agreement; 0.6–0.8 substantial agreement; 0.4–0.6 moderate agreement; 0.2–0.4 slight to fair agreement (18). Intrarater reliability was calculated for the tuck jump total score using ICC, where the same rater scored a sample of the videos from experimental test session 1 on 2 occasions separated by a period of 7 days.

RESULTS

Descriptive statistics and all reliability values are displayed in Table 1. Intrarater reliability for the 10-s repeated tuck jump

TABLE 2. Tuck jump kappa statistics.*

Phase of jump	Criterion	Kappa value (pre-PHV)	Kappa value (post-PHV)
Knee and thigh motion	Knee valgus	$k = 0.78$	$k = 0.67$
	Not parallel	$k = 0.34$	$k = 0.44$
	Not equal	$k = 0.11$	$k = 0.29$
Foot positioning during landing	Not shoulder width	$k = 0.52$	$k = 0.25$
	Not parallel	$k = 0.62$	$k = 0.20$
	Unequal contact time	$k = 0.04$	$k = 0.05$
	High-contact noise	$k = 0.34$	$k = 0.43$
Plyometric technique	Pause between jumps	$k = 0.65$	$k = 0.00$
	Declines in 10 s	$k = 0.33$	$k = 0.31$
	Excess flight motion	$k = -0.06$	$k = 0.01$

*PHV = peak height velocity.

total score was deemed strong ($ICC = 0.88$) based on previous classifications (15). A significant difference was observed between the total number of jumps performed in each test session only in the postpubertal group ($P < 0.001$) who completed less jumps in experimental test session 2. Relationships between tuck jump total score deficits and the numbers of jumps completed were significantly correlated for the prepubertal group only ($r = 0.37$; $P < 0.001$).

The kappa coefficient used to determine between-session agreement for each of the 10 individual tuck jump criteria are displayed in Table 2. Knee valgus was the only criterion that reached substantial agreement across the two test trials for both groups, suggesting strong reliability. Other measures including unparalleled foot position during landing and a pause between jumps also reported substantial agreement for the pre-PHV players; however, this was not consistent across groups. Moderate agreement ($k = 0.6-0.4$) was present for foot positioning not being shoulder width during landing (pre-PHV group only), thigh motion not reaching parallel and high contact noise (post-PHV group only) suggesting that these criteria could be measured with acceptable reliability. The agreement between all other criteria was either fair ($K < 0.40$) or weak ($K < 0.20$) indicating low test-retest reliability based on previous suggestions (18).

DISCUSSION

To the author's knowledge, this is the first study to assess the reliability of the repeated tuck jump assessment in male youth soccer players which could be included as part of a field-based neuromuscular control screening battery. Research quantifying the effects of age, growth, and maturation on the reliability of this test also does not currently exist. The total score obtained during the repeated tuck jump assessment demonstrated moderate ICC values. However, caution should be applied when using this approach for determining the reliability of a test. A retest correlation measures how closely the values of 2 trials track each other specific to each individual and the reproducibility of the rank order of subjects during the retest (13). Intraclass correlation coefficient statistics is also affected by the heterogeneity of the values between participants and the sample used for measurement (13). A sample which is of a homogenous nature will demonstrate a low value. The subjects in this study are reflective of a homogenous sample because of their status as elite male youth academy soccer players, and thus, this provides a plausible explanation for the low ICC values. Examining the results of previous reliability studies using this approach and interpreting the findings is also difficult.

An alternative and more appropriate approach is to use statistical tests that measure the typical error to show variation in the participant scores between experimental test sessions (2,15). The typical error range reported (0.89–1.01 in pre- and post-PHV players, respectively) would be considered acceptable and indicates that changes in total score >1 following an intervention would be classified as a real

change in injury risk reduction. Previous investigations have reported large percentage changes after augmented feedback interventions for various criteria in the tuck jump assessment (35), suggesting that a change in total score >1 is realistic. However, despite the acceptable typical error values reported, when each of the 10 individual criterion measures was analyzed separately, a number of these demonstrated fair to low agreement. These findings may be explained in part by the greater number of criteria included in the total tuck jump score which may have provided an averaging effect where errors begin to cancel themselves out. This is representative of taking the mean of more trials. However, owing to the paucity of literature which has investigated the reliability of this test, further research is warranted.

Criterion deficits of knee valgus, feet not parallel to shoulder width on landing, and a pause between jumps were the only measures to demonstrate acceptable reliability for pre-PHV players ($k = 0.52-0.78$); however, observed (19) knee and thigh motion variables including: knee valgus on landing, and thighs not parallel in flight and high contact noise were the only criteria to report acceptable reliability in the post-pubertal group ($k = 0.43-0.67$). This suggests that these plyometric technique flaws can be reliably assessed as part of a test-retest screen in pre-PHV children. Thus, caution should be applied in purely interpreting the composite assessment score due to the high within-subject variation in a number of the individual criteria. Respective of a change in total score >1 which was identified as a reliable performance modification, it would be uncertain as to how injury risk has changed.

Although this is the only study to analyze test-retest within-subject reliability for each of the individual criteria during the tuck jump assessment, knee valgus on landing reported the strongest agreement for both groups. This suggests that screening for this criterion risk factor is reliable for test-retest comparison in male youth soccer players who are either pre- or post-PHV. Knee injuries comprise 17% of all injuries in elite male youth soccer players (25) and the valgus mechanism has been suggested as a high-risk movement pattern for incidences of both medial MCL (9,16) and ACL injury (8,12). MCL injuries occur frequently in male youth soccer accounting for up to 85% of knee ligament injuries (19,33), and ACL lesions are the most frequently occurring serious injury (>28 days absence) in this cohort (40). Knee valgus motion has been cited as a key contributing factor to high knee abduction moments and ACL injuries (29) and is a predictor of injury assessed during jumping and landing tasks (12). Therefore, practitioners using the tuck jump assessment to screen male youth soccer players may reliably track changes in knee valgus on landing following an intervention.

There is a paucity of available literature to compare the current findings with other studies, and specifically, no data are available in male youth of different ages or stages of maturation. In their initial review of this assessment, Myer et al. (27) stated that unpublished pilot studies indicated

moderate-to-strong interrater reliability (range = 0.72–0.97). More recently, Herrington et al. (11) assessed the main outcome scores of males and females using two independent assessors demonstrating good intra- and intertester reliability which is in agreement with the findings of this study (intra rater reliability = 0.88). This suggests that variation in test scores is most likely because of the participants and not the rater. However, other investigations have shown poor interrater and poor-to-moderate intrarater reliability (6), and none of these studies investigated within-subject reliability. In the present study, high within-subject variance was present as indicated by low kappa agreement between trials on a number of the test criteria. This finding is not unexpected, as obtaining high reliability for repeated trials during tasks requiring dynamic jumping and landing activities that are reliant on reflexive muscle responses, and proprioceptive and kinesthetic feedback will likely use a range of movement strategies, thus increasing variability (7,23,37). Previous literature has suggested greater variation in jump performance in younger subjects (10); lower levels of consistency in technique can be expected in youth players who may experience difficulty in consistently executing the correct jump-landing pattern.

A significant difference was noted between the numbers of jumps completed in each test in the post-pubertal group. This occurred despite the standardized 10-s protocol which has been previously suggested (27,31,35). Measures need to be reproducible if their purpose is for repeated screening. Intuitively, a player who performs more or less jumps during a retest may heighten the chance of deficit changes (either higher or lower) being reported purely as a result of altering the number of jumps completed. Specifically, a greater number of jumps will typically increase the likelihood of more errors, and this may partly explain the variation in jump performance between the two trials in the present study. This was likely evident in the pre-PHV group supported by a significant correlation between the number of jumps performed and the tuck jump total score. In addition, fewer jumps during the 10-s test period may be representative of more powerful jump performance because of increases in flight time. Cumulatively, these findings suggest that for practitioners who wish to use this assessment for regular athlete monitoring and comparison between groups, standardizing the number of jumps performed may be recommended.

PRACTICAL APPLICATIONS

This study is novel as data are now available to determine the within-subject reliability of the repeated tuck jump assessment in male youth soccer players. The total score comprised of 10 criterion measures reported acceptable within-subject variation and indicated that a change in score >1 is needed to reliably state that performance has improved. However, when each criterion was analyzed individually, moderate-substantial agreement between trials was noted in some of the assessed criterion, but these differed in each age group and other criteria reported low agreement.

Therefore, caution should be applied when solely interpreting the composite score because of uncertainty of how their tuck jump technique has changed and subsequently which potential injury risk factors are present.

Knee valgus was the only measure to demonstrate substantial agreement in both groups and may be reliably used to screen male youth soccer players for this potential injury risk factor and for test-retest comparison, on the provision that the total number of jumps performed is standardized. Thus, it is suggested that the tuck jump could be used to effectively assess jump-landing technique but only the presence of knee valgus should be evaluated. This provides a time-efficient tool for practitioners who screen large groups of athletes as part of regular field-based screening protocols. In addition, lower total scores reported in the post-PHV players on both trials tentatively suggest that performance in the tuck jump assessment may improve as a result of growth and maturation.

REFERENCES

- Alentorn-Geli, E, Myer, GD, Silvers, HJ, Samitier, G, Romero, D, Lázaro-Haro, C, and Cugat, R. Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surg Sports Traumatol Arthrosc* 17: 705–729, 2009.
- Atkinson, G and Nevill, AM. Statistical methods for addressing measurement error (reliability) in variables relevant to sports medicine. *Sports Med* 26: 217–238, 1998.
- Barber-Westin, SD, Noyes, FR, and Galloway, M. Jump land characteristics and muscle strength development in young athletes. A gender comparison of 1140 athletes 9–17 years of age. *Am J Sports Med* 34: 375–382, 2006.
- Cloke, D, Spencer, S, Hodson, A, and Deehan, D. The epidemiology of ankle injuries occurring in English Football Association Academies. *Br J Sports Med* 43: 1119–1125, 2009.
- Daniel, DM, Stone, ML, Dobson, BE, Fithian, DC, Rossman, DJ, and Kaufman, RF. Fate of the ACL injured patient: A prospective outcome study. *Am J Sports Med* 22: 632–644, 1994.
- Dudley, LA, Smith, CA, Olson, BK, Chimera, NJ, Schmitz, B, and Warren, M. Interrater and intrarater reliability of the tuck jump assessment by health professionals of varied educational backgrounds. *J Sports Med* 2013: 5, 2013.
- Ebben, WP, VanderZanden, T, Wurm, BJ, and Petushek, EJ. Evaluating plyometric exercises using time to stabilization. *J Strength Cond Res* 24: 300–306, 2010.
- Ford, KR, Myer, GD, and Hewett, TE. Valgus knee motion during landing in high school female and male basketball players. *Med Sci Sports Exerc* 35: 1745–1750, 2003. **AU10**
- Gardiner, JC, Weiss, JA, and Rosenberg, TD. Strain in the human medial collateral ligament during valgus loading of the knee. *Clin Orthop Relat Res* 391: 266–274, 2001.
- Gerodimos, V, Zafeiridis, A, Perkos, S, Dipla, K, Manou, V, and Kellis, S. The contribution of stretch-shortening cycle and arm-swing to vertical jumping performance in children, adolescents, and adult basketball players. *Pediatr Exerc Sci* 20: 379–389, 2008.
- Herrington, L, Myer, GD, and Munro, A. Intra and inter-tester reliability of the tuck jump assessment. *Phys Ther Sport* 14: 152–155, 2013.
- Hewett, TE, Myer, GD, Ford, KR, Heidt, RS, Colosimo, AJ, McLean, SG, van den Bogert, AJ, Paterno, MV, and Succop, P. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: A prospective study. *Am J Sports Med* 33: 492–501, 2005.

- AU11**
13. Hopkins, WG. Measures of reliability in sports medicine and science. *Sports Med* 30: 1–15, 2000.
 14. Hopkins, WG. Reliability from consecutive pairs of trials (Excel spreadsheet). In: *A New View of Statistics*. 2000. sports.org: Internet Society for Sport Science. Available at: sports.org/resource/stats/xrely.xls.
 15. Hopkins, WG, Hawley, JA, and Burke, LM. Design and analysis of research on sports performance enhancement. *Med Sci Sports Exerc* 31: 472–485, 1999.
 16. Indelicato, PA. Isolated medial collateral ligament injuries in the knee. *J Am Acad Orthop Surg* 3: 9–14, 1995.
 17. Krosshaug, T, Nakamae, A, Boden, BP, Engebretsen, L, Smith, G, Slauterbeck, JR, Hewett, TE, and Bahr, R. Mechanisms of anterior cruciate ligament injury in basketball: Video analysis of 39 cases. *Am J Sports Med* 35: 359–367, 2007.
 18. Landis, JR and Koch, GG. Agreement measures for categorical data. *Biometrics* 33: 159–174, 1977.
 19. Lazaridis, S, Bassa, E, Patikas, P, Giakas, G, Gollhofer, A, and Kotzamanidis, C. Neuromuscular differences between prepubescent boys and adult men during drop jump. *Eur J Appl Physiol* 110: 67–74, 2010.
 20. Le Gall, F, Carling, C, Reilly, T, Vandewalle, H, Chrucho, J, and Rochcongar, P. Incidence of injuries in elite French youth soccer players: A 10-Season study. *Am J Sports Med* 34: 928–938, 2006.
 21. Li, G, Rudy, TW, Allen, C, Sakane, M, and Woo, LS. Effect of combined axial compressive and anterior tibial loads on in situ forces in the anterior cruciate ligament: A porcine study. *J Orthop Res* 16: 122–127, 1998.
 22. Little, T and Williams, AG. Specificity of acceleration, maximum speed, and agility in professional soccer players. *J Strength Cond Res* 19: 76–78, 2005.
 23. Lloyd, RS, Oliver, JL, Hughes, MG, and Williams, CA. Reliability and validity of field-based measures of leg stiffness and reactive strength index in youths. *J Sports Sci* 27: 1565–1573, 2009.
 24. McMillan, K, Helgerud, J, Macdonald, R, and Hoff, J. Physiological adaptations to soccer specific endurance training in professional youth soccer players. *Br J Sports Med* 39: 273–277, 2005.
 25. Moore, O, Cloke, DJ, Avery, PJ, Beasley, I, and Deehan, DJ. English premiership academy knee injuries: Lessons from a 5 year study. *J Sports Sci* 29: 1535–1544, 2011.
 26. Myer, GD, Brent, JL, Ford, KR, and Hewett, TE. Real-time assessment and neuromuscular training feedback techniques to prevent ACL injury in female athletes. *Strength Cond J* 33: 21–35, 2011.
 27. Myer, GD, Ford, KR, and Hewett, TE. Tuck jump assessment for reducing anterior cruciate ligament injury risk. *Athl Ther Today* 13: 39–44, 2008.
 28. Myer, GD, Ford, KR, Khoury, J, Succop, P, and Hewett, TE. Development and validation of a clinic based prediction tool to identify female athletes at high risk of ACL injury. *Am J Sports Med* 38: 2025–2033, 2010.
 29. Myer, GD, Ford, KR, Khoury, J, Succop, P, and Hewett, TE. Biomechanics laboratory-based prediction algorithm to identify female athletes with high knee loads that increase risk of ACL injury. *Br J Sports Med* 45: 245–252, 2011.
 30. Myer, GD, Paterno, MV, Ford, KR, and Hewett, TE. Neuromuscular training techniques to target deficits before return to sport after anterior cruciate ligament reconstruction. *J Strength Cond Res* 22: 987–1014, 2008.
 31. Myer, GD, Stroube, BW, DiCesare, CA, Brent, JL, Ford, KR, Heidt, RS, and Hewett, TE. Augmented feedback supports skill transfer and reduces high-risk injury landing mechanics. A double-blind, randomized controlled laboratory study. *Am J Sports Med* 41: 669–677, 2013.
 32. Padua, DA, Marshall, SW, Boling, MC, Thigpen, CA, Garrett, WE, and Beutler, AI. The Landing Error Scoring System (LESS) is a valid and reliable clinical assessment tool of jump-landing biomechanics. *Am J Sports Med* 37: 1996–2002, 2009.
 33. Paterno, MV, Schmitt, LC, Ford, KR, Rauh, MJ, Myer, GD, Haug, B, and Hewett, TE. Biomechanical measures during landing and postural stability predict a second ACL injury after ACL reconstruction and return to sport. *Am J Sports Med* 38: 1968–1978, 2010.
 34. Price, RJ, Hawkins, RD, Hulse, MA, and Hodson, A. The football association and medical research programme: An audit of injuries in academy youth football. *Br J Sports Med* 38: 466–471, 2004.
 35. Stroube, BW, Myer, GD, Brent, JL, Ford, KR, Heidt, RS, and Hewett, TE. Effects of task-specific augmented feedback on deficit modification during performance of the tuck-jump exercise. *J Sport Rehabil* 22: 7–18, 2013.
 36. Sugimoto, D, Alternton-Geli, E, Mediguchia, J, Samuelsson, K, Karlsson, J, and Myer, GD. Biomechanical and neuromuscular characteristics of male athletes: Implications for the development of anterior cruciate ligament injury prevention programs. *Sports Med*. In press. **AU12**
 37. Wikstrom, EA, Tillman, MD, Chmielewski, TL, and Borsa, PA. Measurement and evaluation of dynamic joint stability of the knee and ankle after injury. *Sports Med* 36: 393–410, 2006.
 38. Wisløff, U, Helgerud, J, and Hoff, J. Strength and endurance of elite soccer players. *Med Sci Sports Exerc* 30: 462–467, 1998.
 39. van der Sluis, A, Elferink-Gemser, MT, Coelho-e-Sliva, MJ, Nijboer, JA, Brink, MS, and Visscher, C. Sports injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med* 35: 351–355, 2014.
 40. Volpi, P, Pozzoni, R, and Galli, M. The major traumas in youth football. *Knee Surg Sports Traumatol Arthrosc* 11: 399–402, 2003.