



UNIVERSITY OF  
GLOUCESTERSHIRE

This is additional metadata of the following published document and is licensed under All Rights Reserved license:

**Hamilton, R. Liz ORCID: 0000-0003-2532-2236, Trimmer, Mark, Bradley, Chris and Pinay, Gilles (2016) Deforestation for oil palm alters the fundamental balance of the soil N cycle. Soil Biology and Biochemistry, 95. pp. 223-232.  
doi:10.1016/j.soilbio.2016.01.001**

Official URL: <http://dx.doi.org/10.1016/j.soilbio.2016.01.001>

DOI: <http://dx.doi.org/10.1016/j.soilbio.2016.01.001>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/4438>

#### **Disclaimer**

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

## Abstract

Expansion of commercial agriculture in equatorial regions has significant implications for regional nitrogen (N) budgets. Here we investigate changes in N availability and turnover in Southeast Asia following the replacement of tropical forest with oil palm plantations along a chronosequence of oil palm maturity (3-months to 15-year-old stands) and secondary to primary forest succession in Sabah, Malaysian Borneo. Ten sites were sampled during March and April 2012 and rates of gross ammonium ( $\text{NH}_4^+$ ) and nitrate ( $\text{NO}_3^-$ ) production (mineralisation and nitrification) and consumption ( $n = 8$ ), potential denitrification and “anaerobic ammonium oxidation” (“anammox”) ( $n = 12$ ) were determined using  $^{15}\text{N}$  isotope additions to soil cores and slurries respectively. Gross mineralisation rates ( $0.05\text{--}3.08 \text{ g N m}^{-2} \text{ d}^{-1}$ ) remained unchanged in oil palm relative to forests. However, a significant reduction in gross nitrification ( $0.04\text{--}2.31 \text{ g N m}^{-2} \text{ d}^{-1}$ ) and an increase in  $\text{NH}_4^+$  immobilisation disrupt the pathway to nitrogen gas ( $\text{N}_2$ ) production substantially reducing (by  $> 90\%$ ) rates of denitrification and “anammox” in recently planted oil palm relative to primary forest. Potential nitrous oxide ( $\text{N}_2\text{O}$ ) emissions were greater than potential  $\text{N}_2$  production and remained unchanged across the chronosequence indicating a potentially increased ratio of  $\text{N}_2\text{O}:\text{N}_2$  emission when soils were first disturbed. These results are an important precursor to studies that could yield improved estimates of regional N turnover and loss in Southeast Asia which will have global implications for N biogeochemical cycling.