



UNIVERSITY OF  
GLOUCESTERSHIRE

This is a peer-reviewed, final published version of the following document and is licensed under Creative Commons: Attribution 4.0 license:

**Mills, Claire D ORCID: 0000-0003-4156-4593 and Eglon, Hannah (2018) Heart Rate Intensity in Female Footballers and its effect on playing position based on external workload. Sports and Exercise Medicine Open Journal, 4 (1). pp. 24-34. doi:10.17140/SEMOJ-4-157**

Official URL: <https://openventio.org/wp-content/uploads/Heart-Rate-Intensity-in-Female-Footballers-and-its-Effect-on-Playing-Position-based-on-External-Workload-SEMOJ-4-157.pdf>  
DOI: 10.17140/SEMOJ-4-157  
EPrint URI: <https://eprints.glos.ac.uk/id/eprint/5575>

#### **Disclaimer**

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

## Original Research

# Heart Rate Intensity in Female Footballers and its Effect on Playing Position based on External Workload

Claire D. Mills, PhD\*; Hannah J. Eglon, BSc

School of Sports and Exercise, University of Gloucestershire, Oxstalls Campus, Gloucester, GL2 9HW, UK

### \*Corresponding author

Claire D. Mills, PhD

Senior Lecturer, School of Sports and Exercise, University of Gloucestershire, Oxstalls Campus, Gloucester, GL2 9HW, UK; Tel. +44 (0)1242 715156;

Fax: +44 (0)1242 715222; E-mail: [clairem@glos.ac.uk](mailto:clairem@glos.ac.uk)

### Article information

Received: April 20<sup>th</sup>, 2018; Revised: May 11<sup>th</sup>, 2018; Accepted: May 18<sup>th</sup>, 2018; Published: June 4<sup>th</sup>, 2018

### Cite this article

Mills CD, Eglon HJ. Heart rate intensity in female footballers and its effect on playing position based on external workload. *Sport Exerc Med Open J.* 2018; 4(1): 24-34. doi: <http://dx.doi.org/10.17140/SEMOJ-4-157>

## ABSTRACT

### Introduction

Female football is the world's fastest developing sport, and due to the rise in magnitude, female football, of all levels, must embrace scientific applications allowing an increase in performance through training, technique, and preparation.

### Purpose

The purpose of the study was to examine the physiological external workload, of amateur female footballers, across varying heart rate intensities, as well as, interpret fatigue between each half of the Soccer-Specific Aerobic Field Test (SAFT<sup>90</sup>) protocol.

### Methods

A sample of n=24 amateur female football players (mean±SD; age: 20.7±4.0 years; stretched stature=165.6±5.8 cm, body mass=58.1±4.7 kg) were recruited during the 2016/2017 competitive season. Maximum heart rate (HR<sub>max</sub>) values were determined using the Yo-Yo intermittent recovery level 1 (Yo-Yo IR Level 1) with the SAFT<sup>90</sup> protocol used to interpret the physiological and mechanical demands displayed during football match. A one-way analysis of variance was applied to determine the differences between each position (defenders, midfielders, and forwards) with the level of significance set at alpha level  $p < 0.05$ .

### Results

There were statistically significant differences between each position and total external workload ( $F(df:23)=9.156$ ; ( $p < 0.05$ ), in addition to average heart rate (HR) across 90-minutes ( $F(df:23)=22.317$ ; ( $p < 0.05$ ). Statistical significance determined differences between each position and the duration of time spent within the prescribed HR intensity zones, including zone 1 (<70% HR<sub>max</sub>), zone 2(70-85% HR<sub>max</sub>) and zone 4(90-95% HR<sub>max</sub>) across the SAFT<sup>90</sup>. The total distance all players completed across the 90-minutes was 10913.7±1076.7 m, whereas the shortest external workload (10020.4±1086.6 m) was completed by defenders and largest (11781.9±324.7 m) by midfielders. The average heart rate of all player's was 161.1±14.7 bpm<sup>-1</sup> resulting in a mean intensity of representing 81% HR<sub>max</sub>.

### Conclusion

Midfielders spent the largest proportion of time between zone 2 and zone 4. Based on these results, coaches are able to determine which type of physiological profile is needed for a specific position and use this information to design specifically prescribed training programmes to maximise the fitness development.

### Keywords

Football; Female football; External workload; Heart rate; Intensity.

## INTRODUCTION

Football is the world's most popular sport, practiced across the world.<sup>1</sup> The sports universal appeal is reflected by 260 million, males, females, and children, of all abilities, who participate globally, which corresponds for more than 29 million female participants.<sup>2</sup> Therefore, through obtaining a deeper level of understanding, coaches and players can benefit in terms of their ability to present improved information to modify tactics and training approaches.<sup>3</sup> With female football extensively increasing in stature, the performance expectation has further risen with an increased need for specific scientific research to improve performance.<sup>4</sup> From a physical outlook, football is an intermittent sport<sup>3,5</sup> that requires a well-developed level of conditioning and physical fitness to be played successfully.<sup>6</sup> Throughout the duration of a match, the completed amount of external work completed is characterised by the total distance covered by each player.<sup>7</sup> Nonetheless, given its inability to account for the utility movements and their complete energy costs, total distance alone is not considered to be a valid measure of overall match performance. Rather, the combination of total distance completed, alongside the measurement of heart rate throughout a football match could thus provide a more representative figure.<sup>8,9</sup>

### Physical Demands in Football

Physical demands within football have increased across the last decade<sup>3</sup> with all players now participating within defensive and offensive phases; this has created multifunctional footballers, usable within multiple positions.<sup>10</sup> Physical individual variables have been observed within playing positions in all competitive levels of football.<sup>11,12</sup> Numerous studies have compared playing position, presenting a clear link between playing position and physical capacity both aerobically and anaerobically.<sup>13</sup> Being an intermittent sport,<sup>3</sup> the aerobic system is heavily taxed with peak heart rates of 85% HR<sub>max</sub> and 98% HR<sub>max</sub>.<sup>10</sup> However, oxygen kinetics undergo changes throughout a football match due to the 150-250 anaerobic short intense bouts performed, indicating a high anaerobic energy turnover.<sup>14,15</sup> The benefits of efficient oxygen kinetics to subsequently benefit footballer's performance due to delaying fatigue and improving recovery, due to the characteristics of football, players have to attain physical qualities including high-levels of aerobic and anaerobic endurance.<sup>14,16</sup> However, Hoff and Helgerud<sup>17</sup> stated that football players do not excel in one single physical component.<sup>16,14</sup> Shalfawi et al<sup>18</sup> affirms that the ability to sustain aerobic and anaerobic endurance further to strength, power and agility for the match, particularly in the second half, are vital to aid physical performance and determine the match outcome.

### Positional Variations in External Workload

Studies show the differences in external workload throughout a match.<sup>19</sup> The energy expenditure in performance is directly linked to the mechanical external work output. Conflicting research debates the total distance travelled throughout a match; the concluded distance players cover is 8-12 km, with distances at elite levels reaching 14 km.<sup>20,16</sup> Collective studies, conclude midfielders

performing the largest total distance in comparison to other playing positions.<sup>3,10</sup> External midfielders previously recorded distances covering 11990+776 m whilst central midfielders recorded distances of 12027+625 m throughout the duration of a 90-minute match.<sup>14</sup> As a result, centre backs (10627+893 m), external defenders (11410+708 m) and forwards (11254+894 m) display significantly lower overall distances covered.<sup>14</sup> Whereas, external defenders (402+165 m), external midfielders (446+161 m) and forwards (404+140 m) have presented a noticeable difference in total sprinting distance during the duration of a game.<sup>19,14</sup> Central midfielders only recorded total sprinting distances of 248±116 m with centre backs presenting an even lower total sprinting distance (215±100 m).<sup>14</sup> The study by Di Salvo et al<sup>21</sup> supports centre backs low overall distances covered both anaerobically and aerobically resulting in their lower physical capacity and important tactical role.

Midfielders, through covering the largest distances are unable to work at high intensities and therefore sprint for the same duration as other positions. Covering a larger distance results in working at lower intensities with less opportunity for rest due to midfielder's HR rarely falling below 65% of HR<sub>max</sub>.<sup>14</sup> Therefore, the working muscles have a continuous demand for oxygen prohibiting anaerobic, high intensity sprinting. External defenders and forwards have further rest opportunities, allowing higher intensities to be met on numerous occasions, thus resulting in larger total sprinting distances.

### Positional Variables Affecting External Workload

Total external work completed throughout a game has been found to vary according to a variety of factors which dictate work rate profiles and energy expenditure.<sup>14</sup> Firstly, a player's VO<sub>2max</sub> can influence the distance performed; a higher physical capacity displays a positive correlation to enable further distance to be completed due to positional and tactical roles.<sup>14</sup> The variation between VO<sub>2max</sub> and positions influences the ability to cover greater distances at higher intensities; this is further reflected through the increased VO<sub>2max</sub> values consistently obtained by midfielders, alongside overall largest distances travelled.<sup>14</sup> Secondary factors affecting external workload include the style of play and team formation.<sup>22,20</sup> The evolution concerning style of play within professional football has been suggested by Di Mascio and Bradley,<sup>14</sup> whom produced conclusions which demonstrate an increased distance covered by contemporary English Premier League players (11 km) compared to those in the original First Division (8.2 km) observed by Reilly and Thomas.<sup>23</sup> The increase in 2.8 km supports the physical demands of football increasing with every position involved in defensive and offensive phases across a match.<sup>3,10</sup> Tierney et al<sup>22</sup> stated that there are noticeable differences in the positional demands across a series of formations. Jozak et al<sup>24</sup> claimed that midfielders will always cover the largest total distance in any formation, due to being the direct link between the defensive and offensive part of the whole team.

### Heart Rate of Footballers

Heart rate (HR) is a leading tool guiding the intensity level of exercise and determining physical demands.<sup>9,25</sup> The intensity level is

expressed as a percentage of maximal heart rate ( $HR_{max}$ ) due to the large variations that occur throughout 90 minutes within a match. Changes to the modern game<sup>3</sup> has led researchers into developing specific aerobic conditioning concepts for players.<sup>26</sup> Hoff et al<sup>27</sup> stated that pure running is suitable for players to reach maximum oxygen consumption. However, simply running is not suitable for the development of football specific aerobic capacity, which is more dependent upon aerobic peripheral, rather than central adaptation.<sup>26,28</sup> Within football, it is documented that the maximum benefits are achieved when the conditioning stimuli coincide with specific competitive demands.<sup>26,29</sup>

### Heart Rate Intensity in Football

In football, athletes perform different types of physical movements ranging at a series of intensities.<sup>1,8</sup> Santos et al<sup>1</sup> claimed that 80-90% of football spent within a low to moderate intensity zone with 10-20% spent completing high intensity activities. These findings do not match previous research by Jozaket al<sup>24</sup> whom presented 30% of overall football activities to be spent at a high intensity. In total, the average myocardium reaction throughout the duration of a 90-minute match is 80-90% of the  $HR_{max}$ .<sup>20,30</sup> Coelho et al<sup>30</sup> compared player positions based on intensity within football matches. The researchers concluded that midfielders spent the majority of a 90-minute football match at 85-90%  $HR_{max}$ , thus suggesting higher aerobic requirements than defenders and forwards. As a result, Coelho et al<sup>30</sup> suggested a higher requirement of aerobic activities within midfielders physical training to assist with recovery and consequently, their performance.

### Fatigue of Footballers

Fatigue has become a high profile topic in intermittent team sports.<sup>31</sup> Waldron and Highton<sup>31</sup> discussed the likelihood that performers whom inversely have an increase in overall work performed, must reflect recovery either partially or fully, overall displaying temporary fatigue. The phrase refers to a period of reduced intensity, which occurs directly after the most intense period which can occur through possession or the match score.<sup>31</sup> Therefore, if players are in possession of the ball, their intensity of play is reduced and external workload reduced, this period of recovery overcomes temporary fatigue. Acute fatigue, which is seen within footballers, occurs as a result of both peripheral and central factors.<sup>31</sup> The immediate peripheral assumption to fatigue is assumed to be due to blood lactate accumulation with a reduction in pH.<sup>31</sup> Waldron and

Highton<sup>31</sup> report blood lactate concentrations samples during and after football matches to vary between 7-9  $mmol^{-1}$ . The findings reveal that team sports players tolerate some degree of lactate accumulation; however, blood lactate accumulations do not appear to increase across the course of the matches.<sup>31</sup> Furthermore, whilst a decrease in pH has been reported after the most intense period within a match, a poor correlation has been connected to decrements in sprinting performance.<sup>31</sup> Overall, the evidence therefore concludes that other factors must contribute towards the reduction in work within football matches and other intermittent sports.

Football matches last 90-minutes, indicating that muscle glycogen is the primary source for ATP synthesis; however, glycogen stores are limited and depleted quickly.<sup>32</sup> Kenney et al<sup>32</sup> stated that since the muscle biopsy technique was introduced, studies show a clear correlation between muscle glycogen depletion and fatigue during prolonged activity. Football is an intermittent sport<sup>3</sup> with over 1000 high intensity actions,<sup>10,20</sup> therefore the working muscles constantly rely on a supply of glycogen to meet high energy demands.<sup>32</sup> As the match prolongs, the constant supply of glycogen slows down causing the muscles to fatigue.<sup>32</sup> This study aimed to: (i) To investigate which position completes the most external work across the varying heart rate intensities, (ii) To compare differences in external workload (distance travelled) completed by each position and their average heart rate intensity and, (iii) Compare the average heart rate intensity of female footballers per 45 minute half of a football match to investigate fatigue

## METHODS

### Participants and Recruitment

A sample of n=24 amateur female football players (mean±SD; age: 20.7±4.0 years; stretched stature: 165.6±5.8 cm; body mass: 58.1±4.7 kg) were recruited from an amateur football club within the South West of England during the 2016/2017 competitive season (Table 1). Recruited participants were outfield playing positions including defenders (n=8), midfielders (n=8) and forwards (n=8) and possessed at least 5 years of playing and training footballing experience. All participants were over 18 years of age, free from disease, illness or injury. They all completed a Physical Activity Readiness Questionnaire (PAR-Q) before inclusion<sup>33</sup> and provided written informed consent, and understood their right to withdraw. To minimise risk, ethical approval was sought from the University of Gloucestershire Faculty Research Ethics Committee.

**Table 1.** General summary ( $\bar{x} \pm s$ ) characteristics for n=24 amateur female football players according to playing position (defenders, midfielders and forwards)

Variables	Defenders (n=8) $\bar{x} \pm s$	Midfielders (n=8) $\bar{x} \pm s$	Forwards (n=8) $\bar{x} \pm s$
Age (years)	22.4±5.5	20.9±3.5	20.7±2.1
Stretched Stature (cm)	164.6±2.2	168.3±7.4	163.8±6.1
Body Mass (kg)	59.1±3.7	59.1±3.7	59.1±3.7

## Procedures


All participants followed the same protocols leading up to each test to ensure consistency. In the 24 hour period prior to testing participants did not partake in any vigorous exercise and refrain from consuming any caffeine or alcohol.<sup>34</sup> Furthermore, participants arrived hydrated and having not consume large quantities of food within a 2 hour prior to testing. Body Mass was measured in the morning with electronic digital weighing scales and to the nearest 0.1 kg. Stretched stature was also measured in the morning using a Harpenden Stadiometer to the nearest 0.1 cm. After measurements were taken, the mean value was determined for subsequent data analysis. The Yo-Yo IR Level 1 protocol commenced whilst all players entered the middle of the competitive season to determine the maximal HR values for each participant. The week following, the SAFT<sup>90</sup> procedure was completed.<sup>35</sup> The training load and amount of match play performed for each participant was standardised to the competitive season.

## Experimental Session 1

**(Yo-Yo Level 1 Intermittent Recovery Test):** The Yo-Yo IR Level 1 test was performed on artificial turf in running lanes having a width of 2 m and a length of 20 m.<sup>36,37</sup> Prior to the test, the participants performed a warm up prescribed by Barrett et al<sup>38</sup> and Marshall et al<sup>39</sup> an identical warm up was used prior to the completion of the SAFT<sup>90</sup>. Procedures suggested by Krusturpet al<sup>40</sup> and Castagna et al<sup>41</sup> for the Yo-Yo IR Level 1 were followed, with runs at a progressive speed, controlled by audio bleeps from a pre-recorded tape.<sup>36</sup> After each bout of running, the players had a 10s rest period.<sup>36</sup> The test result was recorded upon the point when a participant has failed twice to reach to finish line within the allocated time.<sup>36</sup>

## Experimental Session 2

**(Soccer-Specific Aerobic Field Test (SAFT<sup>90</sup>)):** The SAFT<sup>90</sup> design uses an agility based course whereby players navigate around the 20 m course in an intermittent fashion through stranding (0 km/h), walking (~5.5 km/h), jogging (~10.7 km/h), striding (~15 km/h) or sprinting (maximal effort).<sup>39,42,43,39</sup> The course is based around a shuttle run with the incorporation of 4 positioned poles in which the participants must navigate around (Figure 1).<sup>38,42,43</sup> Altogether, the course provides the participants with 1332 changes in direction and 1269 changes in speed, eliciting internal loads similar to those reported from match-play.<sup>39,38,42,43</sup>



Prior to the completion of the test, the participants performed a football specific warm up prescribed by Barrett et al<sup>38</sup> and

Marshall et al.<sup>39</sup> The first 8 exercises proposed by Marshall et al<sup>39</sup> involved light jogging further to side-stepping with the inclusion of simple plyometric movements. Dynamic movements followed with each sequence performed twice over a 20 m distance.<sup>39</sup> Final exercises involved sprint-based agility movements before undertaking the SAFT<sup>90</sup>.<sup>39</sup>

## Data Collection Session

**Heart Rate:** Heart rates were monitored throughout the Yo-Yo IR Level 1 and SAFT<sup>90</sup> protocols using a HR monitoring system (Polar Team 2 System, Kempele, Finland).<sup>35</sup> Furthermore, the data collected through the completion of the Yo-Yo IR Level 1 quantified the overall HR intensity zones. The intensity of the match was reported as the percentage of time spent in the five HR zones prescribed by Coelho et al<sup>30</sup>. Each zone was identified as a different percentage of HR<sub>max</sub>: zone 1 equates to <70% HR<sub>max</sub>, zone 2, 70-85% HR<sub>max</sub>; zone 3, 85-90 HR<sub>max</sub>; zone 4, 90-95% HR<sub>max</sub>, with zone 5, 95-100% HR<sub>max</sub>.<sup>30</sup> Heart rates for each participant were recorded every second of each protocol and quantified to produce data for each minute throughout each 45 minute half and across the 90-minute duration with each position being investigated. The amount of time, alongside the percentage of time spent in each zone was reported in relation to the amount of external work completed for each position. Additionally, differences in HR recordings between positions and across the duration of the protocol (per 45 minute half) was measured, to investigate fatigue.

**External Workload:** Throughout both protocols, in order to obtain overall external workloads, the participants were videoed replicating protocols previously implemented by De Ste Croix et al.<sup>43</sup> The footage attained was then interpreted based on the standardised 20 m guidelines.

## Statistical Analysis

Raw data sets were analysed using Microsoft Excel, 2016. Descriptive statistics, including means and standard deviations, were calculation for each measure including, external workload and HR intensities across each 45 minute half. Furthermore, percentage measurements for each position, based on each HR intensity, were calculated. In order to determine the following situations: a) comparison of intensity between the different halves of the match (first half *vs.* second half); b) comparison of each position across the intensity zones; c) comparison of each positions external workload; d) time spent at each intensity across each 45 minute half; a one-way analysis of variance (ANOVA) was applied. The level of significance for statistical testing was set at alpha level  $p < 0.05$ .

## RESULTS

On average, the overall external workload for all positions, during the SAFT<sup>90</sup>, was (mean±SD) 10913.7±1076.7 m. Across the 90-minutes, defenders recorded an external workload of 10020.4±1086.6 m, whilst forwards recorded a distance of 10938.9±865.2 m, and midfielders an external workload of 11781.9±324.7 m. Therefore,

**Table 1.** Total external workload performed during SAFT<sup>90</sup> for each position (p) including defenders (d), midfielders (m) and forwards (f).

P	1 <sup>st</sup> Half (m)		2 <sup>nd</sup> Half (m)		Full Game (m)	
	Mean±SD	Range	Mean±SD	Range	Mean±SD	Range
D	5774.6±783.0	4969-6987	4245.8±856.3	3263-5659	10020.4±1086.6	8562-11984
M	6325.3±645.4	5236-6984	5456.6±635.3	4362-6549	11781.9±324.7	11236-12142
F	6325.3±645.4	5002-6954	4935.1±633.2	4071-5762	10938.9±865.2	9412-11985
ALL	6034.5±727.8	4969-6987	4879.2±851.3	3263-6549	10913.7±1076.7	8562-12142

**Table 2.** Mean heart rate (HR) during SAFT<sup>90</sup> for each position (p) including defenders (d), midfielders (m) and forwards (f).

P	1 <sup>st</sup> Half (bpm <sup>-1</sup> )		2 <sup>nd</sup> Half (bpm <sup>-1</sup> )		Full Game HR (bpm <sup>-1</sup> )	
	Mean±SD	Range	Mean±SD	Range	Mean±SD	Range
D	140.4±6.7	130.0-148.0	148.1±9.4	136.0-160.0	144.3±7.4	136.0-154.0
M	181.0±13.3	146.0-183.0	170.0±9.2	157.0-180.0	168.8±10.8	152.0-181.5
F	170.1±7.5	146.0-183.0	170.4±9.4	154.0-183.0	170.3±7.4	159.0-184.0
ALL	159.4±16.6	130.0-185.0	162.8±13.9	136.0-183.0	161.1±14.7	136.0-184.0

across the 90-minute SAFT<sup>90</sup>, midfielders overall performed the largest proportion of external work. Across each 45 minute halves of the SAFT<sup>90</sup>, midfielders consistently recorded the highest external workload. Across the opening 45 minutes, midfielders recorded distances of 6325.3±645.4 m, with forwards documenting distances of 6003.8±32.2 m and defenders, the lowest of the three outfield positions, 5774.6±783.0 m. This trend continued into the second half of the protocol, with midfielders maintaining the highest external workload (5456.6±635.3 m), with forwards distances of 4935.1±633.2 m and repeatedly, defenders the lowest of the three positions (4245.8±856.3 m). Furthermore, between the two separate halves, all players demonstrated a decreased external workload in the second half (6034.5±727.8 m) in comparison with the first 45 minutes completed (4879.2±851.3 m). A one-way ANOVA showed that the effect of position within female football on the external workload performed during a 90-minute SAFT<sup>90</sup> was significant (F(df:23)=9.156; p<0.05). However, across the two separate 45 minute halves, significance results differed. Within the first half, the effect of position on the external workload performed was not significant, whereas, the second half showed there to be a significant difference between the playing positions and overall external workload completed (F(df:23)=5.758; p< 0.05).

Based on the results illustrated in Table 1, defenders present the lowest average HR (144.3±7.4 bpm<sup>-1</sup> in comparison with forwards (170.3±7.4 bpm<sup>-1</sup> and midfielders (168.8±10.8 bpm<sup>-1</sup>) across the 90-minutes of recorded data. This trend is further present between the two 45 minute halves. Across the first 45 minutes, defenders recorded an average HR of 140.4±6.7 bpm<sup>-1</sup> significantly lower than midfielders (181.0±13.3 bpm<sup>-1</sup> and forwards (170.1±7.5 bpm<sup>-1</sup>). In the second half, forwards displayed the highest average HR of 170.4±9.4 bpm<sup>-1</sup> in comparison with midfielders (170.0±9.2 bpm<sup>-1</sup>) and forwards (148.1±9.4 bpm<sup>-1</sup>). Across the two halves of the 90 minutes, defenders and forwards each presented average HR data showing an increase, from the first half to the second, midfielders however, demonstrated a decreased. The average HR recorded for all positions across the SAFT<sup>90</sup> was 161.1±14.7


bpm<sup>-1</sup> as shown in Table 2, with the average intensity during the protocol for all defenders, midfielder and forward represented 81% HR<sub>max</sub>.

Figure 2 and Figure 3 demonstrate that midfielders and defenders continually displayed lower average heart rates than forwards per 15-minute bout of the SAFT<sup>90</sup>. Within the opening 15 minute bout (Figure 2), all three positions present an increase in HR, following into the second bout (15-30 minutes), only at the 23<sup>rd</sup> minute did each positions HR plateau. Figure 1 illustrates average heart rates from midfielders and forwards maintaining 150-200 bpm<sup>-1</sup>, however defenders continually work between 100-150 bpm<sup>-1</sup>. Defenders further display periods of increased and decreased heart rates with decreases falling in both the 35<sup>th</sup>-37<sup>th</sup> minute and 41<sup>st</sup>-43<sup>rd</sup> minute. Towards the end of the opening half of the SAFT<sup>90</sup>, both midfielders and defenders show a decrease in average HR.


Across the entire bout, defenders displayed much lower heart rates showing a range between 135-160 bpm<sup>-1</sup> in comparison with midfielders at 160-180 bpm<sup>-1</sup> and forwards at 170-190 bpm<sup>-1</sup>. The final bout displayed both the midfielders and forwards HR decreasing in comparison with the previous 15-minutes (Figure 3). Similarly, defenders showed periods an increased HR of 160 bpm<sup>-1</sup> at 77 minutes and 165 bpm<sup>-1</sup> at 85 minutes, with periods of a decreased HR of 140 bpm<sup>-1</sup> at 80 minutes and then 87 minutes. However, the final 3 minutes of the entire protocol presents all three positions average heart rates decreasing.

A one way ANOVA showed the effect of position, within female football, on the average HR during a 90 minute SAFT<sup>90</sup> was significant (F(df:23 )=22.317; p<0.05). Furthermore, results showed a significant difference between defenders, midfielders and forwards average HR across both the first 45 minute half (F(df:23)=23.405; p<0.05) and second 45 minute half (F(df:23)=14.925; p<0.05).

**Figure 2.** Average HR for defenders, midfielders and forwards per 15-minute bout of the first 45 minute half of the SAFT<sup>90</sup> including 1-15 minutes, 15-30 minutes and 30-45 minutes



**Figure 3.** Average HR for defenders, midfielders and forwards per 15 minute bout of the second 45 minute half of the SAFT<sup>90</sup> including 45-60 minutes, 60-75 minutes and 75-90 minutes




### Heart Rate Intensities

Across the total SAFT<sup>90</sup>, the average intensity for all positions represents 81% HR<sub>max</sub>, which is in line with the investigation means within zone 2 (70-85% HR<sub>max</sub>). Within the first half, the average intensity for all positions represents 80% HR<sub>max</sub>; however, in the second half, average intensity increased to 82% HR<sub>max</sub> but was maintained in zone 2. Through the completion of a one way ANOVA, significance results showed position, to have an effect on the level of intensity and time spent at each intensity. Across the SAFT<sup>90</sup>, there was a significant difference between playing position and the duration spent working within zone 1, <70% HR<sub>max</sub> (F(df:23)=5.271; p<0.05); zone 2 between 70-85% HR<sub>max</sub> (F(df:23)=5.965; p<0.05) and zone 4 at 90-95% HR<sub>max</sub> (F(df:23)=4.433; p<0.05). However, across zone 3 between 85-90% HR<sub>max</sub> and zone 5 at 95-100% HR<sub>max</sub>, no significant difference was found between each position and the duration spent working at each intensity. Within the opening half, defenders average intensity was 73% HR<sub>max</sub>, working within zone 2 (70-85% HR<sub>max</sub>). However, midfielders and forwards average intensity represented that of zone 3 (85-90% HR<sub>max</sub>), with

midfielder's average intensity at 88% HR<sub>max</sub> and forwards 86% HR<sub>max</sub>. In comparison with the second half, forwards display no change in average heart rate intensity. Midfielders however moved into zone 2 (70-85% HR<sub>max</sub>) at 83% HR<sub>max</sub>. Defenders nonetheless display an increase in average heart rate intensity to 78% HR<sub>max</sub>; however, still maintained working within zone 2 (70-85% HR<sub>max</sub>).

Across the 90-minutes (Figure 3), all three positions spent time in each HR zone, from zone 1 to zone 5. All positions each spent the largest percentage of the SAFT<sup>90</sup> in zone 2 (70-85% HR<sub>max</sub>). Specifically, within zone 1 (<70% HR<sub>max</sub>), defenders spent the larger percentage of time with forwards the least, whereas, at the highest intensity, in zone 5 (95-100% HR<sub>max</sub>), forwards spent the larger percentage of time with defender's the least. Midfielders specifically spent the largest percentage of the 90-minutes within zone 3 (85-90% HR<sub>max</sub>) and zone 4 (90-95% HR<sub>max</sub>) in comparison with the further outfield positions.

Figure 3 displays differences between the HR intensities between the two 45-minute halves of the SAFT<sup>90</sup>. Within the first



half, all three positions worked across each HR zone; however, in the second half, defenders never worked with in zone 5 (95-100% HR<sub>max</sub>). Similarly, between the first half and second half, all three positions spent the largest percentage of the game at 70-85% HR<sub>max</sub> in zone 2. Specifically, across both halves of the protocol, midfielder's largest duration of time was spent within zone 2 and 3 working in total across 70-90% HR<sub>max</sub> whereas defender's largest percentage of time was spent within zone 1 and 2. Forwards in comparison spent the smallest percentage of time within zone 1; however, increasingly more from zones 2 to zone 5 with HR intensities varying from 70-100% HR<sub>max</sub>.

Significance results of a completed one-way ANOVA showed that position has an effect on the level of intensity and time spent at each intensity. Across the first 45-minute half of the SAFT<sup>90</sup>, there was a significant difference between playing position and the duration spent working within zone 1 (<70% HR<sub>max</sub>, (F(df:23)=6.379; *p*<0.05); zone 2 between 70-85% HR<sub>max</sub> (F(df:23)=5.076; *p*<0.05); zone 3 (85-90% HR<sub>max</sub>) (F(df:23)=4.997; *p*<0.05) and zone 4 (90-95% HR<sub>max</sub>) (F(df:23)=4.635; *p*<0.05). However, no significant difference was found between position and duration working at the intensity of zone 5 (95-100% HR<sub>max</sub>). Across the second 45-minute half, no significant difference was found between each position, for the duration of time spent within, zone 1 (<70% HR<sub>max</sub>), zone 3 (85-90% HR<sub>max</sub>) further to zone 4 (90-95% HR<sub>max</sub>) and zone 5 (95-100% HR<sub>max</sub>). Within the second 45-minutes, significant differences were found between all three positions and the duration of time spent within zone 2 (70-85% HR<sub>max</sub>), (F(df:23)=4.580; *p*<0.05).

## CONCLUSION

### External Workload

In total, for all three positions, the total external workload complet-

ed across 90-minutes was 10913.7±1076.7 m with a range of 8562-12142 m. The findings correspond to previous research showing the average distance players cover during a match is 8-12 km.<sup>16,7,20</sup> Statistical analysis across the 90-minutes concluded a significant difference between the workloads completed between the three positions. On average, the largest external workload completed was by midfielders (11781.9±324.7 m) followed by forwards with defenders covering the smallest distance covered (10020.4±1086.6 m). Other studies have also found midfielders to cover the greatest external workload.<sup>3,10,44</sup> The completed distance for midfielders (11781.9±324.7 m) further corresponds to that of previous studies within the range of 10-12 km.<sup>10,44</sup>

The protocol removes oppositional effects and tactics, resulting in the physiological and mechanical demands displayed during football match-play being observed; therefore, it can be concluded that the greater distances were not due to tactics and styles of play.<sup>34,42</sup> Midfielders however, to have developed higher physical capacities or VO<sub>2max</sub> levels due to their overall roles within match play, through involvement in both attacking and defensive phases of play in comparison with their counterparts, who rarely cross or retreat past the half way line.<sup>45</sup> Therefore, with an increased physical capacity, more external work is able to be completed. However, with regards to total external workload, other variables further to VO<sub>2max</sub> may have influenced the results. These variables include changes in the intensity of activity as well as motivation.<sup>8</sup> Between the three positions, defenders covered the lowest external workload (10020.4±1086.6 m). Indeed, several studies have previously found defenders to obtain, in general, lower fitness levels, resulting in lower external loads performed in comparison with midfielders and forwards.<sup>20,44</sup> Research by Di Salvo et al<sup>21</sup> supports defender's low overall distances covered both anaerobically and aerobically as the result of their lower physical capacity and tactical role to complete small bouts of movement in winning duels and aerial challenges.<sup>24</sup>



## Heart Rate

Considering the load intensity from a physiological point of view, average HR results across the 90-minute SAFT<sup>90</sup> for all players was  $161 \pm 14.7 \text{ bpm}^{-1}$  corresponding to  $81 \pm 7.4\% \text{ HR}_{\text{max}}$ . This matched heart rates previously obtained within research by Andersson et al<sup>44</sup> who found the average HR of female footballers within domesticated matches to correspond to  $162 \pm 6 \text{ bpm}^{-1}$  or  $85 \pm 3\%$  of  $\text{HR}_{\text{max}}$ . However, the participants of the study were international female players from Sweden and Denmark, playing professionally, in comparison with amateur footballers.

The football simulated match intensity found in the present study did not differ markedly from those reported in other investigations. Coelho et al<sup>30</sup> evaluation of 26 footballers, during a championship game, reported an overall intensity of  $84\% \text{ HR}_{\text{max}}$ . Furthermore, Mohr et al<sup>46</sup> reported an intensity of  $85\% \text{ HR}_{\text{max}}$  during a friendly match of the 4<sup>th</sup> division, whilst O'Connor<sup>47</sup> evaluated HR during two women's and men's soccer games finding an intensity of  $85\% \text{ HR}_{\text{max}}$ . However, both studies evaluated friendly matches rather than a simulated football match. Intensity observed in the present study further matches than that obtained by Reilly and Keane,<sup>48</sup> who evaluated senior soccer players in a specific simulated test with the mean intensity presented at  $80\% \text{ HR}_{\text{max}}$ . Although maximum effort tests are suitable for the determination of  $\text{HR}_{\text{max}}$ , a previous study by Antonacci et al<sup>49</sup> showed that  $\text{HR}_{\text{max}}$  is lower in simulated tests, than during official matches.

## Heart Rate Intensities

Focusing on the total 90 minute SAFT<sup>90</sup>, the average intensity for all positions was  $81\% \text{ HR}_{\text{max}}$  which in line with the investigation means within zone 2 ( $70\text{-}85\% \text{ HR}_{\text{max}}$ ). The data, as a result, corresponds with other studies findings.<sup>30,44,46</sup> Throughout the first half, the average intensity for all positions was  $80\% \text{ HR}_{\text{max}}$ ; however, within the second half the average intensity increased to  $82\% \text{ HR}_{\text{max}}$ . This meant across both 45-minute halves working within zone 2 ( $70\text{-}85\% \text{ HR}_{\text{max}}$ ). However, the results contradict the research conducted by Coelho et al<sup>30</sup> whom observed a decrease in game intensity from the first to second half of a championship football game. Mortimer et al<sup>50</sup> stated that a reduction in game intensity occurs during the second half of a football match, due to the role of fatigue as a result of the progressive utilisation of glycogen, which decreases during the second half. With focus on each of the position, forwards average HR did not differ between either of the halves of football and maintained  $86\%$  within zone 3 ( $85\text{-}90\% \text{ HR}_{\text{max}}$ ), whereas defenders rose from the first half ( $73\% \text{ HR}_{\text{max}}$ ) to the second half ( $78\% \text{ HR}_{\text{max}}$ ). Midfielders average HR from the first half ( $88\% \text{ HR}_{\text{max}}$ ) in comparison to the second half ( $83\% \text{ HR}_{\text{max}}$ ) was the only decrease supporting previous research. A similar decline had been reported in other studies.<sup>30,46</sup> The lack of significant difference between the two halves of the game for forwards is however supported by Vencurik et al<sup>51</sup> who witnessed no changes between certain positions physiological demands across the two halves of a female basketball match.

Across the 90-minutes, results displayed differences between each of the HR intensity zones, however significant differences were only found between the duration each position spent within zone 1, zone 2 and zone 4. Nonetheless, across the 90-minutes, each position spent time within each zone. All three positions each spent the largest percentage of the SAFT<sup>90</sup> within zone 2 ( $70\text{-}85\% \text{ HR}_{\text{max}}$ ) in comparison with the other intensity zones. Defenders spent the largest percentage of time within zone 1, with forwards the least, whereas, at the high intensities, in zone 5, forwards spent the larger percentage of time with defenders the least. This result demonstrates the importance of HIR for forwards and LIT for defenders. Di Mascio and Bradley,<sup>14</sup> analysed the evolution of football describing forwards movements to focus around high intensities and short-duration efforts, with defenders maintaining a low intensity with minimal HIR, therefore, the current study backs up that of previous research.<sup>10,4</sup>

The results however, contradict the research by Jozaket al<sup>24</sup> who studied 600 players during the world cup finals; results showed defenders and forwards to have completed the most LIR due to the positions having the least contact with the ball. However, the comparison must be taken with caution due to the sample using male international athletes during competitive matches, resulting in formations and tactics potentially interfering with play.<sup>22,20</sup> A further finding includes midfielders specifically spending the largest percentage of the 90-minutes within zone 3 and zone 4. Research by Coelho et al<sup>30</sup> and Goncalves et al<sup>25</sup> reported the similar findings. Furthermore, Orendurff et al<sup>52</sup> investigated differences in positions during a game using a step rate as an intensity parameter; the results obtained were similar to those observed in both the current study but further the investigation by Coelho et al.<sup>30</sup> Midfielders presented a similar game pattern characterised by few maximum intensity bouts (step rate=7) and short recovery bouts of moderately high intensity (step rate=4), thus in agreement with studies showing a higher aerobic capacity of midfielders.<sup>30,45</sup>

## First Half Versus Second Half Fatigue

It is generally accepted that physical performance declines during a match because of increased fatigue towards the end of a game.<sup>36,42</sup> Within the current study, first half differences in comparison second half suggest all positions to experience fatigue. External workload decreased for all positions by  $11\%$  with a first half overall workload of  $6034.5 \pm 727.8 \text{ m}$  in comparison to a second half workload of  $4879.2 \pm 851.3 \text{ m}$ . Furthermore, defenders, midfielders and forwards individual workloads completed each decreased. The findings further support numerous research that indicates that the fatigue induced by the physical activity completed during the first half is related to the physical activity completed during the second.<sup>4,19,44</sup>

Within the first half, all three positions worked across each HR zone; however, in the second half defenders never worked with in zone 5 displaying a slowly declining intensity. Furthermore, the percentage of time defenders spent within zone 1 and zone 2 increase in the second half. Additionally, midfielder's

intensity decreased at the highest intensity (95-100% HR<sub>max</sub>) by 2%, further to forwards by 4%. Moreover, forwards time within zone 2 and zone 3 and zone 4 increased within the second half in comparison with the first. Specifically, forwards duration of time spent in zone 3 increased by 2% and in zone 4, by 3%. Martinez-Lagunas et al<sup>4</sup> stated that the amount of HIR completed in the second half, compared to the first half of match-play, decreases due to temporary and permanent forms of fatigue. Since football game intensity and fatigue is influenced by the opponent and tactical preferences of the coach, the data should be considered carefully due to using a SAFT<sup>90</sup>. Nonetheless, Jones et al<sup>42</sup> stated that the fatigue induced by the SAFT<sup>90</sup> protocol is representative of the responses associated with actual football match-play.<sup>53</sup>

In summary, overall external distance during a SAFT<sup>90</sup> is similar to that of competitive match-play. The results indicated that all positions workloads externally are different, with midfielders achieving the highest overall distance, in comparison with forwards and defenders, who completed the least. Defenders spent a higher percentage of the protocol within the lowest intensity zones with forwards rarely performing below zone 2 (70-85% HR<sub>max</sub>) resulting in forwards spending the largest percentage of the game at the highest intensities. This data should be included in specific training programmes addressing the tactical positions of players. In the case of forwards, the objective is to increase performance at short-durations and at the highest intensities, since their main focus is to move at pace to create space to receive the ball or break through oppositions defensive lines. Midfielders are the players who spent the most time between zone 2 (70-85% HR<sub>max</sub>), zone 3 (85-90% HR<sub>max</sub>) and zone 4 (90-95% HR<sub>max</sub>). In addition, across both halves and the total 90-minutes, they spent a higher percentage of time in zone 4 (90-95% HR<sub>max</sub>) than defenders and forwards, suggesting higher aerobic requirements of these players. This higher requirement of aerobic activities should be included in the physical training of midfielders to improve their recovery and consequently their performance.

## REFERENCES

- Santos R, Sousa C, Olher R, Neto I, Pereira L, Sales M. Comparison of the anaerobic power of Brazilian Professional Football Players Grouped by tactical position. *J Sports Sci Med*. 2014; 3:13-18.
- Marshall P, Lovell R, Jeppesen G, Kristoffer A, Sieger J. Hamstring muscle fatigue and central motor output during a simulated soccer match. *PLoS One*. 2014; 9(7): 1027-53. doi:[10.1371/journal.pone.0102753](https://doi.org/10.1371/journal.pone.0102753)
- Ade J, Fitzpatrick J, Bradley P. High-intensity efforts in elite soccer matches and associated movement patterns, technical skills and tactical actions. *J Sports Sci*. 2016; 34: 2205-2214.
- Martinez-Lagunas V, Niessan M, Hartmann U. Women's football: Characteristics and demands of the game. *J Sport Health Sci*. 2014; 3: 258-272. doi: [10.1016/j.jshs.2014.10.001](https://doi.org/10.1016/j.jshs.2014.10.001)
- Selmi MA, Sassi H, Yahmed H, Moalla W, Elloumi M. Effect of between-set recovery durations on repeated sprint ability in young soccer players. *Biol Sport*. 33(2): 165-172. doi: [10.5604/20831862](https://doi.org/10.5604/20831862)
- Botek M, Krejčí J, Weisser R. Autonomic cardiac regulation and morpho-physiological responses to eight week training preparation in junior soccer players. *Acta Gymnica*. 2014; 44(3): 155-163. doi: [10.5507/ag.2014.016](https://doi.org/10.5507/ag.2014.016)
- Andrzejewski M, Chmura J, Dybek T, Pluta B. Sport exercise capacity of soccer players at different levels of performance. *Biol Sport*. 2012; 29(3): 185-191.
- Belka J, Hulka K, Safar M, Weisser R. External and internal load on playing positions of elite female handball players (U19) during competitive matches. *Acta Gymnica*. 2016; 46(1): 12-20. doi: [10.5507/ag.2015.025](https://doi.org/10.5507/ag.2015.025)
- Galy O, Zoubir B, Hambli M, Chaouachi A, Hue O, Chamari K. Relationships between heart rate and physiological parameters of performance in top level water polo players. *Biol Sport*. 2014; 31: 33-38.
- Ademovic A. Differences in the quantity and intensity of playing in elite soccer players of different positions in the game. *Homo Sporticus*, 2016; 18: 26-31.
- Bojkowski L, Śliwowski R, Wieczorek A, Eider J. Analysis of the longest distances run by the best soccer players at The Fifa World Cup in Brazil in 2014. *European Journal of Sport Sciences and Medicine*. 2015; 11(3):145-151. doi: [10.18276/cej.2015.3-15](https://doi.org/10.18276/cej.2015.3-15)
- Nilsson J, Cardinale D. Aerobic and anaerobic test performance among elite male football players in different team positions. *Lase Journal of Sport Science*. 2015; 6(2): 73-92. doi: [10.1515/ljss-2016-0007](https://doi.org/10.1515/ljss-2016-0007)
- Bangsbo J, Mohr M, Krstrup P. Physical and metabolic demands of training and match-play in the elite football player. *J Sports Sci*. 2006; 24(7): 665-674. doi: [10.1080/02640410500482529](https://doi.org/10.1080/02640410500482529)
- Di Mascio M, Bradley P. (2013). Evaluation of the most intense high-intensity running period in english FA premier league soccer matches. *J Strength Cond Res*. 2013. 27(4), 909-915. doi: [10.1519/JSC.0b013e31825ff099](https://doi.org/10.1519/JSC.0b013e31825ff099)
- Andersen T, Bendiksen M, Pedersen J, et al. Kicking velocity and physical, technical, tactical match performance for U18 female football playerseffect of a new ball. *Hum Mov Sci*. 2012; 31(6): 1624-1638. doi: [10.1016/j.humov.2012.07.003](https://doi.org/10.1016/j.humov.2012.07.003)
- Dallaway N. Movement profile monitoring in professional football. [Unpublished master's thesis]. Birmingham, UK: University of Birmingham; 2013.
- Hoff J, Helgerud J. Endurance and strength training for soccer players: Physiological Considerations. *Sports Med*. 2004; 34(3): 165-180.

18. Shalfaw S, Ingebrigtsen J, Dillern T, Tonnessen E, Kristofer T, Enoksen E. The effect of 40m repeated sprint training on physical performance in young elite male soccer players. *Serbian Journal of Sports Sciences*. 2012; 6(3): 111-116. doi: [10.1519/JSC.0b013e3182023a65](https://doi.org/10.1519/JSC.0b013e3182023a65)
19. Bradley PS, Bendiksen M, Dellal A, et al. The application of the Yo-Yo intermittent endurance level 2 test to elite female soccer populations. *Scand J Med Sci Sport*. 2014; 24(1): 43-54. doi: [10.1111/j.1600-0838.2012.01483](https://doi.org/10.1111/j.1600-0838.2012.01483)
20. Andrzejewski M, Chmura J, Pluta B, Konarski J. Sprinting activities and distance covered by top level Europa league soccer players. *Internal Journal of Sports Science and Coaching*. 2015; 10(1): 39-50. doi: [10.1260/1747-9541.10.1.39](https://doi.org/10.1260/1747-9541.10.1.39)
21. Di Salvo V, Tschan H, Bachl N. Performance characteristics according to playing position in elite soccer. *Int J Sports Med*. 2007; 28(3): 222-227. doi: [10.1260/1747-9541.10.1.39](https://doi.org/10.1260/1747-9541.10.1.39)
22. Tierney P, Young A, Clarke N, Duncan M. Match play demands of 11 versus 11 professional football using global positioning system tracking: Variations Across Common Formations. *Hum Mov Sci*. 2016; 49: 1-8. doi: [10.1016/j.humov.2016.05.007](https://doi.org/10.1016/j.humov.2016.05.007)
23. Reilly T, Thomas V. A motion analysis of work-rate in different positional roles in professional football match-play. *Journal of Human Movement Studies*. 1976; 2(2): 87-97.
24. Jozak R, Peric A, Bradic A, Dizdar D. Position-related differences in the amount, intensity and speed of movement in elite football players. *Homo Sporticus*. 2011; 13(2): 16-22.
25. Goncalves B, Figueira B, Macas V, Sampaio J. Effect of player position on movement behaviour, physical and physiological performance during an 11-a-side football game. *J Sports Sci*. 2014; 32(2): 191-199. doi: [10.1080/02640414.2013.816761](https://doi.org/10.1080/02640414.2013.816761)
26. Hulka K, Weisser R, Belka J. Effect of the pitch size and presence of goalkeepers on the workload of players during small-sided soccer games. *J Hum Kinet*. 2016; 51(1): 175-181. doi: [10.1515/hukin-2015-0180](https://doi.org/10.1515/hukin-2015-0180).
27. Hoff J, Wisloff U, Engen L, Kemi O, Helgerud J. Soccer specific aerobic endurance training. *J Sports Med*. 2002; 36(3): 218-221. doi: [10.1136/bjism.36.3.218](https://doi.org/10.1136/bjism.36.3.218)
28. Gamble P. Challenges and games-related solutions to metabolic conditioning for team sports. *Strength and Conditioning Journal*. 2007; 29(4): 60-65. doi: [10.1519/00126548-200708000-00010](https://doi.org/10.1519/00126548-200708000-00010)
29. Campos-Vazquez M, Mendez-Villanueva A, Gonzalez-Jurado J, Leon-Prados J, Santalla A, Suarez-Arrones L. Relationships between rating of perceived exertion and heart rate deprived internal training load in professional soccer players. *Int J Sports Physiol Perform*. 2015; 10(5): 587-592. doi: [10.1123/ijspp.2014-0294](https://doi.org/10.1123/ijspp.2014-0294)
30. Coelho D, Mortimer L, Condessa L, et al. Intensity of real competitive soccer matches and differences among player positions. *Artigo Original*. 2011; 13(5): 341-347. doi: [10.5007/1980-0037.2011v13n5p341](https://doi.org/10.5007/1980-0037.2011v13n5p341)
31. Waldron M, Highton J. Fatigue and pacing in high-intensity intermittent team sport: An update. *Sports Med*. 2014; 44(12): 1645-1658. doi: [10.1007/s40279-014-0230-6](https://doi.org/10.1007/s40279-014-0230-6)
32. Kenney WL, Wilmore JH, Costil DL. *Physiology of Sport and Exercise*. IL, USA: Human Kinetics. 2012.
33. Taskin H, Erkmen N, Cicioglu I. The examination of the heart rate recovery after anaerobic running in soccer players. *Coll Antropol*. 2014. 38(1): 207-211.
34. Azidin R, Sankey S, Drust B, Robinson M, Vanrenterghem J. Effects of treadmill versus overground soccer match simulations on biomechanical markers of anterior cruciate ligament injury risk in side cutting. *J Sports Sci*. 2015; 33(1): 1332-1341. doi: [10.1080/02640414.2014.990491](https://doi.org/10.1080/02640414.2014.990491)
35. Deprez D, Franssen J, Lenior M, Philippaerts R, Vaeyens R. The Yo-Yo intermittent recovery test 1 is reliable in young high-level soccer players. *Biol Sport*. 2015; 32(1): 65-70. doi: [10.5604/20831862.1127284](https://doi.org/10.5604/20831862.1127284).
36. Mohr M, Krustup P. Yo-Yo intermittent recovery test performances within an entire football league during a full season. *J Sports Sci*. 32(4): 315-327 doi: [10.1080/02640414.2013.824598](https://doi.org/10.1080/02640414.2013.824598)
37. Bradley O, Bendiksen M, Dellal A, Mohr M, Wilkie A, Datson, N. The application of the Yo-Yo intermittent endurance Level 2 test to elite female soccer populations. *Scand J Med Sci Sports*. 2012; 24(1): 43-54. doi: [10.1111/j.1600-0838.2012.01483](https://doi.org/10.1111/j.1600-0838.2012.01483)
38. Barrett S, Midgley A, Towson C, Garrett A, Portas M, Lovell R. Within-match-play load patterns during a simulated soccer match. *Int J Sports Physiol Perform*. 2015; 10: 135-140. doi: [10.1123/ijspp.2014-0582](https://doi.org/10.1123/ijspp.2014-0582).
39. Marshall P, Lovell R, Siegler J. Changes in passive tension of the hamstring muscles during a simulated soccer match. *Int J Sports Physiol Perform*. 2016; 11(5): 594-601. doi: [10.1123/ijspp.2015-0009](https://doi.org/10.1123/ijspp.2015-0009)
40. Krustup P, Mohr M, Amstrup T, Rysgaard T, Johansen J, Steensberg A, Bangsbo J. The Yo-Yo intermittent recovery test: Physiological response, reliability, and validity. *Med Sci Sports Exerc*. 2003; 35: 697-705. doi: [10.1249/01.MSS.0000058441.94520.32](https://doi.org/10.1249/01.MSS.0000058441.94520.32)
41. Castagna C, Impellizzeri F, Chamari K, Carlomagno D, Rampinini E. Aerobic fitness and Yo-Yo continuous and intermittent test performances in soccer players: A correlation study. *J Strength Cond Res*. 2006; 20(2): 320-325. doi: [10.1519/R-18065.1](https://doi.org/10.1519/R-18065.1)
42. Jones R, Bennett R, Todd A. Muscle fatigue induced by a soccer

- er match play simulation in amateur black South African players. *J Sports Sci.* 2015; 33(12): 1305-1311. doi:[10.1080/02640414.2015.1022572](https://doi.org/10.1080/02640414.2015.1022572)
43. De Ste Croix M, Priestly M, Lloyd R, Oliver J. ACL injury risk in elite female youth soccer: Changes in neuromuscular control of the knee following soccer-specific fatigue. *Scand J Med Sci Sports* 2014; 25(5): 531-538. doi: [10.1111/sms.12355](https://doi.org/10.1111/sms.12355)
44. Andersson H, Randers M, Krstrup P, Mohr M. Elite female soccer players perform more high-intensity running when playing in international games compared with domestic league game. *J Strength Cond Res.* 2010; 24(4): 912-919. doi: [10.1519/JSC.0b013e3181d09f21](https://doi.org/10.1519/JSC.0b013e3181d09f21).
45. Sporis G, Jukic I, Ostojic S, Milanovic D. Fitness profiling in soccer: Physical and physiologic characteristics of elite players. National Strength and Conditioning Association. *J Strength Cond Res.* 2009; 23(7): 1947-1953. doi: [10.1519/JSC.0b013e3181b3e141](https://doi.org/10.1519/JSC.0b013e3181b3e141)
46. Mohr M, Krstrup P. Yo-Yo intermittent recovery test performances within an entire football league during a full season. *J Sports Sci.* 2014; 32(4): 315-327. doi: [10.1080/02640414.2013.824598](https://doi.org/10.1080/02640414.2013.824598)
47. O'Connor DT. Motion analysis of elite touch players. In: *Fourth World Congress of Science and Football*. London, England: E&FN Spon: 2002.
48. Reilly T, Keane S. Estimation of physiological strain on Gaelic football players during match-play. In: *Science and Football IV*. London, England: E&FN Spon: 2002.
49. Antonacci L, Mortimer L, Rodrigues V, Coelho DB, Soares DD, Silami-Garcia E. Competition, estimated and test maximum heart rate. *J Sports Med Phys Fitness.* 2007; 47(4): 418-421.
50. Mortimer L, Condessa L, Rodrigues V, Coelho DB, Soares DD, Silami-Garcia E. Comparison between the effort intensity of young soccer players in the first and second halves of the soccer Game. *Portuguese Journal of Sport Sciences.* 2006; 6: 154-159.
51. Vencurik T, Nykodym J, Struhar I. Heart rate response to game load of U19 female basketball players. *International Network of Sport and Health Science.* 2015; 10(Proc1): 410-417. doi:[10.14198/jhse.2015.10.Proc1.33](https://doi.org/10.14198/jhse.2015.10.Proc1.33)
52. Orendurff MS, Walker JD, Jovanovic M, Tulchin KL, Levy M, Hoffmann DK. Intensity and duration of intermittent exercise and recovery during a soccer match. *J Strength Cond Res.* 2010; 24(10): 2683-2692. doi:[10.1519/JSC.0b013e3181bac463](https://doi.org/10.1519/JSC.0b013e3181bac463)
53. Small K, McNaughton L, Lovell R. The effects of multidirectional soccer-specific fatigue on markers of hamstring injury risk. *J Sci Med Sport.* 2010; 13(1): 120-125. doi: [10.1016/j.jsams.2008.08.005](https://doi.org/10.1016/j.jsams.2008.08.005)