

This is a peer-reviewed, final published version of the following document and is licensed under Creative Commons: Attribution 4.0 license:

Jarman, Rob ORCID logoORCID: <https://orcid.org/0000-0002-3560-1266>, Mattioni, Claudia, Russell, Karen, Chambers, Frank M ORCID logoORCID: <https://orcid.org/0000-0002-0998-2093>, Bartlett, Debbie, Martin, M. Angela, Villani, Fiorella, Webb, Julia C ORCID logoORCID: <https://orcid.org/0000-0002-1652-965X> and Cherubini, Marcello (2019) DNA analysis of *Castanea sativa* (sweet chestnut) in Britain and Ireland: Elucidating European origins and genepool diversity. PLoS ONE, 14 (9). pp. 1-26. doi:10.1371/journal.pone.0222936

Official URL: <http://dx.doi.org/10.1371/journal.pone.0222936>
DOI: <http://dx.doi.org/10.1371/journal.pone.0222936>
EPrint URI: <https://eprints.glos.ac.uk/id/eprint/7265>

Disclaimer

The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.

RESEARCH ARTICLE

DNA analysis of *Castanea sativa* (sweet chestnut) in Britain and Ireland: Elucidating European origins and genepool diversity

Rob Jarman^{1☯*}, Claudia Mattioni^{2☯}, Karen Russell^{3☯}, Frank M. Chambers^{1☯}, Debbie Bartlett^{4☯}, M. Angela Martin^{5☯}, Marcello Cherubini^{2☯}, Fiorella Villani^{2‡}, Julia Webb^{1‡}

1 Centre for Environmental Change and Quaternary Research, School of Natural & Social Sciences, University of Gloucestershire, Cheltenham, United Kingdom, **2** Istituto di Ricerca sugli Ecosistemi Terrestri (IRET), Consiglio Nazionale delle Ricerche, Porano, Italy, **3** K Russell Consulting Ltd, Leighton Bromswold, Huntingdon, United Kingdom, **4** Faculty of Engineering & Science, University of Greenwich, Chatham Maritime, United Kingdom, **5** Department of Genetics, University of Cordoba, Cordoba, Spain

☯ These authors contributed equally to this work.

‡ These authors also contributed equally to this work.

* rjarman1@glos.ac.uk, robinajarman@gmail.com



OPEN ACCESS

Citation: Jarman R, Mattioni C, Russell K, Chambers FM, Bartlett D, Martin MA, et al. (2019) DNA analysis of *Castanea sativa* (sweet chestnut) in Britain and Ireland: Elucidating European origins and genepool diversity. PLoS ONE 14(9): e0222936. <https://doi.org/10.1371/journal.pone.0222936>

Editor: Filippas A. Aravanopoulos, Aristotle University of Thessaloniki, GREECE

Received: March 20, 2019

Accepted: September 10, 2019

Published: September 25, 2019

Copyright: © 2019 Jarman et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: RJ received funding for this study from the University of Greenwich (UK). The Future Trees Trust (FTT) funded Karen Russell 'KR' (K Russell Consulting Ltd) to provide samples from the FTT collection for DNA analysis. DNA analysis was undertaken by Dr Claudia Mattioni 'CM' (IRET), whose laboratory costs for this project were

Abstract

Castanea sativa is classified as non-indigenous in Britain and Ireland. It was long held that it was first introduced into Britain by the Romans, until a recent study found no corroborative evidence of its growing here before c. AD 650. This paper presents new data on the genetic diversity of *C. sativa* in Britain and Ireland and potential ancestral sources in continental Europe. Microsatellite markers and analytical methods tested in previous European studies were used to genotype over 600 *C. sativa* trees and coppice stools, sampled from ancient semi-natural woodlands, secondary woodlands and historic cultural sites across Britain and Ireland. A single overall genepool with a diverse admixture of genotypes was found, containing two sub groups differentiating Wales from Ireland, with discrete geographical and typological clusters. *C. sativa* genotypes in Britain and Ireland were found to relate predominantly to some sites in Portugal, Spain, France, Italy and Romania, but not to Greece, Turkey or eastern parts of Europe. *C. sativa* has come to Britain and Ireland from these western European areas, which had acted as refugia in the Last Glacial Maximum; we compare its introduction with the colonization/translocation of oak, ash, beech and hazel into Britain and Ireland. Clones of *C. sativa* were identified in Britain, defining for the first time the antiquity of some ancient trees and coppice stools, evincing both natural regeneration and anthropogenic propagation over many centuries and informing the chronology of the species' arrival in Britain. This new evidence on the origins and antiquity of British and Irish *C. sativa* trees enhances their conservation and economic significance, important in the context of increasing threats from environmental change, pests and pathogens.

contributed to by FTT; RJ and KR prepared a preliminary report to FTT. FTT also supported Rob Jarman 'RJ' (University of Gloucestershire) by contributing to the laboratory costs of DNA analysis of the Historic trees samples, undertaken by Dr Claudia Mattioni (IRET). The specific roles of these authors are articulated in the 'author contributions' section. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript

Competing interests: Karen Russell Consulting Ltd provided samples for DNA analysis. There are no patents, products in development or marketed products to declare. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Introduction

Castanea sativa (sweet chestnut) is presently classified as non-indigenous in Britain and Ireland, and until a recent re-evaluation [1] it was thought introduced to Britain during the Roman period (AD 43–410), and to Ireland sometime thereafter: a test of the 'Roman introduction to Britain' thesis has not found any corroborative evidence [1]. The role of natural and anthropogenic vectors for the expansion of various tree and shrub species from refugia since the Last Glacial Maximum (LGM) in Europe is increasingly being reviewed, including *C. sativa* [2], *Quercus robur* and *Q. petraea* [3] and *Corylus avellana* [4]. DNA analysis provides a powerful tool, additional to archaeological and palaeoenvironmental analyses, to track the movement of trees across the post-LGM landscape [5, 6].

Castanea sativa in continental Europe

Palaeoenvironmental evidence indicated that *C. sativa* survived in refugia during the LGM in Iberia, S France, Italy, the Balkans, Greece, Turkey and the Caucasus [2, 7]. Genetic studies of 'wild' *C. sativa* populations in these areas [8, 9, 10, 11] have described these same refugia groups as three distinct genepools: 'eastern' (east Turkey to the Caucasus), 'central' (west Turkey, Greece, Bulgaria) and 'western' (Italy, S Switzerland, S France, Spain, Portugal). Recent studies [12, 13] have corroborated the indigenous status of *C. sativa* populations in parts of continental Europe and their derivation from these refugia; and described the redistribution of genotypes via natural colonization and anthropogenic translocation. *C. sativa* was used for wood and fruit production from before the Classical Greek and Roman periods, when it was translocated across southern Europe; and was then more intensively cultivated by monastic, royal and noble estates, by farmers and by foresters across Europe [14, 15, 16, 17, 18].

Castanea sativa in Britain

C. sativa has been classified in accounts of the British flora as an archaeophyte of Roman introduction [19, 20, 21, 22]: 'an honorary native' [23] is how it is typically described in some ancient semi-natural woodland contexts. *C. sativa* trees are presently recorded growing across most of Britain (Fig 1) [24], but *C. sativa* woodland is only locally abundant where slightly acidic (pH 4.5–5.5) soils over moist but well-drained substrates predominate [25].

There are approximately 28,200 ha of *C. sativa*-dominated woodland in Britain [26], located predominantly in the east, south and west regions of England, occurring as coppice and high forest in ancient semi-natural woodlands and secondary woodlands. *C. sativa* can presently set viable seed and self-regenerate as far north as Sutor, Cromarty, Scotland (57° 40'N, 4° 01'W) (Diane Gilbert 2018 pers. com.).

In many historic landscapes in Britain, sweet chestnut trees and pollards are amongst the oldest and largest of any species, situated in wood pastures, medieval deer parks, historic designed parks and gardens, and on ancient boundaries within farmed landscapes [23]. In Britain there are over 1,400 individual ancient *C. sativa* trees of >6 metres girth (measured at 1.3–1.5 m above the root collar) [27].

Despite its oft-asserted status as a Roman archaeophyte, only nine acceptable records of *C. sativa* remnants have been found in Britain up to AD 650 (including only one find of nut fragments): none could be verified as grown in Britain [1]. The early history of *C. sativa* growing in Britain is unknown, despite three centuries of speculation [23, 28]. The earliest written record of *C. sativa* growing in Britain presently dates from AD 1113 [29]: this and two other written records from the 12th and 13th centuries AD indicate specific *C. sativa* trees and woods that were established in at least three places in southern England and SE Wales sometime



Fig 1. Distribution of *Castanea sativa* in Britain and Ireland.

<https://doi.org/10.1371/journal.pone.0222936.g001>

before the 12th century AD [1]. The oldest extant British *C. sativa* trees that have been dendrochronologically cross-dated presently date from c. AD 1640 [30].

***Castanea sativa* in Ireland**

C. sativa is conventionally presumed to have been introduced to Ireland sometime in the medieval period [31, 32]. The largest extant *C. sativa* tree in Ireland is the 10.78 m girth 'Wesley tree' (Rossanna, Co. Wicklow); thirty-eight ancient *C. sativa* trees are recorded for Eire [33]. In Northern Ireland, there are twenty-two ancient *C. sativa* trees >6 m girth [27]. In Eire, Coillte (the State-sponsored forestry company) manages c. 100 ha of *C. sativa* woodland (c. 95% of the *C. sativa* woodland in Eire), mostly in the southern and south-eastern counties (Ted Horgan 2017 pers. com.)—Fig 1.

Genetic analysis of *Castanea sativa* in Britain and Ireland

Despite many studies of *C. sativa* across Europe [12, 13], British and Irish populations have largely been ignored. Two previous studies assessed a few samples from *C. sativa* trees in England: six coppice groups in southern England [34] were used in the European CASCADE project to establish reference SSR markers for *C. sativa* [35, 36, 37] and subsequently [38]; and some preliminary data from the present research were included in an Europe-wide landscape genetics study of *C. sativa* [11].

Aims of the paper

This paper aims to describe the genetic diversity of *C. sativa* trees in Britain and Ireland, to discover any geographical and typological associations and, by comparison with *C. sativa* genotypes from sites in continental Europe, indicate potential ancestral sources. Genetic information can locate close kinships and clonal groups of plants; and help to determine the antiquity and management history of individual trees and coppice stools. The combined information on genetic diversity, relatedness and antiquity can help elucidate whence *C. sativa* arrived in the British Isles, and possibly when.

Materials and methods

Castanea sativa trees were sampled from sites across Britain and Ireland, which were selected to represent a broad range of ecological, cultural and geographical contexts. Plant material for DNA analysis was collected from ancient trees and coppice stools in ancient semi-natural woodlands and historic cultural sites in England and Wales, hereinafter referred to as the 'Historic trees' sample set; and from trees and coppice of mixed antiquities selected from sites included in the Future Trees Trust (FTT) Sweet Chestnut Improvement Programme [39], which covers England, Wales, Scotland and Ireland, hereinafter referred to as the 'FTT trees' sample set.

Sampling strategy and site selection

For the 'Historic trees' samples, fresh leaf samples were collected from *Castanea sativa* plants in ancient semi-natural woodlands and historic non-wooded landscapes ('plant' here means 'standard' single-trunked trees, multi-stemmed trees including pollards and multi-stemmed 'coppiced' trees growing from 'stools') [40]. These 'historic' sites were identified from a range of sources, including archives and published contemporary accounts; the Ancient Tree Inventory [27]; maps of ancient semi-natural woodlands; designated historic monuments, parks and gardens; Forestry Commission and National Trust records; and peer review. A preliminary survey in 2013 tested the methods, followed by extensive sampling across England and Wales during 2014–2018. Wherever possible, several trees were sampled within each site to represent its characteristics: within an ancient woodland coppice, several discrete stools were sampled from different parts of the wood, and some stools were multi-sampled (to test for clonality); within an historic parkland, several ancient trees were sampled from specific features such as an avenue or grove, and some individual trees were multi-sampled (to test for grafting or bundle planting). For some historic parkland/garden sites, where only a single ancient *C. sativa* tree existed, samples were taken from several parts of the tree to test its genetic integrity. Several historic trees and coppice stools were re-sampled in consecutive years to test whether the analysis of leaves from the same stem but in different annual growth seasons produced consistently replicable results. Several trees were re-sampled by collecting dormant buds (during winter) instead of leaves, to check their replicability.

For the 'FTT trees', fresh leaf samples were collected from the FTT clonal archive of plants, which had been established from outstanding individual timber trees and coppice stools located in long-established woodland (not recent plantations) [39].

DNA extraction and Microsatellite analysis

For direct comparison and integration of British and Irish samples with samples from an earlier Europe-wide study [11], the analytical procedures and materials used were identical with that study. Eight polymorphic microsatellite markers developed and used for *Castanea sativa*

[10, 34, 35] (CsCAT-1, -2, -3, -6, -14, -16; and EMCs-25, -38) were used to genotype the British and Irish samples: six of these markers had been used by the European study [11] (CsCAT-1, -3, -6, -16; and EMCs-25, -38). Multiplex-PCRs were performed exactly as described in that study: the samples were run on an ABI Prism 3130 Avant DNA sequencer and the resulting raw data were analyzed using GeneMapper software (Life Technologies). The alleles were determined by automated binning and checked by visual inspection.

Data analysis

The microsatellite data were used to estimate genetic diversity, to determine whether the British and Irish samples constituted a single genepool or several genepools: the British and Irish samples were then integrated with the European dataset [11] and analysed to investigate the potential continental European origin(s) of the British and Irish germplasm. The first step was to identify discrete genotypes from all the samples, using GenAlEx 6.51b2 Multilocus Matching Programme, so that the diversity analyses could be run without any matching samples. Groups of matching samples were assessed separately for clonal characteristics.

Genetic diversity and population structure

Null allele frequencies were estimated for each locus and population using FreeNA [41]. Allelic richness (A_r) and private alleles (P_{Ar}) were calculated by the statistical method of rarefaction to avoid bias due to different sampling size, implemented by the software HP-Rare 1.1 [42]. Genetic diversity parameters were measured using the software GenAlEx 6.51b2 [43]: observed (N_a) and effective (N_e) number of alleles, observed (H_o) and expected (H_e) heterozygosity [44], unbiased estimate of mean expected heterozygosity (U_{He}) and the diversity index of Shannon (I) were computed; and the inbreeding coefficient (F_{IS}) was derived using GENEPOP 4.2 software Option 5 [45]. The values of the population differentiation coefficient (F_{ST}) were assessed using FreeNA [41] to detect genotypic differences, employing ENA correction for null allele bias and F_{ST} overestimation. A threshold for determining relevance of F_{ST} estimates was derived from upper and lower (10% and 90%) percentiles of the dataset of pairwise values, to indicate 'high' and 'low' differentiation relative to the actual range of genetic difference in the populations observed. AMOVA was performed using GenAlEx 6.51b2.

The genetic distance (GD) among individuals and the Nei genetic distance among groups of individuals were calculated using GenAlEx 6.51b2; Principal Coordinates Analyses (PCoA) were run to test various parameters that might define postulated populations. Unweighted Neighbour Joining Cluster analysis (UPGMA) used DARwin ver.6 [46] to construct a dendrogram of hierarchical relationships between samples.

Pairwise relatedness (r) was estimated using Queller and Goodnight [47] and Lynch and Ritland [48] analyses run in GenAlEx v.6.51b2, to detect kinships within and between populations. The Lynch and Ritland method (LRM) states that an r value close to the maximum calculable 0.5 indicates an identical twin, 0.25 indicates a full sibling relationship (parent–offspring, or siblings sharing the same parents), and 0.125 indicates a half sibling (one shared parent). In the Queller and Goodnight method (QGM), an r value close to the maximum 1, 0.5 and 0.25 correspond to identical twin, full sibling/parent–offspring, and half sibling, respectively.

The samples were analysed using GenAlEx 6.51b2 Multilocus Matching programme to identify samples with matching multilocus genotypes for codominant data, at two hierarchical stages of division: samples matching exactly at all loci ('clonal'), and matching at all loci except one (a 'near match'). The analysis was run for the 8–SSR dataset and then separately for the 6–SSR dataset. Unexpected results (based on the tree/stool descriptions) were checked by visual assessment of the GeneMapper automated binning output, followed by repeat analyses

of leaf material from that sample. Samples recorded as 'near matches' were tested for the degree of allelic difference; and checked on site for evidence of direct biological linkage between potentially identical clonal plants (in biological terms, between a 'genet' and a putative 'ramet'). Where only a single allele difference at a single locus was recorded (such as '232' instead of '230'), a somatic mutation from the parent plant (genet) was accepted and the sample pair was considered 'clonal'. All 'near matches' greater than a single allele difference ('234' instead of '230') were rejected as 'non-clonal'. These results were compared with the r pairwise relatedness results (LRM and QGM) to check the compatibility of these methods for determining very close or clonal relationships.

A Bayesian analysis (STRUCTURE v.2.3.4 [49]) was run with the option of including prior information on the spatial location of populations, using the admixture model, with the number of tested clusters (K) from 1 to the purported number of provenances plus 2. Six independent runs were performed for each K value, with a burn-in period of 10^3 steps followed by 10^5 MCMC replicates. The analysis assigns individuals into each of the K clusters based on the membership coefficient (Q -value). To identify the number of clusters that best explained the data, the rate of change on $L(K)$ (termed ΔK) between successive K values was calculated [50] using STRUCTURE HARVESTER software [51]. The six runs for each simulation were averaged using CLUMPP and displayed using DISTRUCT, provided by CLUMPAK [52]. To delineate genetic repartition of sweet chestnut populations, a phylogenetic dendrogram UPGMA [53] was constructed, using Nei genetic distance corrected for F_{ST} using the software POP-TREE2 [54] and drawn with FigTree v1.4.3 [55].

Site characterisation for British and Irish samples

Five parameters were selected to provide geographical, ecological and cultural frameworks within which to analyse the British and Irish genotypes for any spatial or typological pattern of distribution: Ordnance Survey 100km square Map Grid; Administrative Counties and Regions; Seed Zones [56]; Cultural landscapes [57]; and Site Types (see below). Each parameter dataset was analysed using GenAlEx 6.51b2 to generate a 'genetic distance by population' table, PCoA chart and UPGMA dendrogram. GENEPOP 4.2 produced F_{IS} values of the inbreeding coefficient for each site, and F_{ST} estimates of pairwise differentiation between sites.

Results

In total, 753 samples were collected from 259 sites across Britain and Ireland (S1 File; S1 Table); in addition, 22 samples were collected from the Tortworth Chestnut tree (South Gloucestershire, England) for clonal analysis and antiquity tests. Five types of site were identified in the survey: 'Type A' (high forest standard trees < c. 200 years old); 'Type B' (coppice stools < c. 200 years old); 'Type C' (ancient coppice stools in ancient semi-natural woodland > c. 200 years old); 'Type D' (ancient trees in woods and historic landscapes > c. 200 years old); and 'Type E' (historic garden trees, typically singular and of great antiquity). These Site Types are fully described in S3 File (para 5).

Genetic diversity of British and Irish *Castanea sativa* trees

The eight microsatellite loci comprised 91 alleles: CsCAT-2, -3, -6 and EMCs38 had the highest number of alleles (A and Ae) and values of genetic diversity (H_e); EMCs25 showed the lowest number of alleles (5), low values of H_o and high values of F_{IS} . FreeNA check for null alleles showed EMCs25 consistently produced null alleles.

Of the 753 samples, 611 were identified as discrete genotypes (GenAlEx 6.51b2 Multilocus Matching Programme) and used for the diversity analyses, for Britain and Ireland as a single

Table 1. Genetic diversity measures for *Castanea sativa* in Britain and Ireland.

Population	N	A	Ae	I	Ho	He	UHe	Ar	PAr	F _{IS} P<0.001	F _{ST} without ENA	F _{ST} with ENA
England	516	14.5	4.957	1.792	0.713	0.74	0.741	10.07	0.97	0.041	0.0081	0.0083
Ireland	48	10	4.431	1.669	0.661	0.7	0.708	9.97	0.89	0.062	0.006	0.0057
Wales	47	9.25	4.102	1.604	0.657	0.706	0.713	9.25	0.56	0.074	0.0141	0.0138
Britain & Ireland	611	14.75	4.915	1.69	0.704	0.737	0.738	14.75	—	0.05		

N = Number of samples, A = Mean Number of different alleles, Ae = Mean Number of effective alleles, I = Shannon's Information Index, Ho = Observed heterozygosity, He = expected heterozygosity, UHe = Unbiased expected heterozygosity, Ar = allelic richness (average over loci), PAr = private allelic richness (average over loci), F_{IS} = inbreeding coefficient, F_{ST} = genetic differentiation, ENA = FreeNA estimate of null allele frequency per locus.

<https://doi.org/10.1371/journal.pone.0222936.t001>

site group and for three separate site groups 'England', 'Ireland' and 'Wales' (Table 1); 88 groups of matching samples were identified for separate clonal analyses.

Comparable values of genetic diversity parameters were observed for England, Ireland and Wales; slightly higher values of allelic richness (Ar) and private allelic richness (PAr) were observed for England samples. F_{IS} was higher for Wales (0.074) than for England or Ireland, as also was F_{ST} (Wales 0.014), indicating a relatively higher degree of inbreeding and isolation for Wales sites. AMOVA analysis of the England, Ireland and Wales groups of samples (S2 Table) found the majority of variability (95%) within individuals, with 4% amongst individuals and 1% amongst groups.

Genetic structure

The genetic structure of England, Ireland and Wales samples was revealed by three statistical approaches:

1) The Bayesian analysis implemented by STRUCTURE indicated the presence of two gene-pools, 'Genepool A' and 'Genepool B' (K = 2 Evanno, $\Delta K = 700$)—Fig 2.

The England, Ireland and Wales samples were mapped according to their 'Genepool A' or 'Genepool B' membership—Fig 3 and S2 File (for viewing in GIS mapping). Local and regional spatial clusters of 'Genepool A' and 'Genepool B' sites were evident, indicating local area fidelity, but overall there was no evidence of broad geographical separation between the two gene-pools.




Fig 2. STRUCTURE analysis of England, Ireland and Wales (611 samples, 8 SSRs): (2a) 'Genepool A' (red) and 'Genepool B' (blue); (2b) K = 2 (Evanno); (2c) differentiation between Ireland and Wales in 'Genepool A' and 'Genepool B'.

<https://doi.org/10.1371/journal.pone.0222936.g002>




Fig 3. England, Ireland and Wales samples in ‘Genepool A’ (red) and ‘Genepool B’ (blue). Map base open-sourced from ‘EuroGeographics and UN-FAO @EuroGeographics’.

<https://doi.org/10.1371/journal.pone.0222936.g003>

2) The dendrogram of Nei genetic distance, using the Unweighted Neighbour Joining Method (DARwin ver. 6), indicated two main clades and two minor clades: genetic distance among individuals was low—Fig 4 (see S1 Fig for expandable image).

Some samples from the same site clustered together, whilst other samples from the same site dispersed through the dendrogram: ‘top down’ linkages within and between some sites were identifiable. Comparison of the dendrogram with the STRUCTURE output allowed




Fig 4. Dendrogram for England, Ireland and Wales samples (Nei genetic distance) indicating two major and two minor clades (‘Genepool A’—red; ‘Genepool B’—blue).

<https://doi.org/10.1371/journal.pone.0222936.g004>




Fig 5. PCoA plot of samples from England, Ireland and Wales, showing differentiation between Wales and Ireland.

<https://doi.org/10.1371/journal.pone.0222936.g005>

matching of ‘Genepool A’ and ‘Genepool B’ members to the two main clades and to the two minor clades of the dendrogram, shown ringed in red (Genepool A) and blue (Genepool B) on Fig 4, such that the dendrogram depicts three main clades, with ‘Genepool A’ to the left, ‘Genepool B’ centrally in 2 clades, and ‘Genepool A’ minor clade to the right.

3) PCoA evaluation of the England, Ireland and Wales groups of samples corroborated the differentiation between Wales and Ireland, with a high degree of separation on the x -axis (74.7%)—Fig 5.

Relatedness of the England, Ireland and Wales samples and sites

Three measures were used to test for relatedness between samples and between sites: F statistics, relatedness estimation (r) and clonal analysis. Pairwise F_{ST} estimates, using a threshold value of $F_{ST} = 0.05$, found no great differentiation (all scores were <0.05) between the three country groups, with the highest differentiation between Wales and Ireland ($F_{ST} = 0.016$). Pairwise F_{ST} estimates using the five parameter groups (S3 File; S3 Table) found no differentiation between any of the samples at $F_{ST} > 0.05$ for any of the five parameters, except in the ‘Counties’ dataset, where 22 out of 300 (7.3%) sample pairs measured >0.05 , to a maximum 0.093, again differentiating between Wales and Ireland sites. Some sites (including iconic historic trees at Bushy House, Seven Sisters Penshurst and the Tortworth Chestnut) had values of $F_{ST} > 0.15$ across all five parameters, and were highly differentiated from all other British and Irish sites. F_{IS} values were estimated for each site with >2 samples (S3 Table): they ranged from -0.857 (Warren Plantation A, Essex) to 0.467 (America Wood, Isle of Wight).

Kinship r estimation between individual samples, using pre-selected values of $r > 0.4$ (LRM) and >0.75 (QGM) to define a threshold above which samples were deemed to have very close kinship, identified the ‘highest related’ pairs of samples (S4 Table); these were corroborated by the pairwise F_{ST} and clonal ‘near matches’ analyses. For example, 73 samples from coppice stools and ancient trees in ‘Castiard’ (Forest of Dean, Gloucestershire) that had been analysed as clonal groups produced pairwise r scores of 0.500 for LRM and 1.000 for QGM; and very close kinship scores were obtained for ‘near match’ samples (up to 0.383 in LRM and 0.924 in QGM)—S5 Table. Some of the very close kinship samples from across Britain grew on adjacent trees/stools, but some were from widely disjunct trees (such as between Croft

Castle in Herefordshire and Seven Sisters Penshurst in Kent; and between Godinton in Kent and Hagley Hall in Worcestershire), perhaps evincing historical vegetative and/or seed translocation at an inter-regional scale.

Clonal analysis determined the relatedness of samples between and within sites: from across England and Wales, 185 samples formed 88 clonal groups. Thirty samples formed 15 pairs of 'near matches', four of which differed by a single allele at a single locus, so could be presumed somatic mutations and re-classified as clones. This clonal analysis of tree and stool components enabled the definition of the 'genotypic' size of large-girthed coppice stools and multiple-stemmed/collapsed trees, as opposed to their 'visual' size, thereby indicating their true antiquity. For example, a 16 m girth ring-form stool in Welshbury Wood was found genetically identical in all its parts, so is formed of a single-genotype original plant that has grown over centuries of repeated cutting into a very large stool; whereas a 15 m girth ring-stool growing nearby, which visually/structurally appeared like a single entity, was found to consist of three genetically different plants, indicating that it was probably much younger than the 'genetically entire' stool. A massive ancient tree at Baldwyns (Kent) appeared visually to be a single tree base with four main trunks with a basal girth of 12 m, but clonal analysis showed it was at least four genetically separate plants (one of which was a 'near match' with a large coppice stool in ancient woodland 10 km distant). In another case, two trees growing hundreds of metres apart (at Abenhall, Gloucestershire) were found to be clonal and so were presumed translocations of vegetative material. Two examples of grafted trees were revealed on two sites in England. The clonal information was used to confirm the antiquity of ancient trees and stools in the Site Type classification, below.

Site-based relationships within the British and Irish genepool

The England, Ireland and Wales samples were sorted into five parameter-defined groups. In the four geographically defined groups, the PCoA and UPGMA analyses indicated genetic differentiation between samples in Ireland and Wales. In the fifth group, the Site Type evidence of tree/stool size and site history indicated that older (>200 years) trees and stools (Types C and D) were genetically differentiated from younger (<200 years) trees and stools (Types A and B). Type B (younger coppice) was the most differentiated of the five Site Types: STRUCTURE analysis placed it predominantly (64%) within 'Genepool B'. Ancient historic garden trees (Type E) were differentiated by the PCoA from other ancient trees (Type D) as well as from the younger sites (Types A and B): STRUCTURE analysis placed Type E trees predominantly (60%) within 'Genepool A'. The detailed results from the five parameter analyses are presented in [S3 File](#).

The 'Historical trees' group was compared with the 'FTT trees' group: no genetic differentiation was found.

European dimension for the British and Irish samples

The six SSR markers used for the previous European research [11] were applied to the British and Irish samples, identifying 608 discrete genotypes (560 from Britain, 48 from Ireland) for integration and comparative analysis with the continental European samples. [S6 Table](#) contains the full dataset (1332 samples, 41 site groups) of continental European, British and Irish samples.

Genetic diversity analysis of the European samples

The continental European, British (subdivided into England and Wales) and Irish dataset, comprising 1332 samples in 41 site groups, was assessed for genetic diversity ([S7 Table](#)). The

number of alleles (N_a) and number of effective alleles (N_e) were greater for the England, Ireland and Wales groups than for any of the continental European groups; as were the values of I and H_e and allelic richness (A_r). Private allelic richness (PAr) was higher in Wales than in 61% of the continental European groups and in Ireland higher than in 44% of those groups. Null allele frequency estimation showed EMCs25 produced null alleles. Molecular variance was 11% among populations, 6% within groups and 83% within populations ($P < 0.001$). The combined continental European, British and Irish dataset produced a F_{IS} value of 0.067 and F_{ST} value of 0.114 (F_{ST} with ENA was 0.108).

STRUCTURE analysis of the continental European, England, Ireland and Wales samples

For an initial STRUCTURE analysis, the England, Ireland and Wales samples were incorporated into the original full dataset of continental European populations (eastern and western) [11], comprising 2199 samples from 319 site groups. The results previously published [11] were confirmed, indicating a clear separation of 'eastern' from 'western' European site groups, with no connection between British and Irish sites and the 'eastern' European sites (S4 File).

A 'western' European sites subset was defined, containing the British and Irish samples with Portugal (PT), Spain (SP), France (FR), Italy (IT) and some sites from Slovakia (SK), Romania (RO) and Hungary (HU): it comprised 1332 samples in 41 groups (38 continental European groups plus 'England', 'Ireland' and 'Wales'). STRUCTURE analysis identified two main groups ($K = 2$ Evanno), termed 'Genepool C' and 'Genepool D': each site was allocated to 'Genepool C' or 'Genepool D' by the 'K2-pop' CLUMPP output, as shown in Fig 6, using $Q = 0.5$ as a threshold; and mapped (Fig 7; S5 File for GIS data in .kml format).

The England, Ireland and Wales samples fitted predominantly (78%) within 'Genepool C', whereas in the Britain and Ireland analysis (using 8 SSRs) they had divided almost equally into two groups ('Genepool A' and 'Genepool B')—Fig 8.

STRUCTURE analysis of the 'western' European groups indicated that 'Genepool C' sites in England, Ireland and Wales derived predominantly from northern Iberia, southern France, central/southern Italy and northern Romania; whereas 'Genepool D' sites in England, Ireland and Wales derived predominantly from western and southern Iberia, northern Italy and Hungary, Romania and Slovakia. 'Genepool D' affiliations were stronger for Wales (notably with PT03, SP07, SP14) than for Ireland, so genetic differentiation between Wales and Ireland was again evinced. Typologically, Site Type E (ancient historic garden trees) was predominantly affiliated with 'Genepool D' (PT01, PT03, SP07), whereas the other four Site Types (A–D) were more affiliated with 'Genepool C'. (Table 2).




Fig 6. Continental Europe, England, Ireland and Wales sites, STRUCTURE output defining 'Genepool C' (green) and 'Genepool D' (purple).

<https://doi.org/10.1371/journal.pone.0222936.g006>

The UPGMA analysis

A dendrogram of the continental European, England, Ireland and Wales sites was produced using UPGMA analysis (F_{ST} Corrected option) (Fig 9). The England ('ENG'), Ireland ('IRE') and Wales ('WAL') groups were most closely associated with FR01, FR02, FR03, RO01, IT05, SP17, SP07 and SP12.

For comparison, the England, Ireland and Wales samples were re-grouped according to their Site Types A–E (above; S3 File) to test whether older generation trees might derive from different parts of Europe cf. younger generation trees. The resulting dendrogram (Fig 10) indicates broadly the same clustering as Fig 9, but with PT01 and PT03 aligned closer to England, Ireland and Wales; and SP03, SP08, SP13 and SP04 separated from them. The separation of Site Types A and B (younger trees) from Site Types C, D and E (older trees) corroborated the British and Irish data results, above, as summarised in Table 2.

The PCoA assessment

PCoA of the continental European, England, Ireland and Wales samples produced Fig 11: England and Wales straddled the centre of the plot (separated from Ireland), most closely associated with FR01, FR03, SP07, SP17, SK03, RO01, IT03, IT05 (similar to the STRUCTURE and UPGMA results). The percentage of variation explained by either axis was relatively low but the outlying positions of RO02, RO03 and SK02 were noteworthy, as these sites appeared similarly anomalous in the F_{ST} and UPGMA analyses.

Relationships within and between European sites

F statistics (F_{IS} and F_{ST}) were used to test for relatedness. For the F_{IS} estimation, 38 continental European sites were compared with 143 England, Ireland and Wales sites (excluding single-




Fig 7. Map of European sites, STRUCTURE output 'Genepool C' (green) and 'Genepool D' (purple). Map base open-sourced from 'EuroGeographics and UN-FAO @EuroGeographics'.

<https://doi.org/10.1371/journal.pone.0222936.g007>




Fig 8. Dendrogram for England, Ireland and Wales samples (Genepool A–red; Genepool B–blue) overlain with their continental European classification into ‘Genepool C’ (green) and ‘Genepool D’ (purple).

<https://doi.org/10.1371/journal.pone.0222936.g008>

sample sites): values of F_{IS} ranged from -0.8333 to 0.4783 , indicating a wide range presumed caused by diverse numbers of samples per site. (S8 Table provides detailed site values). The pairwise F_{ST} estimation compared 38 continental European sites with the ‘England’, ‘Ireland’ and ‘Wales’ groups and with selected site groups. (S9 Table provides detailed site values). The 10percentile and 90percentile points calculated for each dataset (S10 Table) were found compliant with thresholds for $F_{ST} < 0.05$ (= ‘low’) and > 0.15 (= ‘high’) determined in previous studies [8, 10, 58, 59]. Considering samples with pairwise $F_{ST} < 0.05$, a consistent pattern of low differentiation was identified between specific continental European and England, Ireland and Wales sites (S9 Table)—summarised in Table 2. Continental European sites having the closest relatedness with England, Ireland and Wales sites were FR02 and FR03, followed by FR01, RO01, SP12 and PT03; those with consistently high differentiation from England, Ireland and Wales sites were RO03, SK02, IT07, SP01, SP09 and PT02. The PT02 (Portugal) site provides a useful illustration: it was highly differentiated (> 0.15) from all British and Irish sites, apart from Silwood (0.035) and Nashdom (0.045), which linked < 0.05 with only three other European sites, so their specific linkage with PT02 appears meaningful. Some British and Irish sites displayed no evident connection with any sites: the Tortworth Chestnut linked closest with RO01 (0.1026) but with no other sites < 0.148 . Three Sisters Bachymbyd only linked

Table 2. Summary of dominant relationships between continental European and England, Ireland and Wales sites, as defined by Pairwise F_{ST} scores < 0.05 .

Continental European sites	Genepool	Dominant relationship -England/Ireland/Wales	Dominant relationship—site type
PT01	D	England	E—ancient historic garden trees
PT03	D	Wales/England	C, D, E—Ancient trees and stools including historic garden trees
SP06	C	Wales	No differentiation
SP07	D	Wales/England	D, E—ancient trees and historic garden trees
SP12	C	Ireland/England	C, D—ancient trees and coppice
SP14	D	Wales	No differentiation
FR01	C	Ireland/England	A, D—trees (ancient and modern), but not coppice
FR02	C	Ireland/England	A, B—modern trees and coppice
FR03	C	Wales/England	all types
IT05	C	England	B—modern coppice
RO01	C	Wales/England	all types

<https://doi.org/10.1371/journal.pone.0222936.t002>

with IT08 (0.0396); Bushy House had only one link <0.2 , with RO02 (for which r analysis yielded LRM 0.204 and QGM 0.720, indicating a weak full-sib/parent-offspring relationship).

The estimation of relatedness using LRM and QGM, using LRM >0.400 and QGM >0.750 to determine strong relationships [60] (S11 Table), showed high relatedness between continental European and England, Ireland and Wales sites in 60 sample pairs for LRM and in 77 pairs for QGM. Reducing the threshold of LRM to >0.25 (a conventional 'parent-offspring/full sib' class) yielded 6,841 paired matches within the continental European, British and Irish dataset, indicating a broad network of related samples but focussed on specific sites/regions. Interpreting relatedness from the LRM and QGM methods using microsatellite data requires caution [60], because genotypic similarity can vary according to allele frequencies of a population; however, LRM and QGM estimators are informative of relative kinship among individuals and have been endorsed as robust [61, 62, 63].

Clonal analysis determined clonal relations between and within sites and defined individual tree or stool 'genotypic size', supporting the Site Type analyses by determining tree or stool antiquity with an arbitrated girth measurement (below). Clonal matches identified in the England, Ireland and Wales 8-SSR dataset were regarded as superior to any matches identified in the 6-SSR dataset, which were rejected. No clonal matches were found with the continental European dataset, but 56 samples formed 28 pairs of 'near matches', of which three differed by a single allele at a single locus so could be presumed somatic mutations, defining three intra-site clonal pairs (within SP13, IT04 and SK03). Six of the 'near matches' were between continental European and England sites, but all were >1 allele different, so were not deemed clonal, although close relatedness was evident (S12 Table).




Fig 9. Continental European, England, Ireland and Wales dendrogram. The boxed section is explained in the text.

<https://doi.org/10.1371/journal.pone.0222936.g009>



Site Characterisation groupings of British and Irish sites compared with continental European sites

<https://doi.org/10.1371/journal.pone.0222936.g011>

The 'Site Types' grouping yielded PT03, FR03 and RO01 with the highest relatedness with all Site Types at the $F_{ST} < 0.05$ level. Site Type B (modern coppice) sites linked exclusively with IT05 and so appear to originate from continental European sites different from those of 'ancient coppice' (Type C) or 'ancient trees' (Type D). Another noteworthy finding was that only Type E (ancient garden trees) sites in Britain linked with PT01 (Table 2). For the 'Seed Zones' group, there were consistent links between all Seed Zones and FR03 and RO01; and slightly weaker links with PT03, SP12, FR01 and FR02. Seed Zones 304 and 305 (Wales) aligned closest with sites SP06, SP07, SP14 and IT03, with which Ireland did not align (reinforcing the differentiation between Wales and Ireland sites). The 'Counties' group displayed a similar pattern: FR03 was the predominant link for all Counties (except Cork and Tipperary, in Ireland), and FR02, RO01, SP07, SP12 and PT03 formed consistent links with most Counties. Sites RO03 and SK02 were consistently highly differentiated from almost all other sites, with F_{ST} frequently > 0.30 ; whereas FR03 and RO01 were consistently undifferentiated across all parameters with all sites, indicating widespread sharing of their genotypes across western Europe.

Summary of results indicating dominant relationships between British and Irish sites and continental European sites

In summary, the statistical analyses indicated that sites from England, Ireland and Wales linked predominantly (78%) with 'Genepool C' sites in continental Europe; but certain England and Wales sites strongly linked with 'Genepool D' (notably with PT03 and SP07) (Table 2). The STRUCTURE map depicts the continental European sites most strongly linked with England, Ireland and Wales—Fig 12: the dominant links are shown with a red star (see S5 File for GIS data).

Discussion

Discussion will focus on the relationships between *Castanea sativa* genotypes in Britain and Ireland and with those in continental Europe: possible causation for the most likely ancestral populations for British and Irish *C. sativa* lying in France (S and SW), Spain (NW and S), Portugal (N), central Italy and Romania will be sought.

Genotypes of *Castanea sativa* across England, Wales and Ireland

The bipolar genepool identified for Britain and Ireland (Fig 2 and Fig 3), possessing moderate genetic diversity, low levels of inbreeding ($F_{IS} = 0.04$) and low differentiation between sites ($F_{ST} = 0.01$), can be explained by analyses of samples grouped by cultural and environmental parameters. The 'Counties' analysis (respecting historic cultural divisions within Britain and Ireland) and the 'Fields of Britannia' [57] analysis (representing rural landscape regions created in Roman and early medieval periods) consistently differentiated genotypes from Wales, as also revealed by STRUCTURE analysis. The 'Seed Zones' [56] analysis (respecting phenotypic cues for tree adaptation) revealed a very weak influence on genotype distribution—as was found in a study of *Fraxinus excelsior* in Britain [64], which mapped haplotype distribution using the same Seed Zones. We suggest that *C. sativa* genotypes in Britain and Ireland may not have undergone sufficient reproductive cycles to respond to phenotypic cues, given that even the oldest trees and stools ($> c. 500$ years) were found to be single-generation plants: intra-site recruitment of multiple generations of trees from seed was very rarely observed, to be expected as coppice management typically used layering (vegetative reproduction) to establish new stools [65].




Fig 12. Dominant linkages between England, Ireland and Wales and continental European sites. Map base open-sourced from 'EuroGeographics and UN-FAO @EuroGeographics'.

<https://doi.org/10.1371/journal.pone.0222936.g012>

In that context, the genetic differentiation found between 'younger' and 'older' trees and stools in Britain and Ireland (viz. the difference between Site Types A, B and C, D) apparently indicates a preserved pattern of the origins and chronologies of *C. sativa* arrivals from continental Europe. The age differences between the trees and stools in these Site Types were confirmed by the clonal and relatedness analyses, whereby the true dimensions of a tree or stool could be defined genetically rather than visually, providing an accurate 'antiquity' measure. The genetic differentiation that was confirmed between 'modern' trees in woods and 'ancient' trees in parks and historic gardens could be linked with historical nut cultivation: ancient trees renowned for their eating nut qualities were recorded in the historic parks and gardens, indicating historical varietal selection for nuts. Systematic nut production was evinced from archive searches for several sites for the 12th–15th centuries AD [1]; and promotion of *C. sativa* for nut production in England is recorded from the 17th century AD [28].

Genotypic clusters were revealed in the genetic distance dendrogram (Fig 4), representing inter- and intra-site associations of closely related samples. For example, the main 'Group A' clade contained a discrete cluster of samples from the Forest of Dean (Gloucestershire) area known as 'Castiard' whence derived some of the earliest written records for *C. sativa* in Britain, c. AD 1140 [1]. Similarly, the Lydney Park (Gloucestershire) ancient trees formed another cluster, which also included a single tree growing in Ireland. Such clusters could indicate that a narrow range of genotypes had been propagated within a small area and become isolated; or that genotypes were exported from one site to another (as seed and/or vegetatively); or that disjunct sites share close kinship with a common (perhaps geographically remote) ancestor. These findings are similar to those reported for domestication of cultivars of ancient *C. sativa* trees in Italy and Iberia [66]; and comparable with a study of the genetic diversity and domestication of *Olea europaea* ssp. *europaea* in the Mediterranean basin [67].

Some single *C. sativa* trees were very highly differentiated from all others across Britain and Europe, such as the Tortworth, Seven Sisters Penshurst and Bushy House trees. These ancient 'single generation', 'single clone' trees are evidently genetically unchanged since their first establishment, so their present 'difference' represents the disjunct source(s) from which they originated (and which are excluded from the European sites accessed in this study), rather than evolution of *in situ* genetic isolation.

Clonal behaviour of *Castanea sativa* defining tree antiquity

A key outcome from this research has been the novel identification of clonal groups in ancient *C. sativa* trees and coppice stools: some stools that had been described as 'massive' and therefore 'ancient' [23] were found to comprise a single genotype, so confirming their great size and antiquity; others were found to be a composite of genotypes so not a massive, necessarily ancient, plant. The clonal analysis has evinced the natural collapse and regeneration of single trees and the historic management of trees and coppice using grafted boughs or layered stems; it also revealed the translocation of vegetative material to establish new plants. Some trees and stools that had appeared 'clonal' owing to their growth form were found to be non-clonal, demonstrating the unreliability of visual observation: a tree or stool cannot be considered a 'single plant structure' without genetic analysis. The clonal analyses underlined the importance of checking the PCR outputs of the automated binning process using knowledge of the sample's site context.

The clonal behaviour of *C. sativa* recorded in this study was compared with that of other tree species, to verify methods and conclusions. *Prunus avium* [68] and *Populus tremula* [69] can propagate vegetatively by root suckering, with a single clone sometimes occupying thousands of square metres; and *Tilia cordata* [70], *Corylus avellana* [4] and *Quercus pyrenaica* [71] can self-propagate from collapsed boughs and/or root budding. *C. sativa* evidently shares the natural habit of *Tilia cordata* of collapsing and regenerating from rooted layers, establishing large clonal patches in the process: clonal growth by bough layering and by stool expansion were observed in the present study for both species across Britain and continental Europe. A study of *Quercus pyrenaica* [71] mapped clonal groups, each occupying many tens of square metres, and found high genetic diversity preserved alongside high clonality, indicating seed regeneration as well as vegetative reproduction: this parallels a study of *C. sativa* in Greece [72].

Assessment of *Populus tremula* clones [69] revealed that a very long period (>2000 years) of vegetative regeneration from suckers was accompanied by somatic mutations and a reduction of pollen viability. This is comparable with some ancient *C. sativa* trees and stools in the present study, which were evidently single generation plants >500 years old with somatic mutations recorded in some of their vegetatively reproduced components. For some of these sites there may have been only a few *C. sativa* trees, even just one during hundreds of years, many kilometres distant from the next. The pollen of *C. sativa* is generally locally dispersed [73, 74], although it can be wind-blown long distances [2], so genetic isolation might be anticipated for such remote trees. Interestingly, the only small-island population of *C. sativa* sampled within Britain and Ireland in this study (from the Isle of Wight) reported the highest inbreeding coefficient (F_{IS} 0.215).

The European *Castanea sativa* genepool

The analysis of the continental European, England, Ireland and Wales dataset indicated an overall genepool with high diversity, moderate differentiation and low degree of isolation and inbreeding, in which the England, Ireland and Wales sites were amongst the most diverse.

AMOVA results were different from, but comparable with, those from the previous analysis [11] of the European genepool, presumably reflecting the expanded contribution from British and Irish samples.

Two main 'western' European genepools were identified: England, Ireland and Wales samples fitted predominantly into 'Genepool C', including the 'oldest' ancient woodlands and iconic ancient trees; whereas 'Genepool D' contained <25% of the England, Ireland and Wales samples, but principally from Wales and historic garden sites.

The assignment by STRUCTURE of sites to 'Genepool C' or 'Genepool D' was corroborated by the other statistical analyses, except that F_{ST} analysis indicated a strong affiliation between N Portugal (PT03 and PT01) and some England, Ireland and Wales sites. Previous research [11] had assessed Portugal, France, England and Italy as a single 'Sub-cluster 2'. In considering the potential source regions for British and Irish *C. sativa* genepools, N Portugal is an important LGM refugial zone [2] that also shared many cultural connections with Britain, from the Atlantic Bronze Age [75] through to the 18th century AD, when Portuguese sweet chestnuts were especially recommended for propagation [28], perhaps providing linkages between British sites and PT01 and PT03.

The map of final outcomes from the various genetic statistical comparisons (Fig 12) illustrates the importance of the Iberian, Pyrenean and Italian LGM refugial areas as potential sources for the British and Irish *C. sativa* genotypes.

Post-LGM European distribution of *Castanea sativa* compared with other tree species

Previous studies [2, 12, 13] evinced that *C. sativa* behaved like other tree species in continental Europe during and post-LGM: it survived in discrete refugia and, following climatic amelioration, spread according to natural and anthropogenic factors [7, 10, 11, 14].

The present study can be compared with phylogenetic studies of other tree species that colonized Britain and Ireland from LGM refugia in continental Europe: *Fraxinus excelsior* [64, 76]; *Quercus robur*/*Q. petraea* [3, 5, 6]; *Tilia cordata* [77]; *Corylus avellana* [4, 78]; and *Fagus sylvatica* [79, 80]. These studies mapped the genotypes/haplotypes of these species across Europe and considered how natural dispersal from LGM refugia and/or anthropogenic translocations contributed to their present distribution: they appear comparable with the spread of *Castanea sativa* from continental Europe to Britain and Ireland.

Studies of *Fraxinus excelsior* in Europe (including Britain) [66, 76] described 12 haplotypes covering Europe: only one was found in Britain and Ireland, which occurs elsewhere only in Iberia, so was concluded to have spread from Iberian LGM refugia to Britain and Ireland along the Atlantic seaboard. AMOVA and STRUCTURE statistics described a single, genetically diverse population covering most of Britain; F_{ST} values between Britain and France were very low. This work on *F. excelsior* closely parallels the results of the present study for *C. sativa*.

Corylus avellana genotypes have been mapped across western Europe [4, 78], indicating a rapid post-LGM expansion from a refugium in the Biscay area into western Europe, Britain and Ireland, presumed to have been facilitated by nut-eating birds and mammals (jays, nut-hatches, squirrels) with repeated long-distance dispersal of nuts. Genetic, historical and archaeological data indicated that hazelnuts were a food source for early people and *C. avellana* may have been deliberately or accidentally spread [4], potentially similar to *Castanea sativa*.

Studies of *Quercus robur* and *Q. petraea* in Britain and Ireland [3, 5, 6] indicated three predominant haplotypes (see Fig 1 and Fig 2 in [5]) that originated in a LGM refugium in west Iberia and migrated north and west into Britain and Ireland along the Atlantic seaboard. These studies considered the rates of post-LGM migration necessary before Irish Sea and

English Channel land bridges flooded and stated that long-distance dispersal was required, by 'birds and rare climatic storms' [6]. The role of anthropogenic translocation was also considered (see Fig 1A and Fig 1B in [3]): two routes were proposed—the 'Atlantic lineage' spread from Spain to Scandinavia; and the 'Balkan lineage' spread from east to west Europe, north of the Alps. Post-LGM migration routes for hominids and for oak appeared very similar, such that anthropogenic translocation might explain 'islands' of oak genotypes where one haplotype has evidently 'leapfrogged' another, as found in France and in Britain [3]. Acorns were an important food for humans and their livestock and varietal selection and propagation appear to have influenced oak genotype distribution [3]: parallels with *C. sativa* are evident.

The distribution of *Fagus sylvatica* across Europe has been assessed using palaeobotanical and genetic evidence [79, 80]: post-LGM colonization was from mountain refugia in N Spain and SW France to western continental Europe; and from refugia in E Alps-Slovenia-Istria, skirting north of the Alps to France and thence to England. *F. sylvatica* genotypes in Britain [80] show a broad admixture, with regional differentiation inherited from several continental European sources: patterns formed by natural colonization were found to persist despite anthropogenic interventions. *F. sylvatica* nuts were an important food for humans and their livestock, as were *C. sativa* nuts [81].

A study of LGM refugia in the Pyrenees and northwest Iberia [82] indicated that many tree species (including *C. sativa*) survived there and that from 6000 yr BP deciduous broadleaved trees spread to colonize northern Iberia: refugia for *C. sativa* have been corroborated on the Cantabrian coast and in S Galicia and N Portugal [8, 12]. These Pyrenean and Iberian refugia appear to be the predominant regional sources of the *C. sativa* genotypes found by the present study in England, Ireland and Wales.

The expansion of *C. sativa* from LGM refugia in Iberia and S and W France [2, 13], evidently has similarities with the expansion from refugia of *Q. robur/Q. petraea*, *F. sylvatica*, *F. excelsior* and *C. avellana*. The ecological niche of *C. sativa* is broadly similar to *Q. robur/Q. petraea*, *F. sylvatica* and *C. avellana*, in terms of soil and climate preferences and seed (nut) dispersal. Crucially, *C. sativa* has long been favoured by humans for food, and anthropogenic interventions in its dispersal and varietal selection have synergised with natural processes [14], possibly paralleling the natural/anthropogenic spread of *Q. robur/Q. petraea*, *F. sylvatica* and *C. avellana*. However, there is no evidence for natural spread of *C. sativa* to Britain and Ireland.

Conclusions

Analysis of the genetic diversity of *Castanea sativa* in England, Ireland and Wales revealed a single overall genepool, with two sub-clusters: Wales sites, differentiated from Ireland. The England, Ireland and Wales sites had their strongest European connections with sites in south and west France, northern Iberia, central Italy, and a site in Romania: no connections were found with the eastern European genepools in Greece, the Balkans and Turkey. The relatively high diversity of the England, Ireland and Wales *C. sativa* genepool is considered the product of several arrivals of seed and/or living plant material from these European zones. Sites in Britain and Ireland with ancient trees and coppice stools (Types C and D) were genetically differentiated from sites with relatively recent (<200 years age) trees and stools (Types A and B), indicating 'modern' selection of seed/rooted plants from new sources and not from already established stands. Sites in Britain with ancient historic garden trees (Type E) were strongly affiliated with sites in N Portugal and NW Spain.

Clonal analysis of British and Irish samples determined for the first time the 'genotypic size' of ancient trees and stools and thereby their great antiquity, including the largest *C. sativa* stools and ancient trees recorded in Britain. Clonal evidence revealed both natural

regeneration and anthropogenic propagation of trees and coppice, through bough collapse, layering and planting of vegetative material. Historical systematic nut production was evinced for several sites, where extant ancient trees renowned for their eating nut qualities indicate past selection for nuts, including two examples of ancient trees with grafted boughs.

The ancient *C. sativa* trees and coppice stools in Britain are considered single-generation survivors of the original planted or self-sown trees. Genetic stability of *C. sativa* was evinced from the clonal studies, sustained through repeated cutting, demise and regrowth over many centuries: a few somatic mutations were recorded.

The conservation significance of ancient *C. sativa* trees and stools in Britain and Ireland has been highlighted by this study: their genetic diversity is important to considerations of future risks from pathogens and environmental change. The Future Trees Trust *Castanea sativa* collection was shown to consist of a broad genetic base representative of the overall British and Irish genepool.

Supporting information

S1 File. Sample locations England, Ireland and Wales .kml file.
(KML)

S2 File. ‘Genepool A’ and ‘Genepool B’ sample locations England, Ireland and Wales .kml file.
(KML)

S3 File. Results of five parameters analyses England, Ireland and Wales.
(DOCX)

S4 File. STRUCTURE analysis European including England, Ireland and Wales samples.
(DOCX)

S5 File. ‘Genepool C’ and ‘Genepool D’ sample locations Europe including England, Ireland and Wales .kml file.
(KML)

S1 Fig. Dendrogram of England, Ireland and Wales samples using Nei genetic distance.
(PDF)

S1 Table. Database for England, Ireland and Wales samples with site information.
(XLSX)

S2 Table. AMOVA England, Ireland and Wales samples.
(DOCX)

S3 Table. Results of F_{ST} analysis for England, Ireland and Wales samples and site parameter distributions.
(XLSX)

S4 Table. Results of r relatedness analysis for England, Ireland and Wales samples using LRM and QGM.
(XLSX)

S5 Table. LRM and QGM results for ‘Castiard’ study site.
(XLSX)

S6 Table. Database European including England, Ireland and Wales samples.
(XLSX)

S7 Table. Genetic diversity analysis European including England, Ireland and Wales samples.

(DOCX)

S8 Table. Results of F_{IS} analysis of European including England, Ireland and Wales samples.

(XLSX)

S9 Table. Results of F_{ST} analysis of European including England, Ireland and Wales samples.

(XLSX)

S10 Table. F_{ST} threshold values.

(DOCX)

S11 Table. Results of r relatedness analysis for European including England, Ireland and Wales samples using LRM and QGM.

(XLSX)

S12 Table. Results of clonal analysis of European including England, Ireland and Wales samples.

(XLSX)

Acknowledgments

We thank owners in Britain and Ireland for access: National Trust, Woodland Trust, Forest Enterprise England, Coillte, Crown Estates, Basildon Council, Bertholey, Bigsweir, Brompton Bryan, Bushy House, Canford School, Cappelquin, Celtic Manor Estate, Chestnut Street, Chillingham Castle, Cowdray, Dean Heritage Museum Trust, Duchy of Cornwall, East Malling Research, Felbridge Parish Council, Fredville, Glen Upper, Godinton, Goodwood, Hagley Hall, Halswell House, Howletts, Kateshill House (Bewdley), Kentchurch, Littledean Hall, Llanfihangel Crucorney, Lesnes Abbey (London Borough of Bexley), Luton Hoo Hotel, Lydney Park, Margam, Nettlecombe, Neuadd Farm Cwmdu, Penshurst Place, Popes Hill, Powderham, St. Pierre Golf Club (Chepstow), Stoke Edith, Southend on Sea Borough Council, Tortworth, Wyndley Bury.

Information and practical assistance were provided by David Alderman, Graham Bathe, Clive Belsom, Simon Bonvoisin, Pete Byfield, David Bullock, Jill Butler, Mike Calnan, Iain Carter, Jeremy Clarke, Brian Clifford, Roger Clooney, Marco Conedera, Charles Courtenay, Rupert Foley, Llorenç Picornell Gelabert, Diane Gilbert, Ted Green, Ray Hawes, Zöe Hazell, Katherine Hearn, Ian Holt, Della Hooke, Ted Horgan, Lucy Hughes, Brian Jones, Roy Keeler, Patrik Krebs, Josefa Fernandez-Lopez, Santiago Pereira-Lorenzo, David McOmish, Hugh Milner, Yannick Miras, Andy Moir, Robert Moreton, Brian Muelaner, John Leigh-Pemberton, George Peterken, Oliver Rackham R.I.P., Karen Russell, Paolo Squatriti, Ian Standing, Andrés Teira-Brión, Filipe Vaz, Robyn Veal, Charles Watkins, Elisabeth Whittle, Pat Wolseley; members of the Kent Coppice Workers Co-operative; and members of the Future Trees Trust Sweet Chestnut Group.

Joan Cottrell advised on methods and interpretation. Santiago Pereira-Lorenzo provided DNA reference material.

Author Contributions

Conceptualization: Rob Jarman, Karen Russell, Frank M. Chambers.

Data curation: Rob Jarman, Claudia Mattioni, Karen Russell, M. Angela Martin, Marcello Cherubini.

Formal analysis: Rob Jarman, Claudia Mattioni, Karen Russell, M. Angela Martin, Marcello Cherubini.

Funding acquisition: Rob Jarman, Karen Russell, Frank M. Chambers, Debbie Bartlett.

Investigation: Rob Jarman, Claudia Mattioni, Karen Russell, Frank M. Chambers, Debbie Bartlett.

Methodology: Rob Jarman, Claudia Mattioni, Karen Russell, Frank M. Chambers, M. Angela Martin, Marcello Cherubini, Fiorella Villani.

Project administration: Rob Jarman, Karen Russell, Debbie Bartlett.

Resources: Rob Jarman, Claudia Mattioni.

Software: Rob Jarman, Claudia Mattioni, M. Angela Martin, Marcello Cherubini.

Supervision: Rob Jarman, Karen Russell, Frank M. Chambers, Fiorella Villani, Julia Webb.

Validation: Rob Jarman, Claudia Mattioni, Karen Russell, Frank M. Chambers, Debbie Bartlett, M. Angela Martin, Marcello Cherubini, Fiorella Villani, Julia Webb.

Visualization: Rob Jarman.

Writing – original draft: Rob Jarman.

Writing – review & editing: Rob Jarman, Claudia Mattioni, Karen Russell, Frank M. Chambers, Debbie Bartlett, M. Angela Martin, Fiorella Villani, Julia Webb.

References

1. Jarman R, Hazell Z, Campbell G, Webb J, Chambers FM. Sweet chestnut (*Castanea sativa* Mill.) in Britain: re-assessment of its status as a Roman archaeophyte. *Britannia*. 2019; 50: 1–26. <https://doi.org/10.1017/S0068113X19000011>
2. Krebs P, Pezzatti GB, Beffa G, Tinner W, Conedera M. Revising the sweet chestnut (*Castanea sativa* Mill.) refugia history of the last glacial period with extended pollen and macrofossil evidence. *Quat Sci Rev*. 2019; 206: 111–128. <https://doi.org/10.1016/j.quascirev.2019.01.002>
3. Kremer A. Did early human populations in Europe facilitate the dispersion of oaks? *Internat Oaks*. 2015; 26: 19–28.
4. Brown JA, Beatty GE, Montgomery WI, Provan J. Broad-scale genetic homogeneity in natural populations of common hazel (*Corylus avellana*) in Ireland. *Tree Genet Genomes*. 2016; 12: 122. <https://doi.org/10.1007/s11295-016-1079-7>
5. Petit R, Brewer S, Bordács S, Burg K, Cheddadi R, Coart E, et al. Identification of refugia and post-glacial colonisation routes of European white oaks based on chloroplast DNA and fossil pollen evidence. *For Ecol Manage*. 2002; 156: 49–74.
6. Lowe A, Unsworth C, Gerber S, Davies S, Munro R, Kelleher C, et al. Route, speed, and mode of oak postglacial colonisation across the British Isles: integrating molecular ecology, palaeoecology and modelling approaches. *Bot J Scot*. 2005; 57(1+2): 59–81.
7. Huntley B, Birks HJB. *An Atlas of Past and Present Pollen Maps for Europe: 0–13,000 years ago*. Cambridge: Cambridge University Press; 1983.
8. Fernandez-Cruz J, Fernandez-Lopez J. Morphological, molecular and statistical tools to identify *Castanea* species and their hybrids. *Conserv Genet*. 2012; 13: 1589–1600.
9. Martin MA, Mattioni C, Cherubini M, Taurchini D, Villani F. Genetic diversity in European chestnut populations by means of genomic and genic microsatellite markers. *Tree Genet Genomes*. 2010; 6(5): 735–744.
10. Lusini I, Velichkov I, Pollegioni P, Chiocchini F, Hinkov G, Zlatanov T, et al. Estimating the genetic diversity and spatial structure of Bulgarian *Castanea sativa* populations by SSRs: implication for conservation. *Conserv Genet*. 2014; 15: 283–293.

11. Mattioni C, Martin MA, Chiocchini F, Cherubini M, Gaudet M, Pollegioni P, et al. Landscape genetics structure of European sweet chestnut (*Castanea sativa* Mill.): indications for conservation priorities. *Tree Genet Genomes*. 2017; 13: 1–14.
12. López-Sáez JA, Glais A, Robles-López S, Alba-Sánchez F, Pérez-Díaz S, Abel-Schaad D, et al. Unraveling the naturalness of sweet chestnut forests (*Castanea sativa* Mill.) in central Spain. *Veg Hist Archaeobot*. 2017; 26: 167–182. <https://doi.org/10.1007/s00334-016-0575-x>
13. Roces-Díaz JV, Jiménez-Alfaro B, Chytrý M, Díaz-Varela ER, Álvarez-Álvarez P. Glacial refugia and mid-Holocene expansion delineate the current distribution of *Castanea sativa* in Europe. *Palaeogeogr Palaeoclimatol Palaeoecol*. 2018; 491: 152–160.
14. Conedera M, Krebs P, Tinner W, Pradella M, Torriani D. The cultivation of *Castanea sativa* (Mill.) in Europe, from its origin to its diffusion on a continental scale. *Veg Hist Archaeobot*. 2004; 13: 161–179.
15. Pereira-Lorenzo S, Ballester A, Corredoira E, Vieitez AM, Anagnostakis S, Costa R, et al. Chestnut. In: Badenes M, Byrne D, editors. *Fruit Breeding*. Springer; 2012. pp. 729–769.
16. Squatriti P. *Landscape and change in early medieval Italy: chestnut, economy and culture*. 1st ed. Cambridge: Cambridge University Press; 2013.
17. Ledger PM, Miras Y, Poux M, Milcent PY. The palaeoenvironmental impact of prehistoric settlement and proto-historic urbanism: tracing the emergence of the oppidum of Corent, Auvergne, France. *PLoS ONE* 2015; 10(4): e0121517. <https://doi.org/10.1371/journal.pone.0121517> PMID: 25853251
18. Conedera M, Tinner W, Krebs P, de Rigo D, Caudullo G. *Castanea sativa* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayán J, de Rigo D, Caudullo G, Houston Durrant T, Mauri A, editors. *European Atlas of Forest Tree Species*. Publ. Off. EU, Luxembourg, pp. e0125e0+; 2016.
19. Godwin H. *The History of the British Flora*. 2nd ed. Cambridge: Cambridge University Press; 1975.
20. Preston CD, Pearman DA, Hall AR. Archaeophytes in Britain. *Bot J Linn Soc*. 2004; 145: 257–294. <https://doi.org/10.1111/j.1095-8339.2004.00284.x>
21. Rackham O. *Woodlands*. London: Harper Collins, New Naturalist; 2006.
22. Stace CA, Crawley MJ. *Alien plants*. London: Harper Collins; 2015.
23. Rackham O. *Ancient Woodland*. 2nd ed. Dalbeattie: Castlepoint Press; 2003.
24. Botanical Society of Britain & Ireland. BSBI Distribution Database. Accessed 01 March 2019. Available from: <https://bsbi.org/maps?taxonid=2cd4p9h.yx>
25. Buckley P, Howell R. *The Ecological Impact of Sweet Chestnut coppice silviculture on former ancient broadleaved woodland sites in South-east England*. Research Report 627. Peterborough: English Nature; 2004.
26. NFI preliminary estimates of quantities of broadleaved species in British woodlands. National Forest Inventory Report. Edinburgh: Forestry Commission. 2013.
27. Ancient tree inventory. Grantham, UK: Woodland Trust. 2017. Available from: <http://www.ancient-tree-hunt.org.uk/discoveries/TreeSearch>
28. Evelyn J. Silva, or A Discourse of Forest-Trees and the Propagation of Timber in His Majesty's Dominions. 4th ed. London: Royal Society; 1706.
29. TNA: C53/76 (Ch. R. 18 Edward 1) m.10 (in an inspeximus, 1 July 1290), The National Archives, London.
30. Jarman R, Moir AK, Webb JC, Chambers FM and Russell K. Dendrochronological assessment of British veteran sweet chestnut (*Castanea sativa*) trees: successful cross-matching, and cross-dating with British and French oak (*Quercus*) chronologies. *Dendrochronologia*. 2018; 51: 10–21. <https://doi.org/10.1016/j.dendro.2018.07.001>
31. O'Sullivan Beare P. *The Natural History of Ireland 1626*. Translated and edited by O'Sullivan DC. Cork: Cork University Press; 2009.
32. Forbes AC. Tree Planting in Ireland during Four Centuries. *Proc Roy Irish Academy. Section C: Archaeology, Celtic Studies, History, Linguistics, Literature*. 1932; 41: 168–199. Available from: www.jstor.org/stable/25515967.
33. Register Tree. Tree Council of Ireland. 2018. Available from: <http://treecouncil.ie/treeregisterofireland/aaspecies.htm>.
34. Buck EJ. Genetic variation of *Castanea sativa* Mill. Unpub. PhD thesis. University of Wales, Bangor. 2006.
35. Buck EJ, Hadonou M, James CJ, Blakesley D, Russell K. Isolation and characterization of polymorphic microsatellites in European chestnut (*Castanea sativa* Mill.). *Mol Ecol Notes*. 2003; 3: 239–241.
36. Marinoni D, Akkak A, Bounous G, Edwards KJ, Botta R. Development and characterization of microsatellite markers in *Castanea sativa* (Mill.). *Mol Breed*. 2003; 11: 127–136.

37. Barreneche T, Casasoli M, Russell K, Akkak A, Meddour H, Plomion C, et al. Comparative mapping between *Quercus* and *Castanea* using simple-sequence repeats (SSRs). *Theor Appl Genet*. 2004; 108: 558–566. <https://doi.org/10.1007/s00122-003-1462-2> PMID: 14564395
38. Mattioni C, Cherubini M, Micheli E, Villani F, Bucci G. Role of domestication in shaping *Castanea sativa* genetic variation in Europe. *Tree Genet Genomes*. 2008; 4: 563–574.
39. Future Trees Trust. Available from: <http://www.futuretrees.org/our-work/sweet-chestnut/current-research/>.
40. Jarman R, Kofman PD. Coppice in Brief. COST Action FP1301 Reports. Freiburg: Albert Ludwig University of Freiburg. 2017. <https://doi.org/10.13140/RG.2.2.32665.42080>
41. Chapuis MP, Estoup A. Microsatellite null alleles and estimation of population differentiation. *Mol Biol Evol*. 2007; 24: 621–631. FreeNA software. Available from: https://www1.montpellier.inra.fr/CBGP/software/FreeNA/README_FreeNA.pdf <https://doi.org/10.1093/molbev/msl191> PMID: 17150975
42. Kalinowski ST. HP-Rare: a computer program for performing rarefaction on measures of allelic diversity. *Mol Ecol Notes*. 2005; 5: 187–189
43. Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics*. 2012; 28: 2537–2539. <https://doi.org/10.1093/bioinformatics/bts460> PMID: 22820204
44. Nei M. Analysis of gene diversity in subdivided populations. *P Natl Acad Sci USA*. 1973; 70: 3321–3323.
45. Rousset F. Genepop on the web. Available from: <https://kimura.univ-montp2.fr/~rousset/Genepop.htm>.
46. Perrier X, Jacquemoud-Collet JP. DARwin software. 2006. Available from: <http://darwin.cirad.fr/>.
47. Queller DC, Goodnight KF. Estimating relatedness using genetic markers. *Evolution*. 1989; 43: 258–275. <https://doi.org/10.1111/j.1558-5646.1989.tb04226.x> PMID: 28568555
48. Lynch M, Ritland K. Estimation of pairwise relatedness with molecular markers. *Genetics*. 1999; 152: 1753–1766. PMID: 10430599
49. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. *Genetics*. 2000; 155: 945–959. PMID: 10835412
50. Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol*. 2005; 14: 2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x> PMID: 15969739
51. Earl DA, von Holdt BM. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet Resour*. 2012; 4: 359–361.
52. Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I. CLUMPAK: a program for identifying clustering modes and packaging population structure inferences across K. *Mol Ecol Resour*. 2015; 15(5): 1179–1191. <https://doi.org/10.1111/1755-0998.12387> PMID: 25684545
53. Sneath PHA, Sokal RR. Numerical Taxonomy. San Francisco: Freeman. 1973.
54. Takezaki N, Nei M, Tamura K. POPTREE2: software for constructing population trees from allele frequency data and computing other population statistics with windows interface. *Mol Biol Evol*. 2010; 27: 747–752. <https://doi.org/10.1093/molbev/msp312> PMID: 20022889
55. Rambaut A. FigTree, version 1.4.3. Computer program distributed by the author 04–10–2016. Available from: <http://tree.bio.ed.ac.uk/software/figtree/>. Cited 01 June 2018.
56. Commission Forestry. Regions of Provenance and native seed zones. Available from: <https://www.forestry.gov.uk/forestry/inf-d-72kldl>. Cited 02 May 2018.
57. Rippon S, Smart C, Pears B, Fleming F. The Fields of Britannia: continuity and discontinuity in the Pays and Regions of Roman Britain. *Landscapes*. 2013; 14(1): 33–53.
58. Beccaro GL, Torello-Marini D, Binelli G, Donno D, Boccacci P, Botta R, et al. (2012). Insights in the chestnut genetic diversity in Canton Ticino (Southern Switzerland). *Silvae Genetica*. 2012; 61(6): 292–300. <https://doi.org/10.1515/sg-2012-0037>
59. Ramos-Cabrer AM, Caruncho-Picos L, Díaz-Hernández MB, Ciordia-Ara M, Rios-Mesa D, González-Díaz J, et al. Study of Spanish chestnut cultivars using SSR markers. *Adv Hort Sci*. 2006; 20: 113–16.
60. Poljak I, Idžojtić M1, Šatović Z, Ježić M, Čurković-Perica M, Simovski B, et al. Genetic diversity of the sweet chestnut (*Castanea sativa*) in Central Europe and the western part of the Balkan Peninsula and evidence of marron genotype introgression into wild populations. *Tree Genet Genomes*. 2017; 13: 1–13.
61. Konovalov D, Heg D. A maximum-likelihood relatedness estimator allowing for negative relatedness values. *Mol Ecol Resour*. 2008; 8(2): 256–263. <https://doi.org/10.1111/j.1471-8286.2007.01940.x> PMID: 21585767

62. Miller AC, Woeste K, Anagnostakis SL, Jacobs DF. Exploration of a rare population of Chinese chestnut in North America: stand dynamics, health and genetic relationships. *AoB PLANTS*. 2014; 6: plu065; <https://doi.org/10.1093/aobpla/plu065> PMID: 25336337
63. Wang J. Estimating pairwise relatedness in a small sample of individuals. *Heredity*. 2017; 119: 302–313. <https://doi.org/10.1038/hdy.2017.52> PMID: 28853716
64. Sutherland BG, Belaj A, Nier S, Cottrell J, Vaughan SP, Hubert J, et al. Molecular biodiversity and population structure in common ash (*Fraxinus excelsior* L.) in Britain: implications for conservation. *Mol Ecol*. 2010; 19: 2196–2211. <https://doi.org/10.1111/j.1365-294X.2009.04376.x> PMID: 20465580
65. Harmer R. Management of coppice stools. RIN 259. Farnham: Forestry Authority. 1995.
66. Pereira-Lorenzo S, Ramos-Cabrer AM, Barreneche T, Mattioni C, Villani F, Diaz-Hernandez B, et al. Instant domestication process of European chestnut cultivars. *Ann Appl Biol*. 2019; 174: 74–85. <https://doi.org/10.1111/aab.12474>
67. Diez CM, Trujillo I, Martinez-Urdiroz N, Barranco D, Rallo L, Gaut BS. Olive domestication and diversification in the Mediterranean Basin. *New Phytol*. 2015; 206: 436–447. <https://doi.org/10.1111/nph.13181> PMID: 25420413
68. Vaughan SP, Cottrell J, Moodley DJ, Connolly T, Russell K. Clonal structure and recruitment in British wild cherry (*Prunus avium*). *For Ecol Manage*. 2007; 242: 419–430.
69. Ally D, Ritland K, Otto SP. Aging in a long-lived clonal tree. *PLoS Biology*. 2010; 8:e1000454.
70. Pigott D. Lime-trees and basswoods. Cambridge: CUP. 2012.
71. Valbuena-Carabaña M, Gil L. Centenary coppicing maintains high levels of genetic diversity in a root resprouting oak (*Quercus pyrenaica* Willd.). *Tree Genet Genomes*. 2017; 13: 28. <https://doi.org/10.1007/s11295-017-1105-4>
72. Aravanopoulos FA, Drouzas AD, Paraskevi GA. Electrophoretic and quantitative variation in chestnut (*Castanea sativa* Mill.) in Hellenic populations in old-growth natural and coppice stands. *For Snow Landsc Res*. 2001; 76(3): 429–434.
73. Bounous G, Marinoni D. Chestnut: botany, horticulture, and utilization. *Hort Rev*. 2005; 31: 291–347.
74. Peeters AG, Zoller H. Long range transport of *Castanea sativa* pollen. *Grana*. 1988; 27(3): 203–207. <https://doi.org/10.1080/00173138809428927>
75. Cunliffe B. A race apart: insularity and connectivity. *Proc Prehist Soc*. 2009; 75: 55–64.
76. Heuertz M, Fineschi S, Anzidei M, Pastorelli R, Salvini D, Paule L, et al. Chloroplast DNA variation and postglacial recolonization of common ash (*Fraxinus excelsior* L.) in Europe. *Mol Ecol*. 2004; 13: 3437–3452. <https://doi.org/10.1111/j.1365-294X.2004.02333.x> PMID: 15488002
77. Fineschi S, Salvini D, Turchini D, Carnevale S, Vendramin GG. Chloroplast DNA variation of *Tilia cordata* Tiliaceae. *Can J For Res*. 2003; 33: 2503–2508.
78. Boccacci P, Botta R. Investigating the origin of hazelnut (*Corylus avellana* L.) cultivars using chloroplast microsatellites. *Genet Resour Crop Evol*. 2009; 56: 851–859. <https://doi.org/10.1007/s10722-009-9406-6>
79. Magri D, Vendramin GG, Comps B, Dupanloup I, Geburek T, Gomory D, et al. A new scenario for the Quaternary history of European beech populations: palaeobotanical evidence and genetic consequences. *New Phytol*. 2006; 171: 199–221. <https://doi.org/10.1111/j.1469-8137.2006.01740.x> PMID: 16771995
80. Sjolund MJ, Gonzalez-Diaz P, Moreno-Villena JJ, Jump AS. Understanding the legacy of widespread population translocations on the post-glacial genetic structure of the European beech, *Fagus sylvatica* L. *J Biogeogr*. 2017; 1–13. <https://doi.org/10.1111/jbi.13053>
81. Packham JR, Thomas PA, Atkinson MD, Degen T. Biological Flora of the British Isles: *Fagus sylvatica*. *J Ecol*. 2012; 100: 1557–1608. <https://doi.org/10.1111/j.1365-2745.2012.02017.x>
82. Benito Garzon M, Sanchez de Dios R, Sainz Ollero H. Predictive modelling of tree species distributions on the Iberian Peninsula during the Last Glacial Maximum and Mid-Holocene. *Ecography*. 2007; 30: 120–134. <https://doi.org/10.1111/j.2006.0906-7590.04813.x>