



This is a peer-reviewed, final published version of the following document and is licensed under Creative Commons: Attribution 4.0 license:

Staddon, Philip L. ORCID logoORCID: <https://orcid.org/0000-0002-7968-3179> and Faghihinia, Maede (2021) Grazing intensity is key to global grassland carbon sequestration potential. Sustainable Environment, 7 (1). Art 1895474. doi:10.1080/27658511.2021.1895474

Official URL: <http://dx.doi.org/10.1080/27658511.2021.1895474>

DOI: <http://dx.doi.org/10.1080/27658511.2021.1895474>

EPrint URI: <https://eprints.glos.ac.uk/id/eprint/9606>

Disclaimer


The University of Gloucestershire has obtained warranties from all depositors as to their title in the material deposited and as to their right to deposit such material.

The University of Gloucestershire makes no representation or warranties of commercial utility, title, or fitness for a particular purpose or any other warranty, express or implied in respect of any material deposited.

The University of Gloucestershire makes no representation that the use of the materials will not infringe any patent, copyright, trademark or other property or proprietary rights.

The University of Gloucestershire accepts no liability for any infringement of intellectual property rights in any material deposited but will remove such material from public view pending investigation in the event of an allegation of any such infringement.

PLEASE SCROLL DOWN FOR TEXT.



Sustainable Environment

An international journal of environmental health and sustainability

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/oaes21>

Grazing intensity is key to global grassland carbon sequestration potential

Philip L Staddon & Maede Faghihinia

To cite this article: Philip L Staddon & Maede Faghihinia (2021) Grazing intensity is key to global grassland carbon sequestration potential, Sustainable Environment, 7:1, 1895474, DOI: [10.1080/27658511.2021.1895474](https://doi.org/10.1080/27658511.2021.1895474)

To link to this article: <https://doi.org/10.1080/27658511.2021.1895474>



© 2021 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.



Published online: 11 Apr 2021.



Submit your article to this journal [↗](#)



Article views: 42



View related articles [↗](#)

Grazing intensity is key to global grassland carbon sequestration potential

Philip L Staddon^{a,b} and Maede Faghihinia^c

^aCountryside and Community Research Institute, University of Gloucestershire, Cheltenham GL50 4AZ, UK; ^bSchool of Agriculture, Food and Environment, Royal Agricultural University, Cirencester GL7 6JS, UK; ^cLaboratory of Fungal Biology, Institute of Microbiology, Czech Academy of Sciences, Vídeňská 1083, Krč, 14220 Prague 4, Czech Republic

ABSTRACT

Grasslands are coming under ever-increasing pressure worldwide. Many grasslands are degraded due to overgrazing and inappropriate land management. This is impacting belowground biology and soil biological processes. One aspect that deserves far greater attention is the intensity of grazing and how this impacts grassland soil ecosystems. Grazing intensity impacts soil organisms including their diversity and activity, and the soil carbon cycle. However, environmental characteristics determine in part the effects of grazing intensity on soil processes. In addition, many questions remain to be answered in relation to the type of livestock and grazing regime. Only with a fuller understanding of the impacts of grazing on the soil ecosystem will it be possible to advise farmers and land managers on optimal grazing choices for a sustainable future.

ARTICLE HISTORY

Received 15 December 2020
Accepted 18 February 2021



KEYWORDS

Environmental change;
grazing intensity; livestock
density; soil biology; Soil
carbon

Food security and climate change mitigation are arguably the two most significant challenges facing humanity. Because agriculture accounts for an ever-increasing share of the world's net primary production, the management of agricultural land and soil has implications for the global carbon cycle and feedbacks to climate change. This is true for all agricultural systems be they arable, permanent crops or livestock focused. Land used for livestock grazing accounts for about 60% of total agricultural land (FAO 2020) and includes much of the grassland area worldwide, from arid rangelands, such as steppes and savannas, to wetter temperate grasslands. The demand for meat production from an increasingly affluent and growing human population (Ritchie, 2017) is adding pressure to many grasslands, many of which are now classified as degraded as a result of overuse and especially overgrazing. This degradation of grasslands will ultimately impact the sustainability of their use for meat production whilst also leading to positive feedback on climate change as carbon is both lost from grassland soils and less is being sequestered as a result of sparser vegetation. Despite much research being undertaken on the impacts of grazing on ecosystem function in grasslands, including on belowground processes, there is a dearth of information on how grazing intensity *per se* or livestock density impacts grasslands and their carbon balance. Indeed, much of the research to date compares grazed versus non-grazed grassland, although

more recently there has been a belated focus on this issue of grazing intensity. In the last couple of years, a wealth of studies have highlighted how little we actually know about the effects grazing management choices are having on grassland ecosystem functioning.

As traditional livestock farming approaches disappear in favor of intensification, there is increasing evidence of the benefits of low-level livestock grazing in many systems. In Spain, for example, grazing abandonment in *Quercus dehesa* grassland has led to lower soil carbon content (Oggioni et al., 2020). However, in other systems such as alpine steppe in the Tibetan Plateau, it has been reported that the removal of grazing leads to increased soil carbon content (Liu et al., 2020). Zhan et al. (2020) highlight the lack of understanding of how grazing intensity impacts grassland ecosystem functioning, especially belowground. In a meta-analysis across grasslands in China, classified into low, medium and high grazing intensities, they found that soil microbial activity and soil organic carbon were the highest under low grazing intensity. Similarly, Jiang et al. (2020) found that low-intensity grazing increased soil carbon content in Chinese grasslands, whereas moderate to high grazing intensity decreased it. Going hand in hand with impacts on grassland soil carbon stocks, grazing intensity impacts the nitrogen cycle and availability of nutrients, including micronutrients, in the soil (Hou et al., 2020).

CONTACT Philip L Staddon  pstaddon@glos.ac.uk  Countryside and Community Research Institute, University of Gloucestershire, Cheltenham GL50 4AZ, UK

Reviewing editor: Conor Buggy, School of Public Health, Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland

© 2021 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to: Share — copy and redistribute the material in any medium or format. Adapt — remix, transform, and build upon the material for any purpose, even commercially. The licensor cannot revoke these freedoms as long as you follow the license terms. **Under the following terms:** Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use. **No additional restrictions** You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.




Figure 1. Annual scientific production in terms of paper counts (panel a) and the world's most productive countries (panel b) on the interaction between arbuscular mycorrhiza and livestock. The selection of peer-reviewed publications were retrieved from ISI Web of Science (WoS) on the 10 December 2020. The search strings used are as follows: "arbuscul*" OR "mycorrhiz*" AND "*graz*" OR "defoli*" OR "brows*" OR "clip*" OR "live-stock" OR "livestock" OR "herbivor*" OR "cow" OR "cattle" OR "sheep" OR "goat" OR "ewe" NOT "insect*" NOT "ectomycorrhiz*". This search resulted in a list of 169 documents and then the titles were screened to remove unrelated studies which resulted in a final list of 102 documents.

A key component of soil biota in grasslands is the arbuscular mycorrhizal fungi, which form symbiotic associations with plant roots (Staddon et al., 2002). Mycorrhizal fungi assist the plant with nutrient acquisition and, also act as a direct pathway for movement of carbon from the plant to the soil. As such, they are a central component of the soil carbon cycle and play a significant role in determining soil carbon sequestration (Staddon, 2003). Only relatively recently has much effort been spent investigating the impact of livestock on mycorrhizal fungi (Figure 1), leading to a focus on the impact of grazing intensity on mycorrhizas. Faghihinia et al. (2020b) demonstrated that increasing grazing intensity led to decreasing mycorrhizal hyphal density in the soil, which would have implications for both the nutrient uptake capacity of mycorrhizas and carbon translocation into the bulk soil. Focusing on soil microbial biomass more generally, Toledo et al. (2020) showed that high grazing intensity decreased microbial biomass carbon in three ecological areas in Patagonia. They however noted that the effect of grazing intensity on soil microbial communities is moderated by environmental characteristics of the ecological areas, including seasonality (Toledo et al., 2020). This seasonality effect on the impact of grazing intensity was also reported for mycorrhizal fungi in North China steppe (Faghihinia et al., 2020a).

It is also worth highlighting that with regard to grazing impacts on grasslands, it is not only intensity that has often been overlooked but also the make-up of the grazing livestock involved, namely single or multiple species. Indeed, it has recently been shown that grazing by a mix of cattle and sheep can stimulate net carbon sequestration in grassland when compared to either of the species alone (Chang et al., 2020). Due to the dual

challenges of securing food security (especially protein) and mitigating climate change, much more effort is required to better understand the impacts of grazing intensity and types on grassland ecosystem functioning. Only then will it be possible to advise farmers, land-owners and governments on optimal grazing choices to secure the best long-term sustainable outcomes for ecosystem services under a range of conditions, including those resulting from climate change.

Public Interest Statement

Climate change and sustainability are now important considerations for governments and the public worldwide. The understanding that land needs to be managed more sustainably to secure food and water for future generations, as well as mitigate climate change, is now accepted by the public. In many cases, however, there is a lack of evidence as to what are the most appropriate solutions for maximizing target ecosystem services. How grasslands are managed is a case in point. This short communication offers a perspective on the knowledge gaps with regard to livestock management on grasslands. The optimal grazing density or grazing approach for the sustainable use of different grasslands is unknown. Yet without a sound scientific evidence base, it is not possible to advise land managers on which grazing management choice is the most appropriate for the sustainable provision of ecosystem services.

Author description

Dr Staddon is a principal lecturer in environment and sustainability at the Countryside and Community

Research Institute, University of Gloucestershire. He also teaches environmental science at the Royal Agricultural University and Shandong Agricultural University. He has published widely on the impacts of environmental changes, especially climate change, on terrestrial ecosystems. He has also published on the impacts of climate change on human health. Dr Staddon has held visiting positions at the University of Exeter, Xi'an Jiaotong-Liverpool University, University of Liverpool, and Open University.

Dr Faghihinia completed her PhD in *Environmental Sciences* at the University of Liverpool, on the “*Effects of long-term grazing intensity on arbuscular mycorrhizal fungi and soil carbon fluxes in a steppe ecosystem*”. Her work has significant implications for grassland ecosystem management and carbon cycling. She is about to start a postdoctoral fellow position investigating “*the role of bacteria in organic nutrient exploration by arbuscular mycorrhizal fungi*”, in the Laboratory of Fungal Biology at the Czech Academy of Sciences Institute of Microbiology.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The authors received no direct funding for this research.

References

- Chang, Q., Xu, T., Ding, S., Wang, L., Liu, J., Wang, D., Wang, Y., Li, Z., Zhao, X., Song, X., & Pan, D. (2020). Herbivore assemblage as an important factor modulating grazing effects on ecosystem carbon fluxes in a meadow steppe in Northeast China. *Journal of Geophysical Research: Biogeosciences*, 125, e2020JG005652. <https://doi.org/10.1029/2020JG005652>
- Faghihinia, M., Zou, Y., Bai, Y., Marrs, R., & Staddon, P. L. (2020a). Seasonal variation in the response of arbuscular mycorrhizal fungi to grazing intensity. *Mycorrhiza*, 30(5), 635–646. <https://doi.org/10.1007/s00572-020-00974-8>
- Faghihinia, M., Zou, Y., Chen, Z., Bai, Y., Li, W., Marrs, R., & Staddon, P. L. (2020b). The response of grassland mycorrhizal fungal abundance to a range of long-term grazing intensities. *Rhizosphere*, 13, 100178. <https://doi.org/10.1016/j.rhisph.2019.100178>
- FAO (2020) Land use in agriculture by the numbers. *Food and Agriculture Organization of the United Nations*. <http://www.fao.org/sustainability/news/detail/en/c/1274219/> accessed 17.02.21
- Hou, D., Guo, K., & Liu, C. (2020). Asymmetric effects of grazing intensity on macroelements and microelements in grassland soil and plants in Inner Mongolia grazing alters nutrient dynamics of grasslands. *Ecology and Evolution*, 10 (16), 8916–8926. <https://doi.org/10.1002/ece3.6591>
- Jiang, Z.-Y., Hu, Z.-M., Lai, D. Y. F., Han, D.-R., Wang, M., Liu, M., Zhang, M., & Guo, M.-Y. (2020). Light grazing facilitates carbon accumulation in subsoil in Chinese grasslands: A meta-analysis. *Global Change Biology*, 26(12), 7186–7197. <https://doi.org/10.1111/gcb.15326>
- Liu, Y., Tenzintarchen, G. X., Wei, D., Dai, D., & Xu-Ri. (2020). Grazing exclusion enhanced net ecosystem carbon uptake but decreased plant nutrient content in an alpine steppe. *Catena*, 195, 104799. <https://doi.org/10.1016/j.catena.2020.104799>
- Oggioni, S. D., Ochoa-Hueso, R., & Peco, B. (2020). Livestock grazing abandonment reduces soil microbial activity and carbon storage in a Mediterranean Dehesa. *Applied Soil Ecology*, 153, 103588. <https://doi.org/10.1016/j.apsoil.2020.103588>
- Ritchie, H. (2017) *Meat and dairy production*. Published online at OurWorldInData.org with data sourced from the UN Food and Agricultural Organization. <https://ourworldindata.org/meat-production> accessed 17.02.21 .
- Staddon, P. L. (2003). Rapid turnover of hyphae of mycorrhizal fungi determined by AMS microanalysis of ^{14}C . *Science*, 300(5622), 1138–1140. <https://doi.org/10.1126/science.1084269>
- Staddon, P. L., Heinemeyer, A., & Fitter, A. H. (2002). Mycorrhizas and global environmental change: Research at different scales. *Plant and Soil*, 244(1/2), 253–261. <https://doi.org/10.1023/A:1020285309675>
- Toledo, S., Peri, P. L., Correa, O. S., Gargaglione, V., & Gonzalez-Polo, M. (2020). Soil microbial communities respond to an environmental gradient of grazing intensity in south Patagonia Argentina. *Journal of Arid Environments*, 184, 104300. <https://doi.org/10.1016/j.jaridenv.2020.104318>
- Zhan, T., Zhang, Z., Sun, J., Liu, M., Zhang, X., Peng, F., Tsunekawa, A., Zhou, H., Gou, X., & Fu, S. (2020). Meta-analysis demonstrating that moderate grazing can improve the soil quality across China's grassland ecosystems. *Applied Soil Ecology*, 147, 103438. <https://doi.org/10.1016/j.apsoil.2019.103438>